


30. 10. 2004



**Europäisches
Patentamt**

**European
Patent Office**

**Office européen
des brevets**



Bescheinigung

Certificate

Attestation

Die angehefteten Unterla-
gen stimmen mit der
ursprünglich eingereichten
Fassung der auf dem näch-
sten Blatt bezeichneten
europäischen Patentanmel-
dung überein.

The attached documents
are exact copies of the
European patent application
described on the following
page, as originally filed.

Les documents fixés à
cette attestation sont
conformes à la version
initialement déposée de
la demande de brevet
européen spécifiée à la
page suivante.

Patentanmeldung Nr. Patent application No. Demande de brevet n°

03018266.1

Der Präsident des Europäischen Patentamts;
Im Auftrag

For the President of the European Patent Office

Le Président de l'Office européen des brevets
p.o.

R C van Dijk

**PRIORITY
DOCUMENT**
SUBMITTED OR TRANSMITTED IN
COMPLIANCE WITH RULE 17.1(a) OR (b)

BEST AVAILABLE COPY



Anmeldung Nr:
Application no.: 03018266.1
Demande no:

Anmeldetag:
Date of filing: 11.08.03
Date de dépôt:

Anmelder/Applicant(s)/Demandeur(s):

Kweek-en Researchbedrijf Agrico B.V.
P.O. Box 40
8300 AA Emmeloord
PAYS-BAS

Bezeichnung der Erfindung/Title of the invention/Titre de l'invention:
(Falls die Bezeichnung der Erfindung nicht angegeben ist, siehe Beschreibung.
If no title is shown please refer to the description.
Si aucun titre n'est indiqué se referer à la description.)

Resistant plants and uses thereof

In Anspruch genommene Priorität(en) / Priority(ies) claimed /Priorité(s)
revendiquée(s)
Staat/Tag/Aktenzeichen/State/Date/File no./Pays/Date/Numéro de dépôt:

Internationale Patentklassifikation/International Patent Classification/
Classification internationale des brevets:

C12N15/00

Am Anmeldetag benannte Vertragstaaten/Contracting states designated at date of
filing/Etats contractants désignées lors du dépôt:

AT BE BG CH CY CZ DE DK EE ES FI FR GB GR HU IE IT LU MC NL
PT RO SE SI SK TR LI

Resistant plants and uses thereof

The present invention relates to a novel method for increasing the resistance of a plant, in particular of a Solanaceae, preferably of potato and tomato, to plant pathogens of the phylum Oomyceta comprising increasing the activity of the polypeptid of the present invention. The invention further relates to polynucleotides and vectors comprising these polynucleotides. The invention furthermore relates to corresponding vectors, cells transgenic plants and transgenic propagation material derived from them, methods to produce them and to their use for the production of foodstuffs, feeding stuffs, seed, pharmaceuticals or fine chemicals.

The aim of plant biotechnology work is the generation of plants with advantageous novel properties, for example for increasing agricultural productivity, increasing the quality in the case of foodstuffs, or for producing specific chemicals or pharmaceuticals (Dunwell JM (2000) J Exp Bot 51 Spec No:487-96). The plant's natural defense mechanisms against pathogens are frequently insufficient. Fungal diseases alone result in annual yield losses of many billions of US\$. The introduction of foreign genes from plants, animals or microbial sources can increase the defenses. Examples are the protection of tobacco against feeding damage by insects by expressing *Bacillus thuringiensis* endotoxins under the control of the 35S CaMV promoter (Vaeck et al. (1987) Nature 328:33-37) or the protection of tobacco against fungal infection by expressing a bean chitinase under the control of the CaMV promoter (Broglie et al. (1991) Science 254:1194-1197). However, most of the approaches described only offer resistance to a single pathogen or a narrow spectrum of pathogens.

Despite the notorious Irish potato famine of the mid-19th century, late blight still continues to be one of the most devastating of all diseases in crop plants. Late blight is caused by the oomycete fungus *Phytophthora infestans*, a specialised pathogen, primarily causing disease on the foliage and fruits of a range of Solanaceae species, especially potato and tomato. The fungus was first observed in Mexico and for several reasons Mexico is believed to be the centre of origin of the fungus. Both of the mating types A1 and A2 are permanently present in for example the Toluca area. Also, *P. infestans* is reported on native *Solanum* species in remote areas of Mexico. Furthermore, many species of tuber bearing *Solanum* with a high level of resistance to late blight are found in Mexico. Prevailing measures to prevent crop failures or reduced yields imply the application of fungicides that prevent or cure an infection by *P. infestans*. Instead of the massive use of chemical pesticides an alternative approach for controlling late blight could be advantageous: the use of cultivars, which harbour partial or complete resistance to late blight. To obtain late blight resistance, breeders have in the past focussed on the introgression of dominant R genes from *Solanum demissum*, a wild potato species indigenous to Mexico. Eleven such R genes have been

2

identified, several of which have been mapped to specific loci on the genetic map of potato (reviewed in Gebhardt and Valkonen, 2001) and recently the R1 gene has been cloned. R¹ and R² are located on chromosomes 5 and 4, respectively. R³, R⁶ and R⁷ are located on chromosome 11. Unknown R genes conferring race specific resistance to late blight have also been described in *S. tuberosum* ssp. *Andigena* and *S. berthaultii* and *pennapisecpum*. The resistance induced by these R-genes was (nearly) complete but appeared not to be durable in any case. Because of the high level of resistance and ease of transfer, many cultivars contain *S. demissum* derived resistance. Unfortunately, *S. demissum* derived race specific resistance, although nearly complete, is not durable. Once newly bred potato cultivars were grown on larger scale in commercial fields, new virulences emerged in *P. infestans*, which rendered the pathogen able to overcome the introgressed resistance. More durable field resistance to late blight, often quantitative in nature and presumed to be race non-specific, can be found in several Mexican and Central and South American *Solanum* species. However this type of resistance is difficult to transfer into potato cultivars through crossing and phenotypic selection.

Diploid *S. bulbocastanum* from Mexico and Guatemala is one of the tuber bearing species that is long known for its high levels of resistance to late blight. Unfortunately, classic transfer of resistance from wild *Solanum* species to cultivated potato is frequently prevented due to differences in ploidy and Endosperm Balance Number (EBN). Despite these problems, introgression of the *S. bulbocastanum* resistance trait has been successful. Recently, somatic hybrids of *S. bulbocastanum* and *S. tuberosum* and backcrossed germplasm were found to be highly resistant to late blight, even under extreme disease pressure (Helgeson *et al.*, 1998). Some of these hybrids and backcrossed germplasm were found to be highly resistant to late blight, even under extreme disease pressure. Despite reports of suppression of recombination, resistance in the backcrossed material appeared to be on chromosome 8 within an approximately 6 cM interval between the RFLP markers CP53 and CT64. A CAPS marker derived from the tomato RFLP probe CT88 cosegregated with resistance.

Accordingly, in the recent years the development of plants resistant to pathogens of the phylum Oomyceta forged ahead. However, 40 years of intense and continuous research and breeding efforts with this germplasm has still not resulted in market introduction of resistant cultivars. The prevailing number of genes identified in the recent years confers merely race specific resistance. Further, the achieved resistance was not durable. In addition, the application of crop protectants is widely considered to be a burden for the environment. Thus, in several Western countries, legislation becomes more restrictive and partly prohibitive to the application of specific fungicides, making chemical control of the disease more difficult. Further, chemical control is expensive. Finally, another restriction is the development of resistance by the fungus to specific

fungicides such as metalaxyl, which has been reported from many countries in the world.

- 5 Accordingly, the problem underlying the present invention is to provide novel means and methods for a efficient protection of plants against late blight and related diseases.

The solution of the technical problem is achieved by providing the embodiments characterized in the claims.

- 10 Accordingly, the present invention relates to a method for generating or increasing the resistance of a plant to plant pathogen of the phylum Oomyceta comprising increasing the activity of *Rpi-blb2* protein in the plant or a tissue, organ or cell of the plant or a part thereof.

- 15 *Rpi-blb2* is a LZ-NBS-LRR type of R gene and shows sequence homology to the tomato gene *Mi-1*, that confers resistance to three species of root knot nematodes (*Meloidogyne* spp.) as well as to the potato aphid *Macrosiphum euphorbiae* (Vos *et al.*, 1998; Rossi *et al.*, 1998; Milligan *et al.*, 1998) and to both B- and Q-biotypes of whitefly *Bemisia tabaci* (Nombela *et al.*, 2003). As was found for *Rpi-blb*, *Rpi-blb2* also confers full resistance to a range of *P. infestans* isolates carrying multiple virulence factors and
20 race-specificity has not yet been demonstrated.

By the term "generating" or "increasing" or "stimulating" "the resistance of a plant" is meant that the resistance of a plant or a part thereof is increased in comparison to a reference.

- 25 "Conferring", "existing", "generating", "stimulating" or "increasing" a pathogen resistance means that the defense mechanisms of a specific plant species or variety is increasingly resistant to one or more pathogens due to the use of the method according to the invention in comparison with the wild type of the plant, to which the method according to the invention has not been applied, under otherwise identical conditions
30 (such as, for example, climatic conditions, growing conditions, pathogen species and the like). The increased resistance manifests itself preferably in a reduced manifestation of the disease symptoms, disease symptoms comprising - in addition to the abovementioned adverse effects - for example also the penetration efficiency of a pathogen into the plant or plant cells or the proliferation efficiency in or on the same. In
35 this context, the disease symptoms are preferably reduced by at least 10% or at least 20%, especially preferably by at least 40% or 60%, very especially preferably by at least 70% or 80% and most preferably by at least 90% or 95%. Further by the term "increased" it is hereby meant that the resistance or activity is higher than in a reference or the resistance or activity is generated de novo, if no resistance could be observed in the reference. The stimulation of a resistance or activity is also under the
40 scope of the term "increased resistance" or "increased activity". The activation of the

a gene, i.e. the promoter, can be stimulated, e.g. by applying chemicals or by biotic stress, e.g. by transfection with *P. infestans*, conferring resistance to a pathogen. In the following, the term "increasing" also comprises the term "stimulating".

- 5 "Pathogen resistance" denotes the reduction or weakening of disease symptoms of a plant following infection by a pathogen. The symptoms can be manifold, but preferably encompass those which directly or indirectly have an adverse effect on the quality of the plant, the quantity of the yield, the suitability for use as feeding stuff or foodstuff, or else which make sowing, planting, harvesting or processing of the crop difficult.

10

"Pathogen" within the scope of the invention means by way of example but not by limitation viruses or viroids, bacteria, fungi, animal pests such as, for example, insects or nematodes.

- 15 The term "Rpi-blb2 protein" relates to a protein or polypeptid which expression in a plant or a part confers resistance of the plant or a part of the plant to one of pathogens described herein in comparison to a non-resistant strain.

- The plant or a tissue, organ or cell of the plant or a part thereof comprising increased activity of Rpi-blb2 protein is less susceptible to an infection by a pathogen, in particular to pathogen of the phylum Oomyceta preferably to *P. infestans*, than a plant or a part thereof which has the identical genetic background but not the genetic elements necessary to allow an expression of Rpi-blb2 (herein named as "wild type" or "reference"). Assays for the testing of the resistance of a plant or a part thereof are well known to a person skilled in the art. The resistance to *P. infestans* can be defined as
- 20 Sporulation index according to Flier, 2001. Flier describes the sporulation index as level of sporulation per 1 cm². Thus, a reduction of sporulation per 1 cm² of 20% compared to a wild type is herein defined as resistance. In the examples illustrating the present invention, the sporulation index was defined as level of sporulation per lesion. Thus, by the term "resistance" is alternatively meant a reduction of sporulation per lesion of 20% compared to a wild type. The later definition is preferred. In preferred embodiments the sporulation in an assay is reduced by 30%, more preferred is a reduction of 50%, even more preferred are 70%, even more preferred are more than 80%, more preferred are 85% and 90%. Most preferred is 95% or more.

- 35 Accordingly, in the present invention by "activity" of a Rpi-blb2 protein is meant, that the protein expression confers said reduction in the sporulation index. Further, it was observed, that a typical response for plants containing Rpi-blb2 to a *P. infestans* infection is the presence of small lesions, without any clear sporulation, at the end of the growing season. Thus, in one embodiment, the activity of Rpi-blb2 is defined as the
- 40 presence of small lesions without any clear sporulation.

The term "reference" relates to an organism which is genetically identical to the Organism of the present invention or a part thereof but a expression of Rpi-blb2 can not be observed as there is a little difference in the genom, proteom or metabolom. Therefore
5 the reference is a plant or a part thereof which does not encode Rpi-blb2 or does not transcribe Rpi-blb2 gene or does not translate a Rpi-blb2 mRNA or does not provide any modification for creating an active mRNA or polypeptide. Whether two plants are genetically identical can be tested with assays known to a person skilled in the art, e.g. via fingerprint analysis, e.g. as described in Roldan-Ruiz, Theor. Appl. Genet., 2001,
10 1138-1150. In order to increase the resistance the reference is susceptible to the infection with a plant pathogen of said group.

Preferably, the reference is a clone of that organism in which the polynucleotide of the invention or an activator of the expression of said polynucleotide or a derivate of said polynucleotide or a polypeptide of the present invention or a corresponding vector has
15 been introduced. For example, a preferred reference is the strain before it has been transfected or transformed with the polynucleotid or vector of the invention. If the strain can not be identified it is state of the art to cleave the elements which mediate Rpi-blb2 expression out of the genome of the plant or to inhibit the expression of the Rpi-blb2 protein by other methods. Such a strain can than be compared with the resistant, Rpi-blb2
20 expressing strain.

The term "plant" as used herein refers to all genera and species of higher and lower plants of the Plant Kingdom. The term includes the mature plants, seed, shoots and seedlings and their derived parts, propagation material, plant organs, tissue, proto-
25 plasts, callus and other cultures, for example cell cultures, and any other type of plant cell grouping to give functional or structural units. Mature plants refers to plants at any desired developmental stage beyond that of the seedling. Seedling refers to a young immature plant at an early developmental stage. "Plant" encompasses all annual and perennial monocotyledonous and dicotyledonous plants. Preferred within the scope of
30 the invention are those plants which are employed as foodstuffs or feeding stuffs, very especially preferably monocotyledonous genera and species like the above-described cereal species.

As known to a person skilled in the art, the method of the present invention comprises
35 further selecting those plants in which, as opposed or as compared to the original plant, the resistance to at least one said pathogen exists or is increased.

"Selection" with regard to plants in which - as opposed or as compared to the original plant - resistance to at least one pathogen exists or is increased means all those meth-
40 ods which are suitable for recognizing an existing or increased resistance to pathogens. These may be symptoms of pathogen infection but may also comprise the herein

described symptoms which relate to the quality of the plant, the quantity of the yield, the suitability for use as feeding stuff or foodstuff and the like.

Accordingly, in one embodiment of the method of present invention the Rpi-blb2 protein
5 is encoded by a polynucleotide comprising a nucleic acid molecule selected from the group consisting of:

- a) nucleic acid molecules encoding at least the mature form of the polypeptide depicted in SEQ ID NO: 2 or 4;
- b) nucleic acid molecules comprising the coding sequence as depicted in SEQ ID
10 NO: 1 or 3, or 5 or 6 encoding at least the mature form of the polypeptide;
- c) nucleic acid molecules the nucleotide sequence of which is degenerate as a result of the genetic code to a nucleotide sequence of (a) or (b);
- d) nucleic acid molecules encoding a polypeptide derived from the polypeptide en-
15 coded by a polynucleotide of (a) to (c) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence of the polypeptide encoded by a polynucleotide of (a) to (c);
- e) nucleic acid molecules encoding a polypeptide the sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleic acid molecule of (a) or (b);
- 20 f) nucleic acid molecules comprising a fragment or a epitope-bearing portion of a polypeptide encoded by a nucleic acid molecule of any one of (a) to (e);
- g) nucleic acid molecules comprising a polynucleotide having a sequence of a nucleic acid molecule amplified from a nucleic acid library using the primers as listed in Tab 3b, in particular ARF1F and ARF1R;
- 25 h) nucleic acid molecules encoding a fragment beginning with amino acid: 1, 30, 50, 100, 200, 300, 500, or 1000 and stopping with amino acid 1276, 1000, 500, 300, 200, 50, or 1 of a polypeptide encoded by any one of (a) to (g);
- i) nucleic acid molecules comprising at least 20 nucleotides of a polynucleotide of any one of (a) or (d);
- 30 j) nucleic acid molecules encoding a polypeptide being recognized by a monoclonal antibody that have been raised against a polypeptide encoded by a nucleic acid molecule of any one of (a) to (h);
- k) nucleic acid molecules obtainable by screening an appropriate library under stringent conditions with a probe having the sequence of the nucleic acid molecule of
35 any one of (a) to (j) or of a fragment thereof of at least 15, preferable 30, 60, 100 or more nucleotides; and

Agrico B.V.

20030596

7

- l) nucleic acid molecules the complementary strand of which hybridises under stringent conditions with a nucleic acid molecule of any one of (a) or (k);
or the complementary strand of any one of (a) to (l);
or expressing a polypeptide encoded by a segment or linkage group 6 of *Solanum bulbocastanum* which co-segregates with a marker selected from table 3A and which mediates resistance to pathogens, in particular to pathogens selected from the group consisting of phylum Oomyceta;
and whereby the polynucleotide has not the sequence of Mi1.1 or Mi1.2 as depicted in Seq. ID NO.: 7 and 9 or encoding a Mi1.1 or Mi1.2 protein as depicted in Seq. ID NO.: 8 and 10.

Accordingly, in one embodiment the present invention relates to the method of the present invention, wherein the Rpi-blb2 protein is encoded by the polynucleotide of the present invention.

15

On basis of a BLASTX search the genes with the highest homology identified were the Mi1.1- and Mi1.2-genes. Both genes have a high identity to the sequence depicted in Seq. ID NO.: 1 or 3 or 5 or 6 but do not confer resistance to the plant pathogen of the phylum Oomyceta. Therefore the activity of Mi1.1 and Mi1.2 is an other activity as the activity of the polypeptide of the present invention. The sequence of Mi1.1 and Mi1.2 is herein shown in Seq. ID NO.: 7 to 10. Further, the application EP 401764.4 relates to the Mi-genes. The sequence of prior art Mi1.1- and Mi1.2-genes is excluded from the polynucleotide of the present invention, in particular Seq. ID NO.: 7 and 9 are excluded. Also included may be polynucleotide sequences encoding the polypeptide of Seq. ID NO.: 8 or 10. Thus, in an embodiment also sequences encoding the Mi1.1 and Mi1.2 protein are excluded. Proteins with a lower homology to the polypeptide encoded by the polynucleotide of the present invention are Hero Resistance proteins 1 and 2 (Genbank AccNo.: gi26190252 and gi26190254), Tospovirus resistance proteins A, B, C, D and E [Genbank AccNos.: gi15418709, gi15418710, gi15418712, gi15418713, gi15418714]; R1 [Genbank AccNo.: gi17432423] and Prf [Genbank AccNo.: gi8547237] which sequences or encoded sequences are as well excluded from the sequences of the present invention.

The terms "gene(s)", "polynucleotide", "nucleic acid sequence", "nucleotide sequence" or "nucleic acid molecule(s)" as used herein refers to a polymeric form of nucleotides of any length, either ribonucleotides or deoxyribonucleotides. This term refers only to the primary structure of the molecule.

Thus, this term includes double- and single-stranded DNA, and RNA. It also includes known types of modifications, for example, methylation, "caps" substitution of one

Agrico B.V.

20030596

8

or more of the naturally occurring nucleotides with an analog. Preferably, the DNA sequence of the invention comprises a coding sequence encoding the herein defined polypeptide.

- 5 A "coding sequence" is a nucleotide sequence which is transcribed into mRNA and/or translated into a polypeptide when placed under the control of appropriate regulatory sequences. The boundaries of the coding sequence are determined by a translation start codon at the 5'-terminus and a translation stop codon at the 3'-terminus. A coding sequence can include, but is not limited to mRNA, cDNA, recombinant nucleotide sequences or genomic DNA, while introns may be present as well under certain circumstances.

- 15 By "hybridizing" it is meant that such nucleic acid molecules hybridize under conventional hybridization conditions, preferably under stringent conditions such as described by, e.g., Sambrook (Molecular Cloning; A Laboratory Manual, 2nd Edition, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY (1989)). An example of one such stringent hybridization condition is hybridization at 4XSSC at 65°C, followed by a washing in 0.1XSSC at 65°C for one hour. Alternatively, an exemplary stringent hybridization condition is in 50 % formamide, 4XSSC at 42°C. Further, the conditions during the wash step can be selected from the range of conditions delimited by low-stringency conditions (approximately 2X SSC at 50°C) and high-stringency conditions (approximately 0.2X SSC at 50°C, preferably at 65°C) (20X SSC: 0.3M sodium citrate, 3M NaCl, pH 7.0). In addition, the temperature during the wash step can be raised from low-stringency conditions at room temperature, approximately 22°C, to higher-stringency conditions at approximately 65°C. Both of the parameters salt concentration and temperature can be varied simultaneously, or else one of the two parameters can be kept constant while only the other is varied. Denaturants, for example formamide or SDS, may also be employed during the hybridization. In the presence of 50% formamide, hybridization is preferably effected at 42°C. Some examples of conditions for hybridization and wash step are shown hereinbelow:

- 30 (1) Hybridization conditions can be selected, for example, from the following conditions:
- a) 4X SSC at 65°C,
 - b) 6X SSC at 45°C,
 - 35 c) 6X SSC, 100 mg/ml denatured fragmented fish sperm DNA at 68°C,
 - d) 6X SSC, 0.5% SDS, 100 mg/ml denatured salmon sperm DNA at 68°C,
 - e) 6X SSC, 0.5% SDS, 100 mg/ml denatured fragmented salmon sperm DNA, 50% formamide at 42°C,
 - f) 50% formamide, 4X SSC at 42°C,

Agrico B.V.

20030596

9

- g) 50% (vol/vol) formamide, 0.1% bovine serum albumin, 0.1% Ficoll, 0.1% polyvinylpyrrolidone, 50 mM sodium phosphate buffer pH 6.5, 750 mM NaCl, 75 mM sodium citrate at 42°C,
- h) 2X or 4X SSC at 50°C (low-stringency condition), or
- 5 i) 30 to 40% formamide, 2X or 4X SSC at 42°C (low-stringency condition).
- (2) Wash steps can be selected, for example, from the following conditions:
- a) 0.015 M NaCl/0.0015 M sodium citrate/0.1% SDS at 50°C.
- b) 0.1X SSC at 65°C.
- 10 c) 0.1X SSC, 0.5 % SDS at 68°C.
- d) 0.1X SSC, 0.5% SDS, 50% formamide at 42°C.
- e) 0.2X SSC, 0.1% SDS at 42°C.
- f) 2X SSC at 65°C (low-stringency condition).
- 15 Rpi-blb2 derived from other organisms, may be encoded by other DNA sequences which hybridize to the sequences shown in Seq ID No. 1 or 3 or 5 or 6 under relaxed hybridization conditions and which code on expression for peptides having the activity of Rpi-blb2. Further, some applications have to be performed at low stringency hybridisation conditions, without any consequences for the specificity of the hybridisation. For
- 20 example, a Southern blot analysis of total DNA could be probed with a polynucleotide of the present invention and washed at low stringency (55°C in 2xSSPE, 0.1% SDS). The hybridisation analysis could reveal a simple pattern of only genes encoding Rpi-blb2. A further example of such low-stringent hybridization conditions are 4XSSC at 50°C or hybridization with 30 to 40% formamide at 42°C. Such molecules comprise
- 25 those which are fragments, analogues or derivatives of Rpi-blb2 of the invention and differ, for example, by way of amino acid and/or nucleotide deletion(s), insertion(s), substitution (s), addition(s) and/or recombination (s) or any other modification(s) known in the art either alone or in combination from the above-described amino acid sequences or their underlying nucleotide sequence(s). However, it is preferred to use
- 30 high stringency hybridisation conditions.

The term "homology" means that the respective nucleic acid molecules or encoded proteins are functionally and/or structurally equivalent. The nucleic acid molecules that are homologous to the nucleic acid molecules described above and that are derivatives

35 of said nucleic acid molecules are, for example, variations of said nucleic acid molecules which represent modifications having the same biological function, in particular encoding proteins with the same or substantially the same biological function. They may be naturally occurring variations, such as sequences from other plant varieties or species, or mutations. These mutations may occur naturally or may be obtained by

40 mutagenesis techniques. The allelic variations may be naturally occurring allelic vari-

Agrico B.V.

20030596

10

ants as well as synthetically produced or genetically engineered variants. Structurally equivalents can, for example, identified by testing the binding of said polypeptide to antibodies. Structurally equivalent have the similar immunological characteristic, e.g. comprise similar epitopes.

5

The terms "fragment", "fragment of a sequence" or "part of a sequence" mean a truncated sequence of the original sequence referred to. The truncated sequence (nucleic acid or protein sequence) can vary widely in length; the minimum size being a sequence of sufficient size to provide a sequence with at least a comparable function and/or activity of the original sequence referred to, while the maximum size is not critical. In some applications, the maximum size usually is not substantially greater than that required to provide the desired activity and/or function(s) of the original sequence.

10

Typically, the truncated amino acid sequence will range from about 5 to about 1260 amino acids in length. More typically, however, the sequence will be a maximum of about 1000 amino acids in length, preferably a maximum of about 500 or 100 amino acids. It is usually desirable to select sequences of at least about 10, 12 or 15 amino acids, up to a maximum of about 20 or 25 amino acids.

15

The term "epitope" relates to specific immunoreactive sites within an antigen, also known as antigenic determinates. These epitopes can be a linear array of monomers in a polymeric composition – such as amino acids in a protein – or consist of or comprise a more complex secondary or tertiary structure. Those of skill will recognize that all immunogens (i.e., substances capable of eliciting an immune response) are antigens; however, some antigen, such as haptens, are not immunogens but may be made immunogenic by coupling to a carrier molecule. The term "antigen" includes references to a substance to which an antibody can be generated and/or to which the antibody is specifically immunoreactive. In one embodiment the present invention relates to a epitope of Rpi-b1b2.

25

30

The term "one or several amino acids" relates to at least one amino acid but not more than that number of amino acids which would result in a homology of below 70% identity. Preferably, the identity is more than 75% or 80%, more preferred are 85%, 90% or 95%, even more preferred are 96%, 97%, 98%, or 99% identity.

35

The terms "polynucleotide" and "nucleic acid molecule" also relate to "isolated" polynucleotides or nucleic acids molecules. An "isolated" nucleic acid molecule is one which is separated from other nucleic acid molecules which are present in the natural source of the nucleic acid. Preferably, an "isolated" nucleic acid is free of sequences which naturally flank the nucleic acid (i.e., sequences located at the 5' and 3' ends of the nucleic

40

acid) in the genomic DNA of the organism from which the nucleic acid is derived. For example, in various embodiments, the polynucleotide of the present invention can contain less than about 5 kb, 4kb, 3kb, 2kb, 1 kb, 0.5 kb or 0.1 kb or less of nucleotide sequences which naturally flank the nucleic acid molecule in genomic DNA of the cell from which the nucleic acid is derived. Moreover, the polynucleotides of the present invention, in particular an "isolated" nucleic acid molecule, such as a cDNA molecule, can be substantially free of other cellular material, or culture medium when produced by recombinant techniques, or chemical precursors or other chemicals when chemically synthesized.

Further, the polynucleotide of the invention comprises a nucleic acid molecule which is a complement of one of the nucleotide sequences of above mentioned polynucleotides or a portion thereof. A nucleic acid molecule which is complementary to one of the nucleotide sequences shown in SEQ ID No:1 or 3 or 5 or 6 is one which is sufficiently complementary to one of the nucleotide sequences shown in SEQ ID No: 1 or 3 or 5 or 6 such that it can hybridize to one of the nucleotide sequences shown in SEQ ID No: 1 or 3 or 5 or 6, thereby forming a stable duplex.

The polynucleotide of the invention comprises a nucleotide sequence which is at least about 70%, preferably at least about 75%, more preferably at least about 80%, 90%, or 95%, and even more preferably at least about 96%, 97%, 98%, 99% or more homologous to a nucleotide sequence shown in SEQ ID No: 1 or 3 or 5 or 6, or a portion thereof. The polynucleotide of the invention comprises a nucleotide sequence which hybridizes, preferably hybridizes under stringent conditions as defined herein, to one of the nucleotide sequences shown in SEQ ID No: 1 or 3 or 5 or 6, or a portion thereof.

Moreover, the polynucleotide of the invention can comprise only a portion of the coding region of one of the sequences in SEQ ID No: 1 or 3 or 5 or 6, for example a fragment which can be used as a probe or primer or a fragment encoding a biologically active portion of an Rpi-blb2. The nucleotide sequences determined from the cloning of the the present Rpi-blb2 protein encoding gene allows for the generation of probes and primers designed for use in identifying and/or cloning its homologues in other cell types and organisms. The probe/primer typically comprises substantially purified oligonucleotide. The oligonucleotide typically comprises a region of nucleotide sequence that hybridizes under stringent conditions to at least about 12, 15 preferably about 20 or 25, more preferably about 40, 50 or 75 consecutive nucleotides of a sense strand of one of the sequences set forth, e.g., in SEQ ID No. No: 1 or 3 or 5 or 6, an anti-sense sequence of one of the sequences, e.g., set forth in SEQ ID No.: 1 or 3 or 5 or 6, or naturally occurring mutants thereof. Primers based on a nucleotide of invention can be used in PCR reactions to clone Rpi-blb2 homologues, e.g. as the primers described

in the examples of the present invention, e.g. as shown in tab 3a or 3b, preferably the primers ARF1F and ARF1R are used. A PCR with the primers univ24R and univ14L will result in a fragment of Rpi-blb2 which can be used as described herein. Said primer sets are interchangeable. The person skilled in the art knows to combine said primers to result in the desired product, e.g. in a full length clone or a partial sequence. Probes based on the Rpi-blb2 nucleotide sequences can be used to detect transcripts or genomic sequences encoding the same or homologous proteins. The probe can further comprise a label group attached thereto, e.g. the label group can be a radioisotope, a fluorescent compound, an enzyme, or an enzyme co-factor. Such probes can be used as a part of a genomic marker test kit for identifying cells which express an Rpi-blb2, such as by measuring a level of an Rpi-blb2-encoding nucleic acid molecule in a sample of cells, e.g., detecting Rpi-blb2 mRNA levels or determining whether a genomic Rpi-blb2 gene has been mutated or deleted.

The polynucleotide of the invention encodes a polypeptide or portion thereof which includes an amino acid sequence which is sufficiently homologous to the amino acid sequence of SEQ ID No: 2 or 4 such that the protein or portion thereof maintains the ability to participate in resistance to pathogens, in particular a Rpi-blb2 activity as described in the examples in plants. As used herein, the language "sufficiently homologous" refers to proteins or portions thereof which have amino acid sequences which include a minimum number of identical or equivalent (e.g., an amino acid residue which has a similar side chain as an amino acid residue in one of the sequences of the polypeptide of the present invention, amino acid residues to an amino acid sequence of Seq. ID No.: 2 or 4 such that the protein or portion thereof is able to participate in the resistance of plants to said pathogens. Examples of a Rpi-blb2 activity are described herein. Thus, the function of an Rpi-blb2 protein contributes either directly or indirectly to the resistance to plant pathogens, preferably to the pathogens mentioned herein, more preferred to *P. infestans*.

The protein is at least about 70%, preferably at least about 75%, and more preferably at least about 80%, 90%, 95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of SEQ ID No: 2 or 4.

Portions of proteins encoded by the polynucleotide of the invention are preferably biologically active.

As mentioned herein, the term "biologically active portion" is intended to include a portion, e.g., a domain/motif, that confers resistance to an oomycete plant pathogene and/or *Bemisia tabaci* and/or aphids or has an immunological activity such that it is binds to an antibody binding specifically to Rpi-blb2 or it has an activity as set forth in the Examples or as described herein.

Additional nucleic acid fragments encoding biologically active portions of the polypeptide of the present invention can be prepared by isolating a portion of one of the sequences in SEQ ID No: 1 or 3 or 5 or 6, expressing the encoded portion of the Rpi-blb2 or peptide (e.g., by recombinant expression *in vitro*) and assessing the activity of the encoded portion of the protein.

The invention further encompasses polynucleotides that differ from one of the nucleotide sequences shown in SEQ ID No: 1 or 3 or 5 or 6 (and portions thereof) due to degeneracy of the genetic code and thus encode a Rpi-blb2 as that encoded by the sequences shown in SEQ ID No: 2 or 4. Further the polynucleotide of the invention has a nucleotide sequence encoding a protein having an amino acid sequence shown in SEQ ID No: 2 or 4. In a still further embodiment, the polynucleotide of the invention encodes a full length protein which is substantially homologous to an amino acid sequence of SEQ ID No: 2 or 4.

In addition, it will be appreciated by those skilled in the art that DNA sequence polymorphisms that lead to changes in the amino acid sequences may exist within a population (e.g., the *S. bulbocastanum* population). Such genetic polymorphism in the Rpi-blb2 gene may exist among individuals within a population due to natural variation.

As used herein, the terms "gene" and "recombinant gene" refer to nucleic acid molecules comprising an open reading frame encoding an Rpi-blb2, preferably a *S. bulbocastanum* Rpi-blb2. Such natural variations can typically result in 1-5% variance in the nucleotide sequence of the Rpi-blb2 gene. Any and all such nucleotide variations and resulting amino acid polymorphisms in Rpi-blb2 that are the result of natural variation and that do not alter the functional activity of Rpi-blb2 are intended to be within the scope of the invention.

Polynucleotides corresponding to natural variants and non-*S. bulbocastanum* homologues of the Rpi-blb2 cDNA of the invention can be isolated based on their homology to *S. bulbocastanum* Rpi-blb2 polynucleotides disclosed herein using the polynucleotide of the invention, or a portion thereof, as a hybridization probe according to standard hybridization techniques under stringent hybridization conditions. Accordingly, in another embodiment, an polynucleotide of the invention is at least 20 nucleotides in length. Preferably it hybridizes under stringent conditions to the nucleic acid molecule comprising a nucleotide sequence of the polynucleotide of the present invention, e.g. SEQ ID No: 1 or 3 or 5 or 6. In other embodiments, the nucleic acid is at least 20, 30, 50, 100, 250 or more nucleotides in length. The term "hybridizes under stringent conditions" is defined above and is intended to describe conditions for hybridization and washing under which nucleotide sequences at least 65% identical to each other

Agrico B.V.

20030596

PF 54801

14

typically remain hybridized to each other. Preferably, the conditions are such that sequences at least about 70%, more preferably at least about 75% or 80%, and even more preferably at least about 85%, 90% or 95% or more identical to each other typically remain hybridized to each other. Preferably, polynucleotide of the invention that
5 hybridizes under stringent conditions to a sequence of SEQ ID No: 1 or 3 or 5 or 6 corresponds to a naturally-occurring nucleic acid molecule.

As used herein, a "naturally-occurring" nucleic acid molecule refers to an RNA or DNA molecule having a nucleotide sequence that occurs in nature (e.g., encodes a natural
10 protein). Preferably, the polynucleotide encodes a natural *S. bulbocastanum* Rpi-blb2.

In addition to naturally-occurring variants of the Rpi-blb2 sequence that may exist in the population, the skilled artisan will further appreciate that changes can be introduced by mutation into a nucleotide sequence of the polynucleotide encoding Rpi-blb2, thereby
15 leading to changes in the amino acid sequence of the encoded Rpi-blb2, without altering the functional ability of the Rpi-blb2. For example, nucleotide substitutions leading to amino acid substitutions at "non-essential" amino acid residues can be made in a sequence of the polynucleotide encoding Rpi-blb2, e.g. SEQ ID No: 1 or 3 or 5 or 6. A "non-essential" amino acid residue is a residue that can be altered from the wild-type
20 sequence of one of the Rpi-blb2 without altering the activity of said Rpi-blb2, whereas an "essential" amino acid residue is required for Rpi-blb2 activity. Other amino acid residues, however, (e.g., those that are not conserved or only semi-conserved in the domain having Rpi-blb2 activity) may not be essential for activity and thus are likely to be amenable to alteration without altering Rpi-blb2 activity.

25 Accordingly, a person skilled in the art knows that the codon usage between organism can differ. Therefore he will adapt the codon usage in the polynucleotide of the present invention to the usage of the organism in which the polynucleotide or polypeptide is expressed.

30

Accordingly, the invention relates to polynucleotides encoding Rpi-blb2 that contain changes in amino acid residues that are not essential for Rpi-blb2 activity. Such Rpi-blb2s differ in amino acid sequence from a sequence contained in SEQ ID No: 2 or 4 yet retain the Rpi-blb2 activity described herein. The polynucleotide can comprise
35 a nucleotide sequence encoding a polypeptide, wherein the polypeptide comprises an amino acid sequence at least about 70% identical to an amino acid sequence of EQ ID No: 2 or 4 and is capable of participation in the resistance to a plant pathogen. Preferably, the protein encoded by the nucleic acid molecule is at least about 70%
40 identical to the sequence in SEQ ID No: 2 or 4, more preferably at least about 75% identical to one of the sequences in SEQ ID No: 2 or 4, even more preferably at least

15

bly at least about 80%, 90%, 95% homologous to the sequence in SEQ ID No: 2 or 4, and most preferably at least about 96%, 97%, 98%, or 99% identical to the sequence in SEQ ID No: 2 or 4.

- 5 To determine the percent homology of two amino acid sequences (e.g., one of the sequences of Seq. ID No.: 2 or 4 and a mutant form thereof) or of two nucleic acids, the sequences are aligned for optimal comparison purposes (e.g., gaps can be introduced in the sequence of one protein or nucleic acid for optimal alignment with the other protein or nucleic acid). The amino acid residues or nucleotides at corresponding amino acid positions or nucleotide positions are then compared. When a position in
- 10 one sequence (e.g., one of the sequences of SEQ ID No: 2 or 4) is occupied by the same amino acid residue or nucleotide as the corresponding position in the other sequence (e.g., a mutant form of the sequence selected), then the molecules are homologous at that position (i.e., as used herein amino acid or nucleic acid "homology" is equivalent to amino acid or nucleic acid "identity"). The percent homology between
- 15 the two sequences is a function of the number of identical positions shared by the sequences (i.e., % homology = numbers of identical positions/total numbers of positions x 100).

- Homology can be calculated by comparison with the aid of the program algorithm GAP (Wisconsin Package Version 10.0, University of Wisconsin, Genetics Computer Group
- 20 (GCG), Madison, USA; Altschul et al. (1997) Nucleic Acids Res. 25:3389 et seq.), setting the following parameters:

Gap weight:	50	Length weight:	3
Average match:	10	Average mismatch:	0

- For example a sequence which has at least 80% homology with sequence SEQ ID
- 25 NO: 1 at the nucleic acid level is understood as meaning a sequence which, upon comparison with the sequence SEQ ID NO: 1 by the above program algorithm with the above parameter set, has at least 80% homology.

- Homology between two polypeptides is understood as meaning the identity of the amino acid sequence over in each case the entire sequence length which is calculated
- 30 by comparison with the aid of the program algorithm GAP (Wisconsin Package Version 10.0, University of Wisconsin, Genetics Computer Group (GCG), Madison, USA), setting the following parameters:

Gap weight:	8	Length weight:	2
Average match:	2,912	Average mismatch:	-2,003

- 35 For example a sequence which has at least 80% homology with sequence SEQ ID NO: 2 at the protein level is understood as meaning a sequence which, upon comparison with the sequence SEQ ID NO: 2 by the above program algorithm with the above parameter set, has at least 80% homology.

Agrico B.V.

20030596

PF 54801

17

erably, conservative amino acid substitutions are made at one or more predicted non-essential amino acid residues. A "conservative amino acid substitution" is one in which the amino acid residue is replaced with an amino acid residue having a similar side chain. Families of amino acid residues having similar side chains have been defined in the art. These families include amino acids with basic side chains (e.g., lysine, arginine, histidine), acidic side chains (e.g., aspartic acid, glutamic acid), uncharged polar side chains (e.g., glycine, asparagine, glutamine, serine, threonine, tyrosine, cysteine), non-polar side chains (e.g., alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), beta-branched side chains (e.g., threonine, valine, isoleucine) and aromatic side chains (e.g., tyrosine, phenylalanine, tryptophan, histidine). Thus, a predicted nonessential amino acid residue in an Rpi-blb2 is preferably replaced with another amino acid residue from the same family. Alternatively, in another embodiment, mutations can be introduced randomly along all or part of an Rpi-blb2 coding sequence, such as by saturation mutagenesis, and the resultant mutants can be screened for an Rpi-blb2 activity described herein to identify mutants that retain Rpi-blb2 activity. Following mutagenesis of one of the sequences of SEQ ID No: 1 or 3 or 5 or 6, the encoded protein can be expressed recombinantly and the activity of the protein can be determined using, for example, assays described herein (see Examples).

20

In one embodiment, in the method of present invention the activity of Rpi-blb2 protein and of a further resistance protein is increased.

It is expected, that under field conditions the presence of more than one resistance gene is beneficial, in particular genes conferring resistance to the same pathogen. In case a pathogen isolate, e.g. a *P. infestans* isolate, is present that is able to overcome resistance of one of the R-genes, the other one or more R-gene(s) is/are still functional making it impossible to infect the plant. The present of two undefeated R-genes strongly reduces the chance that a pathogen, in particular a *P. infestans* isolate, is able to mutate into an isolate that can overcome two or more R-genes.

In the following "resistance polypeptide" or "resistance protein" relates to a polypeptide which (increased) activity will confer resistance to a susceptible strain ("wild type" or "reference"). Accordingly, Rpi-blb2 is a resistance protein as well as e.g. Rpi-blb (or RB). A "further resistance protein" relates to an other resistance protein than the protein of the present invention, whereas the term "resistance protein" comprises the polypeptid of the present invention as well as one or more further resistance protein(s). It is further understood, that the term "and a further resistance protein" relates to one or more further resistance proteins. Thus, the activity of one or more resistance proteins can be increased. Further resistance proteins are described below. However, generally any other known resistance protein can be co-expressed with the polypeptid of the pre-

sent invention or its activity can be increased by any of the methods described herein for *Rpi-blb2*.

5 In a preferred embodiment, the further resistance protein comprises a LRR domain and a P-loop.

The cloning and molecular characterisation of over 30 plant disease resistance (*R*) genes conferring resistance to bacteria, fungi, oomycetes, viruses, nematodes, or insects has allowed their classification in structural classes regardless of pathogen specificity (reviewed in Dangl and Jones, 2001). The most abundant class of characterised *R* genes, comprising about 0.5 percent of the genes predicted in the *Arabidopsis* genome, is predicted to encode intracellular proteins that carry leucine-rich repeat (LRR) and nucleotide-binding site (NBS) domains, motifs also found in other receptor and signal transduction proteins. NBS-LRR *R* proteins differ primarily at the N-terminus that either exhibits sequence similarity to the *Drosophila* Toll protein and the mammalian interleukin-1 receptor domain (TIR-NBS-LRR), or code for a coiled-coils structure (CC-NBS-LRR), sometimes in the form of a leucine zipper (LZ-NBS-LRR. Although maybe membrane associated, NBS-LRR proteins are predicted to be cytoplasmic. In contrast, two other classes of *R* proteins that carry LRRs are predicted to span the cell membrane, with an extracellular LRR domain: the LRR-transmembrane (LRR-TM) *Cf* proteins and the LRR-TM-kinase *Xa21* protein. Characterised *R* proteins that lack LRRs are the *Pto* gene from tomato, the *Hs1^{Pro-1}* gene from beet, the *mlo* gene from barley, the *Rpw8* genes from *Arabidopsis* and the *Rpg1* gene from barley.

According to the gene-for-gene hypothesis, disease resistance follows perception by plant *R* proteins of pathogen effector molecules with avirulence (*Avr*) function, thereby initiating through some kind of elicitor recognition complex, signal transduction pathways leading to a hypersensitive response (HR). In common with other receptors it is generally considered that NBS-LRR *R* proteins have a modular structure with separate recognition and signalling domains, whereby the LRR is the candidate recognition domain and the N-terminal region including the NBS, the major signalling domain. Functional analysis of recombinant *R* proteins indicates that recognition specificity indeed resides in the LRR. Moreover, the LRR is the most variable region in closely related NBS-LRR proteins and is under selection to diverge. However, evidence is accumulating that LRRs also contribute to signalling through negative regulation involving putative intramolecular interactions. Currently, five *R* genes have been cloned from potato, including two *R* genes conferring resistance to late blight, and all belong to the CC/LZ-NBS-LRR class of plant *R* genes. While the *S. demissum* derived *R1* gene confers race specific resistance to late blight, the recently cloned *S. bulbocastanum* derived gene *Rpi-blb* (or *RB*) confers full resistance to a range of *P. infestans* isolates carrying multiple virulence factors and race-specificity has not yet been demonstrated. Furthermore, as described before, progeny plants of somatic hybrids containing *Rpi-blb* were

unaffected by late blight on field experiments in Mexico, where nearly every race of the fungus is found. Through complementation of the susceptible phenotype in cultivated potato and tomato the potential of interspecific transfer of broad-spectrum late blight resistance to cultivated *Solanaceae* from sexually incompatible host species by trans-
5 formation with single cloned *R* genes was demonstrated. US 6,127,607 describes resistance proteins with LRR domains and P-loops. The content of US 6,127,607 is herewith incorporated by reference. In particular columns 6 to 8 and col. 11 describe LRR domains and P-loops. Furthermore Song, 2003, PNAS, 100, 16, 9128 shows a comparison of Rpi-blb LRR motifs in Fig. 4 and gives on pages 9132 an overview about
10 LRR domains. The domains of the polypeptid of the present invention are shown in Fig. 14 as well as in Fig. 15.

Preferably the activity of one or more resistance protein(s) selected from the group consisting of Rpi-blb, R1, R-ber, Rpi1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11,
15 Ph-1, Ph-2 and Ph-3 is increased. Preferred is that also the Rpi-blb activity is increased.

In one embodiment of the present invention, the expression of an, e.g. transgenic, Rpi-blb2 protein is increased and further a transgenic resistance gene's expression is in-
20 creased. The resistance protein coexpressed with the Rpi-blb2 is preferably one of the resistance proteins mentioned herein, in particular Rpi-blb, R1, Rpi1, R-ber, R2, R3, R6, R7, Ph-1, Ph-2 or Ph-3 but can also be one of the others resistance to plant pathogens conferring proteins known to a person skilled in the art.

As mentioned, the term "increased expression" according to this invention also includes
25 a de novo-Expression of a polynucleotide or polypeptide.

Most preferred is an increase of resistance via coexpression of the polypeptid of the present invention together with Rpi-blb. Rpi-blb and Rpi-blb2 provide both full resistance in detached leaf assays to *P. infestans* isolates as described in the examples,
30 and in Song 2003, PNAS, 100, 16, 9128.

Said resistance conferring genes are for example described in RB (synonym of Rpi1b): AY336128 [gi: 32693280], (Song et al., 2003). BAC clones 177 013 and CB3A14 comprising the Rpi-blb gene have been deposited in GenBank
35 with accession nos AY303171 and AY303170.

R1: AF447489 [gi: 9117432422], (Ballvora et al., 2002)

Rpi1: Kuhl, J.C., Hanneman, R.E., and Havey, M.J., (2001) Characterization and mapping of Rpi1, a late blight resistance locus from diploid (1EBN) Mexican *Solanum pin-*
natisectum. Molecular genet. Genomics 265: 977-985.

R-ber: Ewing, E.E., Simko, J., Smart, C.D., Bonierbale, M.W., Mizubuti, E.S.G., May, G.D., and Fry, W.E., (200) Genetic mapping from field tests of qualitative and quantitative resistance to *Phytophthora infestans* in a population derived from *Solanum tuberosum* and *Solanum berthaultii*. *Molecular breeding* 6:25-36.

- 5 R2: Li, X., vanEck, H.J., vanderVoort, J.N.A.M., Huigen, D.J., Stam, P., and Jacobsen, E. (1998) Autotetraploids and genetic mapping using common AFLP markers: the R2 allele conferring resistance to *Phytophthora infestans* mapped on potato chromosome 4. *Theoretically and Applied Genetics* 96 (8): 1121-112.

- 10 R3, R6, R7: Elkhartbotly, A., Palominosanchez, C., Salami, F., Jacobsen, E., and Gebhardt, C. (1996) R6 and R7 alleles of potato conferring race-specific resistance to *Phytophthora infestans* (Mont) de Bary identified genetic loci clustering with the R3 locus on chromosome XI. *Theoretical and Applied Genetics* 92 (7): 880-884.

Ph-1: Bonde and Murphy (1952) *Main Agric. Exp. Stn. Bull.* No 497

- 15 Ph-2: Moreau, P., Thoquet, P., Olivier, J., Laterrot, H., and Grimsley, N.H. (1998) Genetic mapping of Ph-2, a single locus controlling partial resistance to *Phytophthora infestans* in tomato. *Molecular Plant Microbe Interactions* 11 (4): 259-269.

Ph-3: Chunwongse, J., Chunwongse, C., Black, L., and Hanson, P. (2002) Molecular mapping of the Ph-3 gene for late blight resistance in tomato. *Journal of Horticultural Science & Biotechnology* 77 (3): 281-286.

20

In one embodiment, the activity of the Rpi-blb2 protein is increased or the polynucleotide of the invention is expressed together with a nucleic acid molecule encoding Rpi-blb, R1, R-ber, Rpi1, R2, R3, R6, R7, Ph-1, Ph-2 and/or Ph-3 selected from the group consisting of:

- 25 a) nucleic acid molecule encoding at least a mature form of a Rpi-blb (or RB-) polypeptide as encoded by the sequence shown in GenBank Accession no.: AY336128 [gi: 32693280], or the R1 polypeptide as encoded by the sequence shown in GenBank Accession no.: AF447489 [gi 9117432422], or a nucleic acid molecule encoding at least a mature form of a resistance conferring protein mapped and characterized as described for Rpi1 in Kuhl, J.C., Hanneman, R.E., and Havey, M.J., (2001) Characterization and mapping of Rpi1, a late blight resistance locus from diploid (1EBN) Mexican *Solanum pinnatisectum*. *Molecular genet. Genomics* 265: 977-985; for R-ber in Ewing, E.E., Simko, J., Smart, C.D., Bonierbale, M.W., Mizubuti, E.S.G., May, G.D., and Fry, W.E., (2000) Genetic mapping from field tests of qualitative and quantitative resistance to *Phytophthora infestans* in a population derived from *Solanum tuberosum* and *Solanum berthaultii*. *Molecular breeding* 6:25-36; for R2 in Li, X., vanEck, H.J., vanderVoort, J.N.A.M., Huigen, D.J., Stam, P., and Jacobsen, E. (1998) Autotetraploids and genetic mapping using common AFLP markers: the R2 allele conferring resistance to *Phytophthora infestans* mapped on potato chromosome 4. *Theoretically and Applied Genetics* 96 (8): 1121-1128, for
- 30
- 35
- 40

R3, R6, R7 in Elkhartbotly, A., Palominosanchez, C., Salamini, F., Jacobsen, E., and Gebhardt, C. (1996) R6 and R7 alleles of potato conferring race-specific resistance to *Phytophthora infestans* (Mont) de Bary identified genetic loci clustering with the R3 locus on chromosome XI. Theoretical and Applied Genetics 92 (7): 880-884; for Ph-1 in Bonde and Murphy (1952) Main Agric. Exp. Stn. Bull. No 497; for Ph-2 in Moreau, P., Thoquet, P., Olivier, J., Laterrot, H., and Grimsley, N.H. (1998) Genetic mapping of Ph-2, a single locus controlling partial resistance to *Phytophthora infestans* in tomato. Molecular Plant Microbe Interactions 11 (4): 259-269; and for Ph-3 in Chunwongse, J., Chunwongse, C., Black, L., and Hanson, P. (2002) Molecular mapping of the Ph-3 gene for late blight resistance in tomato. Journal of Horticultural Science & Biotechnology 77 (3): 281-286.

- b) nucleic acid molecule the nucleotide sequence of which is degenerate as a result of the genetic code to a nucleotide sequence of (a);
 - c) nucleic acid molecule encoding a polypeptide derived from the polypeptide encoded by a polynucleotide of (a) or (b) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence of the polypeptide encoded by a polynucleotide of (a) or (b);
 - d) nucleic acid molecule encoding a polypeptide the sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleic acid molecule of (a);
 - e) nucleic acid molecules comprising a fragment or a epitope-bearing portion of a polypeptide encoded by a nucleic acid molecule of any one of (a) to (d);
 - f) nucleic acid molecule encoding a fragment beginning with amino acid: 1, 30, 50, 100, 200, 500, 1000, or 2000 and stopping with amino acid 2000, 1000, 500, 300, 200, 50, or 1 of a polypeptide encoded by any one of (a) to (e) and with one of said activities;
 - g) nucleic acid molecule comprising at least 20 nucleotides of a polynucleotide of any one of (a) or (b);
 - h) nucleic acid molecule encoding a polypeptide being recognized by a monoclonal antibody that have been raised against a polypeptide encoded by a nucleic acid molecule of any one of (a) to (f);
 - i) nucleic acid molecule obtainable by screening an appropriate library under stringent conditions with a probe having the sequence of the nucleic acid molecule of any one of (a) to (h) or of a fragment thereof of at least 20, preferable 30 or more nucleotides; and
 - j) nucleic acid molecule the complementary strand of which hybridises under stringent conditions with a nucleic acid molecule of any one of (a) or (i);
- or the complementary strand of any one of (a) to (j).

Agrico B.V.

20030596

PF 54801

22

Accordingly, the method of present invention confers resistance of one of said plants, plant tissue or plant cell of the present invention to a plant pathogen of a phylum Oomycetes, preferably to a pathogen of the order Pythiales or Peronosporales, more preferred to the family Pythiaceae or Peronosporaceae, more preferred of the genus

5 Phytophthora or Bremia or Peronospora or Plasmopara, most preferred wherein the pathogen is of the species *Phytophthora parasitica* var. *nicotianae* (causing, amongst others, black shank in tobacco), *Phytophthora sojae* (causing *Phytophthora* root rot in soybean), *Phytophthora capsici* (causing rots in pepper and cucurbits and tomato), *Phytophthora erythroseptica* (causing Pink rot in potato), *Plasmopara viticola* (causing

10 grapevine downy mildew), *Bremia lactuca* (causing downy mildew in lettuce) or *Peronospora tabaci* (causing blue mould in tobacco).

The activity of Rpi-blb2 in a plant, a plant cell, a plant tissue, a plant organ or part thereof according to the present invention can be increased, generated or stimulated

15 via methods which are well known to a person skilled in the art and e.g. are described in Sambrook et al., Cold Spring Harbor Laboratory Press, NY, 1989.

Thus, in a preferred embodiment, the present invention relates to the method of the invention, wherein the expression is a *de novo* expression.

20 The term "de novo-Expression" as understood herein relates to a non-detectability of a polypeptide or polynucleotide. Preferred is that no gene encoding a polypeptide or a polynucleotide which should be *de novo*-expressed is present in the genome. However, if the expression can not be detected, it is generally assumed that no expression

25 occurs. A person skilled in the art, however, knows that the detection methods and means develop to higher sensitivity. Thus, *de novo*-Expression also relates to expression in systems, where the level of expression is too low to confer any resistance to a plant pathogen. A comparison of a knock out strain and a low-expression strain-Phenotyp can show, whether any difference in resistance to any of the herein mentioned pathogens is observable.

30

Accordingly, in another embodiment of the present invention, the endogenous activity of a Rpi-blb2 and/or a further resistance protein is increased.

The level of expression in a cell can be increased by methods known to a person

35 skilled in the art. Several techniques are described herein, e.g. the transgenic expression of the polynucleotide or polypeptide of the present invention. The polynucleotide or polypeptide can be of foreign origin. Preferred that a polynucleotide of the same genetic origin than the host cell, plant cell, plant tissue, plant is introduced.

Agrico B.V.

20030596

PF 54801

23

The activity, in particular an endogenous activity but also the activity of a transgenic expressed Rpi-blb2 can be increased by several methods. Accordingly, in a preferred embodiment, the activity of the resistance proteins described herein is increased by one or more of the following steps

- 5 a) stabilizing the resistance protein;
- b) stabilizing the resistance protein encoding mRNA;
- c) increasing the specific activity of the resistance protein;
- d) expressing or increasing the expression of a homologous or artificial transcription factor for resistance expression;
- 10 e) stimulate resistance protein activity through exogenous inducing factors;
- f) expressing a transgenic resistance gene; and/or
- g) increasing the copy number of the resistance encoding gene.

- 15 In general an activity in a organism, in particular in a plant cell, a plant, or a plant tissue can be increased by increasing the amount of the specific protein, i.e. of the resistance protein, in said organism. "Amount of protein" is understood as meaning the amount of a polypeptide, preferably Rpi-blb2, in an organism, a tissue, a cell or a cell compartment. "Increase" of the amount of protein means the quantitative increase of the amount of a protein in an organism, a tissue, a cell or a cell compartment - for example
- 20 by one of the methods described herein below - in comparison with the wild type of the same genus and species, to which this method had not been applied, under otherwise identical conditions (such as, for example, culture conditions, plant age and the like). The increase amounts to at least 10%, preferably at least 20% or at least 50%, especially preferably at least 70% or 90%, very especially preferably at least 100%, most
- 25 preferably at least 200% or more.

- "Increase" of the activity is understood as meaning the reduction of the total activity of a protein in an organism, a tissue, a cell or a cell compartment in comparison with the wild type of the same genus and species, to which this method had not been applied,
- 30 under otherwise identical conditions (such as, for example, culture conditions, plant age and the like). The increase amounts to at least 10%, preferably at least 20% or at least 50%, especially preferably at least 70% or 90%, very especially preferably at least 100%, most preferably at least 200% or more.

- 35 In this context, the efficacy of the pathogen resistance can deviate both downward or upward in comparison with a value obtained when increasing one of the Rpi-blb2 proteins as shown in SEQ ID NO: 2 or 4. Preferred functional equivalents are those in which the efficacy of the pathogen resistance - measured, for example, by the penetration efficacy of a pathogen or as described herein - differs by not more than 50%,
- 40 preferably 25%, especially preferably 10% from a comparative value obtained by reducing an Rpi-blb2 protein as shown in SEQ ID NO: 2 or 4. Especially preferred are

Agrico B.V.

20030596

PF 54801

24

ducing an Rpi-blb2 protein as shown in SEQ ID NO: 2 or 4. Especially preferred are those sequences where the increase increases the efficacy of pathogen resistance quantitatively by more than 50%, preferably 100%, especially preferably 500%, very especially preferably 1000% based on a comparative value obtained by reducing one
5 of the Rpi-blb2 proteins as shown in SEQ ID NO: 2 or 4.

Any comparison is preferably carried out under analogous conditions. "Analogous conditions" means that all conditions such as, for example, culture or growing conditions, assay conditions (such as buffer, temperature, substrates, pathogen concentration
10 and the like) are kept identical between the experiments to be compared and that the set-ups differ only by the sequence of the Rpi-blb2 polypeptides to be compared, their organism of origin and, if appropriate, the pathogen. When choosing the pathogen, each comparison requires that the pathogen be chosen which is most similar to the other equivalent, taking into consideration the species specificity.

15 Due to the increased Rpi-blb2 activity, the resistance of a plant or a part thereof is increased. In a preferred embodiment, the method of the present invention results in reduction in the sporulation index of at least 30% after infection with *P. infestans* compared to a wild type, more preferred is a reduction of 50%, even more preferred
20 are 70%, even more preferred are more than 80%, more preferred are 85% and 90%. Most preferred is 95% or more.

Accordingly, the present invention also relates to said polynucleotide of the invention, as defined above, is polynucleotide encoding a Rpi-blb2 protein comprising a nucleic
25 acid molecule selected from the group consisting of:

- a) nucleic acid molecules encoding at least the mature form of the polypeptide depicted in SEQ ID NO: 2 or 4;
- b) nucleic acid molecules comprising the coding sequence as depicted in SEQ ID NO: 1 or 3 or 5 or 6 or encoding at least the mature form of the polypeptide;
- 30 c) nucleic acid molecules the nucleotide sequence of which is degenerate as a result of the genetic code to a nucleotide sequence of (a) or (b);
- d) nucleic acid molecules encoding a polypeptide derived from the polypeptide encoded by a polynucleotide of (a) to (c) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence of the polypeptide
35 encoded by a polynucleotide of (a) to (c);
- e) nucleic acid molecules encoding a polypeptide the sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleic acid molecule of (a) or (b);

Agrico B.V.**20030596****PF 54801**

25

- f) nucleic acid molecules comprising a fragment or a epitope-bearing portion of a polypeptide encoded by a nucleic acid molecule of any one of (a) to (e);
- g) nucleic acid molecules comprising a polynucleotide having a sequence of a nucleic acid molecule amplified from a nucleic acid library using the primers as listed in
5 Tab. 3b, preferably ARF1F or ARF1R;
- h) nucleic acid molecules encoding polypeptide fragment beginning with amino acid: 1, 30, 50, 100, 200, 300, 500, or 1000 and stopping with amino acid 1276, 1000, 500, 300, 200, 50, or 30 of a polypeptide encoded by any one of (a) to (g);
- i) nucleic acid molecules comprising at least 20 nucleotides of a polynucleotide of
10 any one of (a) or (d);
- j) nucleic acid molecules encoding a polypeptide being recognized by a monoclonal antibodies that have been raised against a polypeptide encoded by a nucleic acid molecule of any one of (a) to (h);
- k) nucleic acid molecules obtainable by screening an appropriate library under strin-
15 gent conditions with a probe having the sequence of the nucleic acid molecule of any one of (a) to (j) or of a fragment thereof of at least 15, preferable 30, 60, 90 or more nucleotides; and
- l) nucleic acid molecules the complementary strand of which hybridises under strin-
gent conditions with a nucleic acid molecule of any one of (a) or (k);
- 20 or the complementary strand of any one of (a) to (l);
or encoding a polypeptide encoded by a segment of chromosome 6 or of linkage group 6 of *Solanum bulbocastanum* which co-segregates with a marker selected from table 3a or 3b and which mediates resistance to plant pathogens, preferably of the phylum Oomyceta;
- 25 and whereby the polynucleotide has not the sequence of Mi1.1 or Mi1.2, as e.g. depicted in Seq. ID NO.: 7 and 9.

In an further embodiment, the polynucleotide of the present invention is derived or iso-
lated from the genome of a organism selected from the group consisting of Menyan-
30 thaceae, Solanaceae, Sclerophyllacaceae, Duckeodendraceae, Goetzeaceae, Convol-
vulaceae, Cuscutaceae, Polemoniaceae, and Hydrophyllaceae according to the Sys-
teme Naturae 2000, Brands, S.J., Amsterdam or has its origin thereof, more preferably
it is selected from the group consisting of *Atropa*, *Browallia*, *Brunfelsia*, *Capsicum*, *Ces-
trum*, *Cyphomandra*, *Datura*, *Fabiana*, *Franciscea*, *Hyoscyamus*, *Lycium*, *Mandragora*,
35 *Nicandra*, *Nicotiana*, *Petunia*, *Physalis*, *Schizanthus* and *Solanum* according to the
Systema Naturae 2000, Brands, S.J., Amsterdam or has its origin thereof, even more
preferred is a selection out of the group consisting of Solanaceae family, preferably
S. bulbocastanum, potato (*S. tuberosum*), tomato (*S. lycopersicum*), petunia, tree

Agrico B.V.

20030596

PF 54801

26

tomato (*S. betaceum*), pear melon (*S. muricatum*) and eggplant (*S. melongena*). Even more preferred are tomato or potato or *S. bulbocastanum* as source for the polynucleotide of the present invention. Most preferred is *S. bulbocastanum* as source.

- 5 Rpi-blb2 has been isolated from *S. tuberosum* material derived from ABPT. Thus, taxonomic perspective the Rpi-blb2 described is also *S. tuberosum*-derived. However, the gene was present on an introgression fragment presumably derived from *S. bulbocastanum*. A lot of *S. tuberosum* varieties contain introgression fragments of related *Solanum* species, but still are *S. tuberosum*. Therefore, *S. tuberosum* can ac-
- 10 cording to the taxonomical system also be a source for the polynucleotide of the present invention, in particular ABPT-derived *S. tuberosum*, as well as other varieties of other *Solanum* species varieties derived in a similar way.
- Accordingly, in another embodiment the polynucleotide of the present invention is derived from *S. tuberosum*.

15

- A polynucleotide of the present invention, e.g., a nucleic acid molecule having a nucleotide sequence of Seq ID NO: 1 or 3 or 5 or 6, or a portion thereof, can be isolated using standard molecular biology techniques and the sequence information provided herein. For example, Rpi-blb2 cDNA can be isolated from a library using all
- 20 or portion of one of the sequences of the polynucleotide of the present invention as a hybridization probe and standard hybridization techniques (e.g., as described in Sambrook et al., *Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).
- Moreover, a polynucleotide encompassing all or a portion of one of the sequences of
- 25 the polynucleotide of the present invention can be isolated by the polymerase chain reaction using oligonucleotide primers designed based upon this sequence (e.g., a nucleic acid molecule encompassing all or a portion of one of the sequences of polynucleotide of the present invention can be isolated by the polymerase chain reaction using oligonucleotide primers as mentioned above, designed based upon this same
- 30 sequence of polynucleotide of the present invention. For example, mRNA can be isolated from cells, e.g. *S. bulbocastanum* or another plant (e.g., by the guanidinium-thiocyanate extraction procedure of Chirgwin et al. (1979) *Biochemistry* 18: 5294-5299) and cDNA can be prepared using reverse transcriptase (e.g., Moloney MLV reverse transcriptase, available from Gibco/BRL, Bethesda, MD; or AMV reverse transcriptase,
- 35 available from Seikagaku America, Inc., St. Petersburg, FL). Synthetic oligonucleotide primers for polymerase chain reaction amplification can be designed based upon one of the nucleotide sequences shown in SEQ ID No: 1 or 3 or 5 or 6. A polynucleotide of the invention can be amplified using cDNA or, alternatively, genomic DNA, as a template and appropriate oligonucleotide primers according to standard PCR amplifi-

Agrico B.V.**20030596****PF 54801****27.**

cation techniques. The polynucleotide so amplified can be cloned into an appropriate vector and characterized by DNA sequence analysis. Furthermore, oligonucleotides corresponding to an Rpi-blb2 nucleotide sequence can be prepared by standard synthetic techniques, e.g., using an automated DNA synthesizer.

5

In an embodiment of the present invention is the Rpi-blb 2 protein is encoded by a segment of chromosome 6 or linkage group 6 of *Solanum bulbocastanum* or *S. tuberosum*.

10

Further the present invention comprises a segment of chromosome 6 or linkage group 6 of *S. bulbocastanum* or *S. tuberosum*. In one preferred embodiment in the method of the present invention the Rpi-blb2 protein expressed is encoded by a polynucleotide comprising a segment of chromosome 6 or linkage group 6 of *S. bulbocastanum*. Preferably said segment a group comprises further cis acting element, e.g. promoters, enhancers, binding sites etc. or trans acting elements, like cofactors, activators or other resistance proteins, which confer a increased resistance. Genomic fragments comprising the Rpi-blb2 gene and further regulatory elements are depicted in Seq. ID NO.: 5 and 6.

15**20**

A person skilled in the art knows how to obtain a chromosome segment, e.g. by cloning chromosome fragments into BACs, s. for example Song, 2003, PNAS, 100, 16, 9128 or as described herein and in the references cited herein.

25

Accordingly, in a further embodiment, the polynucleotide of the present invention or a polynucleotide encoding the Rpi-blb2 protein co-segregates with a marker selected from table 3a or comprises a replication site or hybridization site for said marker. As described in detail in the examples, the resistance to *P. infestans* could be mapped with the markers depicted in table 3a or 3b. As closer a marker is localized to a gene, as higher is the percentage of lines in which the marker co-s segregates with said marker. Therefore in a preferred embodiment, the polynucleotide of the present invention co-s segregates with the Marker E40M58, CT119 and/or CT216.

30

In a further embodiment, the present invention relates to a method for making a recombinant vector comprising inserting the polynucleotide of the present invention into a vector or inserting said polynucleotide and a further resistance protein into a vector.

35

Accordingly, in one further embodiment, the present invention relates to a vector containing the polynucleotide of the present invention or said polynucleotide and a further resistance gene or produced by the method of the present invention.

40

As used herein, the term "vector" refers to a nucleic acid molecule capable of transporting a polynucleotide to which it has been linked. One type of vector is a "plasmid", which refers to a circular double stranded DNA loop into which additional DNA seg-

Agrico B.V.

20030596

PF 54801

28

ments can be ligated. Another type of vector is a viral vector, wherein additional DNA or PNA segments can be ligated into the viral genome. Certain vectors are capable of autonomous replication in a host cell into which they are introduced (e.g., bacterial vectors having a bacterial origin of replication and episomal mammalian vectors). Other
5 vectors (e.g., non-episomal mammalian vectors) are integrated into the genome of a host cell upon introduction into the host cell, and thereby are replicated along with the host genome. Moreover, certain vectors are capable of directing the expression of genes to which they are operatively linked. Such vectors are referred to herein as "expression vectors". In general, expression vectors of utility in recombinant DNA tech-
10 niques are often in the form of plasmids. In the present specification, "plasmid" and "vector" can be used interchangeably as the plasmid is the most commonly used form of vector. However, the invention is intended to include such other forms of expression vectors, such as viral vectors (e.g., replication defective retroviruses, adenoviruses and adeno-associated viruses), which serve equivalent functions.

15 The present invention also relates to cosmids, viruses, bacteriophages and other vectors used conventionally in genetic engineering that contain a nucleic acid molecule according to the invention. Methods which are well known to those skilled in the art can be used to construct various plasmids and vectors; see, for example, the techniques
20 described in Sambrook, Molecular Cloning A Laboratory Manual, Cold Spring Harbor Laboratory (1989) N.Y. and Ausubel, Current Protocols in Molecular Biology, Green Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, the nucleic acid molecules and vectors of the invention can be reconstituted into liposomes for delivery to target cells.

25 In another embodiment, the vector of the present invention or the method of the present invention the vector or the method is characterized therein, that the polynucleotide encoding Rpi-blb2 protein or a further resistance protein is operatively linked to expres-
30 sion control sequences and/or a linked to a nucleic acid sequence encoding a transgenic expression regulating signal allowing expression in prokaryotic or eukaryotic host cells.

In a preferred embodiment, the present invention relates to a vector of the present in-
35 vention or the method of the present invention in which the polynucleotide encoding Rpi-blb2 protein and/or the further resistance protein is operatively linked to expression control sequences of the same species origin as the polynucleotide encoding Rpi-blb2 protein and/or the further resistance protein.

In the case that a nucleic acid molecule according to the invention is expressed in a
40 cell it is in principle possible to modify the coding sequence in such a way that the protein is located in any desired compartment of the plant cell. These include the nucleus,

Agrico B.V.

20030596

PF 54801

29

endoplasmatic reticulum, the vacuole, the mitochondria, the plastids like amyloplasts, chloroplasts, chromoplasts, the apoplast, the cytoplasm, extracellular space, oil bodies, peroxisomes and other compartments of plant cells (for review see Kermode, Crit. Rev. Plant Sci. 15, 4 (1996), 285-423 and references cited therein). The polynucleotide can then operatively be fused to an appropriate polynucleotide, e.g., a vector, encoding a signal for the transport into the desirable compartment.

In an other preferred embodiment of the present invention relates to a vector in which the polynucleotide of the present invention is operatively linked to expression control sequences allowing expression in prokaryotic or eukaryotic host cells. The nature of such control sequences differs depending upon the host organism. In prokaryotes, control sequences generally include promoter, ribosomal binding site, and terminators. In eukaryotes, generally control sequences include promoters, terminators and, in some instances, enhancers, transactivators; or transcription factors.

The term "control sequence" is intended to include, at a minimum, components the presence of which are necessary for expression, and may also include additional advantageous components.

The term "operably linked" refers to a juxtaposition wherein the components so described are in a relationship permitting them to function in their intended manner. A control sequence "operably linked" to a coding sequence is ligated in such a way that expression of the coding sequence is achieved under conditions compatible with the control sequences. In case the control sequence is a promoter, it is obvious for a skilled person that double-stranded nucleic acid is used.

Operable linkage is to be understood as meaning, for example, the sequential arrangement of a promoter with the nucleic acid sequence to be expressed and, if appropriate, further regulatory elements such as, for example, a terminator in such a way that each of the regulatory elements can fulfil its function when the nucleic acid sequence is expressed recombinantly, depending on the arrangement of the nucleic acid sequences in relation to sense or antisense RNA. To this end, direct linkage in the chemical sense is not necessarily required. Genetic control sequences such as, for example, enhancer sequences, can also exert their function on the target sequence from positions which are further away, or indeed from other DNA molecules. Preferred arrangements are those in which the nucleic acid sequence to be expressed recombinantly is positioned behind the sequence acting as promoter, so that the two sequences are linked covalently to each other. The distance between the promoter sequence and the nucleic acid sequence to be expressed recombinantly is preferably less than 200 base pairs, especially preferably less than 100 base pairs, very especially preferably less than 50 base pairs.

Agrico B.V.

20030596

PF 54801

30

Operable linkage, and an expression cassette, can be generated by means of customary recombination and cloning techniques as are described, for example, in Maniatis T, Fritsch EF and Sambrook J (1989) *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory, Cold Spring Harbor (NY), in Silhavy TJ, Berman ML and Enquist LW (1984) *Experiments with Gene Fusions*, Cold Spring Harbor Laboratory, Cold Spring Harbor (NY), in Ausubel FM et al. (1987) *Current Protocols in Molecular Biology*, Greene Publishing Assoc. and Wiley Interscience and in Gelvin et al. (1990) *In: Plant Molecular Biology Manual*. However, further sequences which, for example, act as a linker with specific cleavage sites for restriction enzymes, or as a signal peptide, may also be positioned between the two sequences. The insertion of sequences may also lead to the expression of fusion proteins. Preferably, the expression cassette, consisting of a linkage of promoter and nucleic acid sequence to be expressed, can exist in a vector-integrated form and be inserted into a plant genome, for example by transformation.

15

Such regulatory sequences are described, for example, in Goeddel; *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990) or see: Gruber and Crosby, in: *Methods in Plant Molecular Biology and Biotechnology*, CRC Press, Boca Raton, Florida, eds.: Glick and Thompson, Chapter 7, 89-108 including the references therein. Regulatory sequences include those which direct constitutive expression of a nucleotide sequence in many types of host cell and those which direct expression of the nucleotide sequence only in certain host cells or under certain conditions. It will be appreciated by those skilled in the art that the design of the expression vector can depend on such factors as the choice of the host cell to be transformed, the level of expression of protein desired, etc. The expression vectors of the invention can be introduced into host cells to thereby produce proteins or peptides, including fusion proteins or peptides, encoded by polynucleotides as described herein.

The recombinant expression vectors of the invention can be designed for expression of said resistance proteins, preferably Rpi-blb2, in prokaryotic or eukaryotic cells. For example, genes encoding the polynucleotide of the invention can be expressed in bacterial cells such as *E. coli*, *C. glutamicum*, *Agrobacterium tumefaciens*, insect cells (using baculovirus expression vectors), yeast and other fungal cells (see Romanos, (1992), *Yeast* 8: 423-488; van den Hondel, (1991) J.W. Bennet & L.L. Lasure, eds., p. 396-428; Academic Press: San Diego; and van den Hondel, (1991) in: *Applied Molecular Genetics of Fungi*, Peberdy, eds., p. 1-28, Cambridge University Press: Cambridge), algae (Falcatore et al., 1999, *Marine Biotechnology*, 1, 3:239-251), and multicellular plant cells (see Schmidt, R. (1988), *Plant Cell Rep.*: 583-586); *Plant Molecular Biology and Biotechnology*, C Press, Boca Raton, Florida, chapter 6/7, S.71-119 (1993); F.F. White, B. Jené et al., *Techniques for Gene Transfer*, in: *Transgenic*

Plants, Vol. 1, Engineering and Utilization, eds.: Kung und R. Wu, Academic Press (1993), 128-43; Potrykus, *Annu. Rev. Plant Physiol. Plant Molec. Biol.* 42 (1991), 205-225 (and references cited therein) or mammalian cells. Suitable host cells are discussed further in Goëddel, *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, CA (1990). Alternatively, the recombinant expression vector can be transcribed and translated *in vitro*, for example using T7 promoter regulatory sequences and T7 polymerase.

Expression of proteins in prokaryotes is most often carried out with vectors containing constitutive or inducible promoters directing the expression of either fusion or non-fusion proteins. Fusion vectors add a number of amino acids to a protein encoded therein, usually to the amino terminus of the recombinant protein but also to the C-terminus or fused within suitable regions in the proteins. Such fusion vectors typically serve three purposes: 1) to increase expression of recombinant protein; 2) to increase the solubility of the recombinant protein; and 3) to aid in the purification of the recombinant protein by acting as a ligand in affinity purification. Further, the fusion vector can also encode for additional proteins, which expression supports an increase of the activity of Rpi-b1b2 or of the resistance of a plant against plant pathogens, e.g. other resistance proteins. Often, in fusion expression vectors, a proteolytic cleavage site is introduced at the junction of the fusion moiety and the recombinant protein to enable separation of the recombinant protein from the fusion moiety subsequent to purification of the fusion protein. Such enzymes, and their cognate recognition sequences, include Factor Xa, thrombin and enterokinase.

Typical fusion expression vectors include pGEX (Pharmacia Biotech Inc; Smith, D.B. and Johnson, K.S. (1988) *Gene* 67:31-40), pMAL (New England Biolabs, Beverly, MA) and pRIT5 (Pharmacia, Piscataway, NJ).

Examples of suitable inducible non-fusion *E. coli* expression vectors include pTrc (Amann (1988) *Gene* 69:301-315) and pET 11d (Studier et al., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 60-89).

One strategy to maximize recombinant protein expression is to express the protein in a host bacteria with an impaired capacity to proteolytically cleave the recombinant protein (Gottesman, S., *Gene Expression Technology: Methods in Enzymology* 185, Academic Press, San Diego, California (1990) 119-128). Another strategy is to alter the nucleic acid sequence of the nucleic acid to be inserted into an expression vector so that the individual codons for each amino acid are those preferentially utilized in the bacterium chosen for expression, such as *E. coli* or *C. glutamicum* (Wada et al. (1992) *Nucleic Acids Res.* 20:2111-2118). Such alteration of nucleic acid sequences of the invention can be carried out by standard DNA synthesis techniques.

Agrico B.V.

20030596

PF 54801

32

Further, the vector can be a yeast expression vector. Examples of vectors for expression in yeast *S. cerevisiae* include pYepSec1 (Baldari, et al., (1987) *Embo J.* 6:229-234), pMFa (Kurjan and Herskowitz, (1982) *Cell* 30:933-943), pJRY88 (Schultz et al., (1987) *Gene* 54:113-123), and pYES2 (Invitrogen Corporation, San Diego, CA).

5

Preferably, the polynucleotide of the present invention or described herein is operatively linked to a plant expression control sequences, e.g. expression cassettes plant expression cassette preferably contains regulatory sequences capable to drive gene expression in plants cells and which are operably linked so that each sequence can fulfil its function such as termination of transcription such as polyadenylation signals. Preferred polyadenylation signals are those originating from *Agrobacterium tumefaciens* t-DNA such as the gene 3 known as octopine synthase of the Ti-plasmid pTiACH5 (Gielen et al., *EMBO J.* 3 (1984), 835 ff) or functional equivalents thereof but also all other terminators functionally active in plants are suitable.

As plant gene expression is very often not limited on transcriptional levels as plant expression cassette preferably contains other operably linked sequences like translational enhancers such as the overdrive-sequence containing the 5'-untranslated leader sequence from tobacco mosaic virus enhancing the protein per RNA ratio (Gallie et al 1987, *Nucl. Acids Research* 15:8693-8711).

20

Accordingly, the polynucleotide described herein can be operably linked to an appropriate promoter conferring gene expression in a timely, cell or tissue specific manner. Preferred are promoters driving constitutive expression (Benfey et al., *EMBO J.* 8 (1989) 2195-2202) like those derived from plant viruses like the 35S CaMV (Franck et al., *Cell* 21(1980) 285-294), the 19S CaMV (see also US5352605 and WO8402913) or plant promoters like those from Rubisco small subunit described in US 4962028.

25

The term plant-specific promoters is understood as meaning, in principle, any promoter which is capable of governing the expression of genes, in particular foreign genes, in plants or plant parts, plant cells, plant tissues or plant cultures. In this context, expression can be, for example, constitutive, inducible or development-dependent.

30

Agrico B.V.**20030596****PF 54801**

33

The following are preferred:

a) Constitutive promoters

- 5 Preferred vectors are those which make possible constitutive expression in plants (Benfey et al. (1989) EMBO J 8:2195-2202). "Constitutive" promoter is understood as meaning those promoters which ensure expression in a large number of, preferably all, tissues over a substantial period of plant development, preferably at all stages of plant development. In particular a plant promoter or a promoter derived from a plant virus are
- 10 preferably used. Particularly preferred is the promoter of the CaMV cauliflower mosaic virus 35S transcript (Franck et al. (1980) Cell 21:285-294; Odell et al. (1985) Nature 313:810-812; Shewmaker et al. (1985) Virology 140:281-288; Gardner et al. (1986) Plant Mol Biol 6:221-228) or the 19S CaMV promoter (US 5,352,605; WO 84/02913; Benfey et al. (1989) EMBO J 8:2195-2202). Another suitable constitutive promoter is
- 15 the "Rubisco small subunit (SSU)" promoter (US 4,962,028), the leguminB promoter (GenBank Acc. No. X03677), the Agrobacterium nopaline synthase promoter, the TR dual promoter, the Agrobacterium OCS (octopine synthase) promoter, the ubiquitin promoter (Holtorf S et al. (1995) Plant Mol Biol 29:637-649), the ubiquitin 1 promoter (Christensen et al. (1992) Plant Mol Biol 18:675-689; Bruce et al. (1989) Proc Natl
- 20 Acad Sci USA 86:9692-9696), the SmaS promoter, the cinnamyl alcohol dehydrogenase promoter (US 5,683,439), the promoters of the vacuolar ATPase subunits or the promoter of a proline-rich protein from wheat (WO 91/13991), and further promoters of genes whose constitutive expression in plants is known to the skilled worker.

25 b) Tissue-specific promoters

Preferred are furthermore promoters with specificity for the anthers, ovaries, flowers, leaves, stems, roots and seeds.

30 Seed-specific promoters

- such as, for example, the phaseolin promoter (US 5,504,200; Bustos MM et al. (1989) Plant Cell 1(9):839-53), the 2S albumin gene promoter (Joseffson LG et al. (1987) J Biol Chem 262:12196-12201), the legumin promoter (Shirsat A et al. (1989) Mol Gen Genet 215(2): 326-331), the USP (unknown seed protein) promoter (Bäumlein H et al. (1991) Mol Gen Genet 225(3):459-67), the napin gene promoter (US 5,608,152; Stalberg K et al. (1996) L Planta 199:515-519), the sucrose binding protein promoter (WO 00/26388) or the legumin B4 promoter (LeB4; Bäumlein H et al. (1991) Mol Gen Genet 225: 121-128; Bäumlein et al. (1992) Plant Journal 2(2):233-9; Fiedler U et al. (1995) Biotechnology (NY) 13(10):1090), the Arabidopsis oleosin promoter
- 40 (WO 98/45461), the Brassica Bce4 promoter (WO 91/13980). Further suitable seed-

Agrico B.V.

20030596

PF 54801

34

specific promoters are those of the genes encoding the high-molecular-weight glutenin (HMWG), gliadin, branching enzyme, ADP glucose pyrophosphatase (AGPase) or starch synthase. Furthermore preferred are promoters which permit seed-specific expression in monocots such as maize, barley, wheat, rye, rice and the like. The following

5 can be employed advantageously: the promoter of the *lpt2* or *lpt1* gene (WO 95/15389, WO 95/23230) or the promoters described in WO 99/16890 (promoters of the hordein gene, the glutelin gene, the oryzin gene, the prolamin gene, the gliadin gene, the glutelin gene, the zein gene, the kasirin gene or the secalin gene).

- 10 Tuber-, storage-root- or root-specific promoters such as, for example, the patatin promoter class I (B33), the potato cathepsin D inhibitor promoter.

Leaf-specific promoters

- such as the potato cytosolic FBPase promoter (WO 97/05900), the Rubisco (ribulose-1,5-bisphosphate carboxylase) SSU (small subunit) promoter or the ST-LSI promoter
- 15 from potato (Stockhaus et al. (1989) EMBO J 8:2445-2451). Very especially preferred are epidermis-specific promoters such as, for example, the OXLP gene (oxalate-oxidase-like protein) promoter (Wei et al. (1998) Plant Mol. Biol. 36:101-112).

20 Flower-specific promoters

such as, for example, the phytoene synthase promoter (WO 92/16635) or the promoter of the *P-rr* gene (WO 98/22593).

Anther-specific promoters

- 25 such as the 5126 promoter (US 5,689,049, US 5,689,051), the glob-I promoter and the γ -zein promoter.

c) Chemically Inducible promoters

- 30 The expression cassettes can also comprise a chemically inducible promoter (review article: Gatz et al. (1997) Annu Rev Plant Physiol Plant Mol Biol 48:89-108), by which the expression of the exogenous gene in the plant at a particular point in time can be controlled. Such promoters such as, for example, the PRP1 promoter (Ward et al. (1993) Plant Mol Biol 22:361-366), a salicylic-acid-inducible promoter (WO 95/19443),
- 35 a benzenesulfonamide-inducible promoter (EP 0 388 186), a tetracyclin-inducible promoter (Gatz et al. (1992) Plant J 2:397-404), an abscisic-acid-inducible promoter (EP 0 335 528) or an ethanol- or cyclohexanone-inducible promoter (WO 93/21334) can likewise be used.

d) Stress- or pathogen-inducible promoters

Further preferred promoters are those which are induced by biotic or abiotic stress such as, for example, the pathogen-inducible promoter of the PRP1 gene (Ward et al. (1993) Plant Mol Biol 22:361-366), the tomato high-temperature-inducible hsp70 or hsp80 promoter (US 5,187,267), the potato low-temperature-inducible alpha-amylase promoter (WO 96/12814), the light-inducible PPDK promoter, or the wounding-induced pinII promoter (EP375091).

- 10 Pathogen-inducible promoters encompass those of genes which are induced as a consequence of infection by pathogens, such as, for example, genes of PR proteins, SAR proteins, β -1,3-glucanase, chitinase and the like (for example Redolfi et al. (1983) Neth J Plant Pathol 89:245-254; Uknes, et al. (1992) The Plant Cell 4:645-656; Van Loon (1985) Plant Mol Biol 4:111-116; Marineau et al. (1987) Plant Mol Biol 9:335-342; Mat-
15 ton et al. (1987) Molecular Plant-Microbe Interactions 2:325-342; Somssich et al. (1986) Proc Natl Acad Sci USA 83:2427-2430; Somssich et al. (1988) Mol Gen Genet- ics 2:93-98; Chen et al. (1996) Plant J 10:955-966; Zhang and Sing (1994) Proc Natl Acad Sci USA 91:2507-2511; Warner, et al. (1993) Plant J 3:191-201; Siebertz et al. (1989) Plant Cell 1:961-968(1989).
- 20 Also encompassed are wounding-inducible promoters such as that of the pinII gene (Ryan (1990) Ann Rev Phytopath 28:425-449; Duan et al. (1996) Nat Biotech 14:494-498), of the wun1 and wun2 gene (US 5,428,148), of the win1 and win2 gene (Stanford et al. (1989) Mol Gen Genet 215:200-208), of systemin (McGurl et al. (1992) Science 225:1570-1573), of the WIP1 gene (Rohmeier et al. (1993) Plant Mol Biol 22:783-792;
25 Eckelkamp et al. (1993) FEBS Letters 323:73-76), of the MPI gene (Corderok et al. (1994) The Plant J 6(2):141-150) and the like.

e) Development-dependent promoters

- 30 Further suitable promoters are, for example, fruit-maturation-specific promoters such as, for example, the tomato fruit-maturation-specific promoter (WO 94/21794, EP 409 625). Development-dependent promoters comprise partly the tissue-specific promoters, since individual tissues develop by nature in a development-dependent fashion.

35

- It can be advantageously that the polypeptide of the present invention is only active or has only an increased activity in the tissue which is transfected or penetrated by the pathogen mentioned herein. Especially preferred are constitutive promoters and leaf- and/or stem-specific, pathogen-inducible and epidermis-specific promoters, with patho-
40 gen-inducible and epidermis-specific promoters being most preferred. Also preferred is

Agrico B.V.

20030596

PF 54801

36

the natural promoter, which is e.g. comprised in the genomic fragment depicted in Seq. ID NO.: 5 and 6.

Furthermore, further promoters may be linked operably to the nucleic acid sequence to be expressed, which promoters make possible the expression in further plant tissues or in other organisms, such as, for example, *E. coli* bacteria. Suitable plant promoters are, in principle, all of the above-described promoters.

The term "genetic control sequences" is to be understood in the broad sense and refers to also all those sequences which have an effect on the materialization or the function of the expression cassette according to the invention. For example, genetic control sequences modify the transcription and translation in prokaryotic or eukaryotic organisms. Preferably, the expression cassettes according to the invention encompass the promoter with specificity for the embryonal epidermis and/or the flower 5'-upstream of the nucleic acid sequence in question to be expressed recombinantly, and 3'-downstream a terminator sequence as additional genetic control sequence and, if appropriate, further customary regulatory elements, in each case linked operably to the nucleic acid sequence to be expressed recombinantly.

Genetic control sequences also encompass further promoters, promoter elements or minimal promoters, all of which can modify the expression-governing properties. Thus, for example, the tissue-specific expression may additionally depend on certain stressors, owing to genetic control sequences. Such elements have been described, for example, for water stress, abscisic acid (Lam E and Chua NH, *J Biol Chem* 1991; 266(26): 17131 -17135) and heat stress (Schoffl F et al., *Molecular & General Genetics* 217(2-3):246-53, 1989).

Further advantageous control sequences are, for example, the Gram-positive promoters *amy* and *SPO2*, and the yeast or fungal promoters *ADC1*, *MFa*, *AC*, *P-60*, *CYC1*, *GAPDH*, *TEF*, *rp28*, *ADH*.

In principle, all natural promoters with their regulatory sequences like those mentioned above may be used for the method according to the invention. In addition, synthetic promoters may also be used advantageously.

Genetic control sequences furthermore also encompass the 5'-untranslated regions, introns or noncoding 3'-region of genes, such as, for example, the actin-1 intron, or the *Adh1-S* introns 1, 2 and 6 (general reference: *The Maize Handbook*, Chapter 116, Freeling and Walbot, Eds., Springer, New York (1994)). It has been demonstrated that they may play a significant role in the regulation of gene expression. Thus, it has been demonstrated that 5'-untranslated sequences can enhance the transient expression of

Agrico B.V.

20030596

PF 54801

37

heterologous genes. Examples of translation enhancers which may be mentioned are the tobacco mosaic virus 5' leader sequence (Gallie et al. (1987) Nucl Acids Res 15:8693-8711) and the like. Furthermore, they may promote tissue specificity (Rouster J et al. (1998) Plant J 15:435-440).

5

The expression cassette may advantageously comprise one or more of what are known as enhancer sequences, linked operably to the promoter, which make possible an increased recombinant expression of the nucleic acid sequence. Additional advantageous sequences, such as further regulatory elements or terminators, may also be inserted at the 3' end of the nucleic acid sequences to be expressed recombinantly. One or more copies of the nucleic acid sequences to be expressed recombinantly may be present in the gene construct.

10

Polyadenylation signals which are suitable as control sequences are plant polyadenylation signals, preferably those which essentially correspond to T-DNA polyadenylation signals from *Agrobacterium tumefaciens*, in particular gene 3' of the T-DNA (octopin synthase) of the Ti plasmid pTiACHS (Gielen et al. (1984) EMBO J 3:835 et seq.) or functional equivalents thereof. Examples of terminator sequences which are especially suitable are the OCS (octopin synthase) terminator and the NOS (nopal synthase) terminator.

15

20

Control sequences are furthermore to be understood as those which make possible homologous recombination or insertion into the genome of a host organism or which permit removal from the genome. In the case of homologous recombination, for example the natural promoter of a particular gene may be exchanged for a promoter with specificity for the embryonal epidermis and/or the flower. Methods such as the cre/lox technology permit a tissue-specific, if appropriate inducible, removal of the expression cassette from the genome of the host organism (Sauer B (1998) Methods. 14(4):381-92). In this method, specific flanking sequences (lox sequences), which later allow removal by means of cre recombinase, are attached to the target gene.

25

30

An expression cassette and the vectors derived from it may comprise further functional elements. The term functional element is to be understood in the broad sense and refers to all those elements which have an effect on the generation, amplification or function of the expression cassettes, vectors or transgenic organisms according to the invention. The following may be mentioned by way of example, but not by limitation:

35

a) Selection markers which confer a resistance to a metabolism inhibitor such as 2-deoxyglucose-6-phosphate (WO 98/45456), antibiotics or biocides, preferably herbicides, such as, for example, kanamycin, G 418, bleomycin or hygromycin,

40

or else phosphinothricin and the like. Especially preferred selection markers are those which confer resistance to herbicides. Examples which may be mentioned are: DNA sequences which encode phosphinothricin acetyl transferases (PAT) and which inactivate glutamin synthase inhibitors (bar and pat genes), 5-enol-pyruvylshikimate-3-phosphate synthase genes (EPSP synthase genes), which confer resistance to Glyphosate (N-(phosphonomethyl)glycine), the gox gene, which encodes Glyphosate-degrading enzymes (Glyphosate oxidoreductase), the deh gene (encoding a dehalogenase which inactivates dalapon), sulfonyl-urea- and imidazolinone-inactivating acetolactate synthases, and bxn genes, which encode bromoxynil-degrading nitrilase enzymes, the aasa gene, which confers resistance to the antibiotic spectinomycin, the streptomycin phosphotransferase (SPT) gene, which allows resistance to streptomycin, the neomycin phosphotransferase (NPTII) gene, which confers resistance to kanamycin or geneticidin, the hygromycin phosphotransferase (HPT) gene, which mediates resistance to hygromycin, the acetolactate synthase gene (ALS), which confers resistance to sulfonylurea herbicides (for example mutated ALS variants with, for example, the S4 and/or Hra mutation).

b) Reporter genes which encode readily quantifiable proteins and, via their color or enzyme activity, make possible an assessment of the transformation efficacy, the site of expression or the time of expression. Very especially preferred in this context are genes encoding reporter proteins (Schenborn E, Groskreutz D. Mol Biotechnol. 1999; 13(1):29-44) such as the green fluorescent protein (GFP) (Sheen et al.(1995) Plant Journal 8(5):777-784; Haseloff et al.(1997) Proc Natl Acad Sci USA 94(6):2122-2127; Reichel et al.(1996) Proc Natl Acad Sci USA 93(12):5888-5893; Tian et al. (1997) Plant Cell Rep 16:267-271; WO 97/41228; Chui WL et al. (1996) Curr Biol 6:325-330; Leffel SM et al. (1997) Biotechniques. 23(5):912-8), chloramphenicol transferase, a luciferase (Ow et al. (1986) Science 234:856-859; Millar et al. (1992) Plant Mol Biol Rep 10:324-414), the aequorin gene (Prasher et al. (1985) Biochem Biophys Res Commun 126(3):1259-1268), β -galactosidase, R locus gene (encoding a protein which regulates the production of anthocyanin pigments (red coloring) in plant tissue and thus makes possible the direct analysis of the promoter activity without addition of further auxiliary substances or chromogenic substrates; Dellaporta et al., In: Chromosome Structure and Function: Impact of New Concepts, 18th Stadler Genetics Symposium, 11:263-282, 1988), with β -glucuronidase being very especially preferred (Jefferson et al., EMBO J. 1987, 6, 3901-3907).

c) Origins of replication, which ensure amplification of the expression cassettes or vectors according to the invention in, for example, E. coli. Examples which may

Agrico B.V.**20030596****PF 54801**

39

be mentioned are ORI (origin of DNA replication), the pBR322 ori or the P15A ori (Sambrook et al.: Molecular Cloning. A Laboratory Manual, 2nd ed. Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989).

- 5 d) Elements which are necessary for Agrobacterium-mediated plant transformation, such as, for example, the right or left border of the T-DNA or the vir region.

10 To select cells which have successfully undergone homologous recombination, or else to select transformed cells, it is, as a rule, necessary additionally to introduce a selectable marker, which confers resistance to a biocide (for example herbicide), a metabolism inhibitor such as 2-deoxyglucose-6-phosphate (WO 98/45456) or an antibiotic to the cells which have successfully undergone recombination. The selection marker permits the selection of the transformed cells from untransformed ones (McCormick et al. (1986) Plant Cell Reports 5:81-84).

15 The introduction of an expression cassette according to the invention into an organism or cells, tissues, organs, parts or seeds thereof (preferably into plants or plant cells, tissue, organs, parts or seeds) can be effected advantageously using vectors which comprise the expression cassettes. The expression cassette can be introduced into the
20 vector (for example a plasmid) via a suitable restriction cleavage site. The plasmid formed is first introduced into E. coli. Correctly transformed E. coli are selected, grown, and the recombinant plasmid is obtained by the methods familiar to the skilled worker. Restriction analysis and sequencing may serve to verify the cloning step.

25 Further promoters for expression in specific plant parts are e.g. the napin-gene promoter from rapeseed (US5608152), the USP-promoter from Vicia faba (Baeumlein et al., Mol Gen Genet, 1991, 225 (3):459-67), the oleosin-promoter from Arabidopsis (WO9845461), the phaseolin-promoter from Phaseolus vulgaris (US5504200), the Bce4-promoter from Brassica (WO9113980) or the legumin B4 promoter (LeB4; Baeumlein et al., 1992, Plant Journal, 2 (2):233-9) as well as promoters conferring seed
30 specific expression in monocot plants like maize, barley, wheat, rye, rice etc. Suitable promoters to note are the lpt2 or lpt1-gene promoter from barley (WO9515389 and WO9523230) or those described in WO9916890 (promoters from the barley hordelin-gene, the rice glutelin gene, the rice oryzin gene, the rice prolamin gene, the wheat gliadin gene, wheat glutelin gene, the maize zein gene, the oat glutelin gene, the Sorghum kasirin-gene, the rye secalin gene).

40 Further, the polynucleotide of the invention can be cloned into the expression vector in an antisense orientation. That is, the DNA molecule is operatively linked to a regulatory sequence in a manner which allows for expression (by transcription of the DNA mole-

- cule) of an RNA molecule which is antisense to the mRNA encoded by the polynucleotide of the present invention. Regulatory sequences operatively linked to a nucleic acid cloned in the antisense orientation can be chosen which direct the continuous expression of the antisense RNA molecule in a variety of cell types, for instance viral promoters and/or enhancers, or regulatory sequences can be chosen which direct constitutive, tissue specific or cell type specific expression of antisense RNA. The antisense expression vector can be in the form of a recombinant plasmid, phagemid or attenuated virus in which antisense nucleic acid molecules are produced under the control of a high efficiency regulatory region, the activity of which can be determined by the cell type into which the vector is introduced. For a discussion of the regulation of gene expression using antisense genes see Weintraub, H. et al., *Antisense RNA as a molecular tool for genetic analysis, Reviews - Trends in Genetics*, Vol. 1(1) 1986 and Mol et al., 1990, *FEBS Letters* 268:427-430.
- 15 In one embodiment the present invention relates to a method of making a recombinant host cell comprising introducing the vector or the polynucleotide of the present invention or said vector or said polynucleotide and a vector for expressing a further resistance protein into a host cell.
- 20 Vector DNA can be introduced into prokaryotic or eukaryotic cells via conventional transformation or transfection techniques. As used herein, the terms "transformation" and "transfection", conjugation and transduction are intended to refer to a variety of art-recognized techniques for introducing foreign nucleic acid (e.g., DNA) into a host cell, including calcium phosphate or calcium chloride co-precipitation, DEAE-dextran-
- 25 mediated transfection, lipofection, natural competence, chemical-mediated transfer, or electroporation. Suitable methods for transforming or transfecting host cells including plant cells can be found in Sambrook, et al. (*Molecular Cloning: A Laboratory Manual*, 2nd, ed., Cold Spring Harbor Laboratory, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, 1989) and other laboratory manuals such as *Methods in Molecular Biology*, 1995, Vol. 44, *Agrobacterium protocols*, ed: Gartland and Davey, Humana Press, Totowa, New Jersey.
- 30 For stable transfection of eukaryotic cells, it is known that, depending upon the expression vector and transfection technique used, only a small fraction of cells may integrate the foreign DNA into their genome. In order to identify and select these integrants, a gene that encodes a selectable marker (e.g., resistance to antibiotics) is generally introduced into the host cells along with the gene of interest. Preferred selectable markers include those which confer resistance to drugs, such as G418, hygromycin and methotrexate or in plants that confer resistance towards a herbicide such as glyphosate or glufosinate. Nucleic acid encoding a selectable marker can be introduced
- 35
- 40

into a host cell on the same vector as that encoding the polypeptide of the present invention or can be introduced on a separate vector. Cells stably transfected with the introduced nucleic acid can be identified by, for example, drug selection (e.g., cells that have incorporated the selectable marker gene will survive, while the other cells die).

5

Further host cells can be produced which contain selection systems which allow for regulated expression of the introduced gene. For example, inclusion of the polynucleotide of the invention on a vector placing it under control of the lac operon permits expression of the polynucleotide only in the presence of IPTG. Such regulatory systems

10

are well known in the art.

Preferably, the introduced nucleic acid molecule is foreign to the host cell.

15

By "foreign" it is meant that the nucleic acid molecule is either heterologous with, respect to the host cell, this means derived from a cell or organism with a different genomic background, or is homologous with respect to the host cell but located in a different genomic environment than the naturally occurring counterpart of said nucleic acid molecule. This means that, if the nucleic acid molecule is homologous with respect to the host cell, it is not located in its natural location in the genome of said host cell, in particular it is surrounded by different genes. In this case the nucleic acid molecule may be either under the control of its own promoter or under the control of a heterologous promoter. The vector or nucleic acid molecule according to the invention which is present in the host cell may either be integrated into the genome of the host cell or it may be maintained in some form extrachromosomally. In this respect, it is also to be understood that the nucleic acid molecule of the invention can be used to restore or create a mutant gene via homologous recombination (Paszowski (ed.), Homologous Recombination and Gene Silencing in Plants. Kluwer Academic Publishers (1994)).

20

25

30

Accordingly, in another embodiment the present invention relates to a host cell genetically engineered with the polynucleotide of the invention or the vector of the invention, or said vector or said polynucleotide and a vector or a polynucleotide for expressing a further resistance protein.

35

The terms "host cell" and "recombinant host cell" are used interchangeably herein. It is understood that such terms refer not only to the particular subject cell but to the progeny or potential progeny of such a cell. Because certain modifications may occur in succeeding generations due to either mutation or environmental influences, such progeny may not, in fact, be identical to the parent cell, but are still included within the scope of the term as used herein.

40

Agrico B.V.

20030596

PF 54801

42

For example, a polynucleotide of the present invention can be introduced in bacterial cells, insect cells, fungal cells or mammalian cells (such as Chinese hamster ovary cells (CHO) or COS cells), algae, ciliates, plant cells or fungi. Suitable host cells are known to those skilled in the art. Preferred are *E. coli*, baculovirus, *Agrobacterium* or plant cells.

Further, the host cell can also be transformed such that further enzymes and proteins are (over)expressed which expression supports an increase of resistance of a plant to pathogens. Preferably, a further resistance genes is also expressed, preferably one or more genes are as mentioned herein is/are expressed. Most preferred is a coexpression of *Rpi-b1b2* and *Rpi-b1b*.

Further preferred are cells of one of herein mentioned plants, in particular, of one of the above-mentioned Solanaceae, most preferred are potato, tomato, petunia, tree tomato, pear melon or egg plant.

In another embodiment, the present invention relates to a process for the production of the polypeptide of the present invention, in particular of a protein having *Rpi-b1b2* activity comprising culturing the host cell of the invention and recovering the polypeptide encoded by said polynucleotide and expressed by the host cell from the culture or the cells.

The term "expression" means the production of a protein or nucleotide sequence in the cell. However, said term also includes expression of the protein in a cell-free system. It includes transcription into an RNA product, post-transcriptional modification and/or translation to a protein product or polypeptide from a DNA encoding that product, as well as possible post-translational modifications.

Depending on the specific constructs and conditions used, the protein may be recovered from the cells, from the culture medium or from both. For the person skilled in the art it is well known that it is not only possible to express a native protein but also to express the protein as fusion polypeptides or to add signal sequences directing the protein to specific compartments of the host cell, e.g., ensuring secretion of the protein into the culture medium; etc. Furthermore, such a protein and fragments thereof can be chemically synthesized and/or modified according to standard methods described, for example hereinbelow.

A host cell of the invention, such as a prokaryotic or eukaryotic host cell in culture, can be used to produce (i.e., express) the polypeptide encoded by the polynucleotide of the invention, preferably a polypeptide having *Rpi-b1b2* activity. An alternate method can be

Agrico B.V.

20030596

PF 54801

43

applied in addition in plants by the direct transfer of DNA into developing flowers via electroporation or Agrobacterium medium gene transfer. Accordingly, the invention further provides methods for producing Rpi-blb2 using the host cells of the invention. In one embodiment, the method comprises culturing the host cell of invention in a suitable medium such that the polypeptide of the present invention is produced. Further, the method comprises isolating recovering said polypeptide from the medium or the host cell.

The polypeptide of the present invention is preferably produced by recombinant DNA techniques. For example, a nucleic acid molecule encoding the protein is cloned into an expression vector (as described above), the expression vector is introduced into a host cell (as described above) and said polypeptide is expressed in the host cell. Said polypeptide can then be isolated from the cells by an appropriate purification scheme using standard protein purification techniques. Alternative to recombinant expression, the polypeptide or peptide of the present invention can be synthesized chemically using standard peptide synthesis techniques. Moreover, native Rpi-blb2 can be isolated from cells (e.g., endothelial cells), for example using the antibody of the present invention as described below, in particular, an anti-Rpi-blb2 antibody, which can be produced by standard techniques utilizing the polypeptide of the present invention or fragment thereof, i.e., the polypeptide of this invention.

In one embodiment, the present invention relates to a Rpi-blb2 protein or a protein having Rpi-blb2 activity.

In one embodiment, the present invention relates to a polypeptide having the amino acid sequence encoded by a polynucleotide of the invention or obtainable by a process of the invention.

The terms "protein" and "polypeptide" used in this application are interchangeable. "Polypeptide" refers to a polymer of amino acids (amino acid sequence) and does not refer to a specific length of the molecule. Thus peptides and oligopeptides are included within the definition of polypeptide. This term does also refer to or include post-translational modifications of the polypeptide, for example, glycosylations, acetylations, phosphorylations and the like. Included within the definition are, for example, polypeptides containing one or more analogs of an amino acid (including, for example, unnatural amino acids, etc.), polypeptides with substituted linkages, as well as other modifications known in the art, both naturally occurring and non-naturally occurring.

Preferably, the polypeptide is isolated. An "isolated" or "purified" protein or biologically active portion thereof is substantially free of cellular material when produced by re-

Agrico B.V.

20030596

PF 54801

44

combinant DNA techniques, or chemical precursors or other chemicals when chemically synthesized.

5 The language "substantially free of cellular material" includes preparations of the polypeptide of the invention in which the protein is separated from cellular components of the cells in which it is naturally or recombinantly produced. In one embodiment, the language "substantially free of cellular material" includes preparations having less than about 30% (by dry weight) of "contaminating protein", more preferably less than about 20% of "contaminating protein", still more preferably less than about 10% of "contaminating protein", and most preferably less than about 5% "contaminating protein". The term "Contaminating protein" relates to polypeptides which are not polypeptides of the present invention. When the polypeptide of the present invention or biologically active portion thereof is recombinantly produced, it is also preferably substantially free of culture medium, i.e., culture medium represents less than about 20%, more preferably less than about 10%, and most preferably less than about 5% of the volume of the protein preparation. The language "substantially free of chemical precursors or other chemicals" includes preparations in which the polypeptide of the present invention is separated from chemical precursors or other chemicals which are involved in the synthesis of the protein. The language "substantially free of chemical precursors or other chemicals" includes preparations having less than about 30% (by dry weight) of chemical precursors or non-Rpi-blb2 chemicals, more preferably less than about 20% chemical precursors or non-Rpi-blb2 chemicals, still more preferably less than about 10% chemical precursors or non-Rpi-blb2 chemicals, and most preferably less than about 5% chemical precursors or non-Rpi-blb2 chemicals. In preferred embodiments, isolated proteins or biologically active portions thereof lack contaminating proteins from the same organism from which the polypeptide of the present invention is derived. Typically, such proteins are produced by recombinant:

30 A polypeptide of the invention can participate in the polypeptide or portion thereof comprises preferably an amino acid sequence which is sufficiently homologous to an amino acid sequence of SEQ ID No: 2 or 4 such that the protein or portion thereof maintains the ability to confer the resistance of the present invention. The portion of the protein is preferably a biologically active portion as described herein. Preferably, the polypeptide of the invention has an amino acid sequence identical as shown in SEQ ID No: 2 or 4. Further, the polypeptide can have an amino acid sequence which is encoded by a nucleotide sequence which hybridizes, preferably hybridizes under stringent conditions as described above, to a nucleotide sequence of the polynucleotide of the present invention. Accordingly, the polypeptide has an amino acid sequence which is encoded by a nucleotide sequence that is at least about 70%, preferably at least about 75%, more preferably at least about 80%, 90%, 95%, and even more preferably at least about 40

Agrico B.V.**20030596****PF 54801**

45

96%, 97%, 98%, 99% or more homologous to one of the amino acid sequences of SEQ ID No: 2 or 4. The preferred polypeptide of the present invention preferably possess at least one of the Rpi-blb2 activities described herein, e.g. its resistance or immunological activities. A preferred polypeptide of the present invention includes an amino acid sequence encoded by a nucleotide sequence which hybridizes, preferably hybridizes under stringent conditions, to a nucleotide sequence of SEQ ID No: 1 or 3 or 5 or 6 or which is homologous thereto, as defined above.

Accordingly the polypeptide of the present invention can vary from SEQ ID No: 2, or 4 in amino acid sequence due to natural variation or mutagenesis, as described in detail herein. Accordingly, the polypeptide comprise an amino acid sequence which is at least about 70%, preferably at least about 75%, and more preferably at least about 80, 90, 95%, and most preferably at least about 96%, 97%, 98%, 99% or more homologous to an entire amino acid sequence of SEQ ID No:1 or 3 or 5 or 6.

Biologically active portions of an polypeptide of the present invention include peptides comprising amino acid sequences derived from the amino acid sequence of an Rpi-blb2, e.g., the amino acid sequence shown in SEQ ID No: 2 or 4 or the amino acid sequence of a protein homologous thereto, which include fewer amino acids than a full length Rpi-blb2 or the full length protein which is homologous to an Rpi-blb2 depicted herein, and exhibit at least one activity of Rpi-blb2. Typically, biologically (or immunologically) active portions i.e. peptides, e.g., peptides which are, for example, 5, 10, 15, 20, 30, 35, 36, 37, 38, 39, 40, 50, 100 or more amino acids in length comprise a domain or motif with at least one activity or epitope of an Rpi-blb2. Moreover, other biologically active portions, in which other regions of the polypeptide are deleted, can be prepared by recombinant techniques and evaluated for one or more of the activities described herein.

Manipulation of the Rpi-blb2 polynucleotide of the invention may result in the production of Rpi-blb2 having functional differences from the wild-type Rpi-blb2. These proteins may be improved in efficiency or activity, may be present in greater numbers in the cell than is usual, or may be decreased in efficiency or activity.

Any mutagenesis strategies for Rpi-blb2 to result in increased said resistance or a resistance to another plant pathogen species or an other strain of a plant pathogen species aforementioned, of said compound are not meant to be limiting; variations on these strategies will be readily apparent to one skilled in the art. Using such strategies, and incorporating the mechanisms disclosed herein, the polynucleotide and polypeptide of the invention may be utilized to generate plants or parts thereof, expressing wildtyp Rpi-blb2 or mutated Rpi-blb2 polynucleotide and protein molecules such that

Agrico B.V.

20030596

PF 54801

46

the yield, production, and/or efficiency of production of a desired compound is improved. This desired compound may be any natural product of plants, which includes the final products of biosynthesis pathways and intermediates of naturally-occurring metabolic pathways, as well as molecules which do not naturally occur in the metabolism of said cells, but which are produced by a said cells of the invention.

The invention also provides chimeric or fusion proteins.

As used herein, an "chimeric protein" or "fusion protein" comprises an polypeptide operatively linked to a non- Rpi-blb2 polypeptide.

An "Rpi-blb2 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to polypeptide having a Rpi-blb2, whereas a "non-Rpi-blb2 polypeptide" refers to a polypeptide having an amino acid sequence corresponding to a protein which is not substantially homologous to the Rpi-blb2, e.g., a protein which does not confer the resistance described herein, in particular does not confer resistance to *P. infestans* and which is derived from the same or a different organism.

Within the fusion protein, the term "operatively linked" is intended to indicate that the Rpi-blb2 polypeptide and the non-Rpi-blb2 polypeptide are fused to each other so that both sequences fulfill the proposed function addicted to the sequence used. The non-Rpi-blb2 polypeptide can be fused to the N-terminus or C-terminus of the Rpi-blb2 polypeptide. For example, in one embodiment the fusion protein is a GST-LMRP fusion protein in which the Rpi-blb2 sequences are fused to the C-terminus of the GST sequences. Such fusion proteins can facilitate the purification of recombinant Rpi-blb2. In another embodiment, the fusion protein is an Rpi-blb2 containing a heterologous signal sequence at its N-terminus. In certain host cells (e.g., mammalian host cells), expression and/or secretion of an Rpi-blb2 can be increased through use of a heterologous signal sequence.

Preferably, an Rpi-blb2 chimeric or fusion protein of the invention is produced by standard recombinant DNA techniques. For example, DNA fragments coding for the different polypeptide sequences are ligated together in-frame in accordance with conventional techniques, for example by employing blunt-ended or stagger-ended termini for ligation, restriction enzyme digestion to provide for appropriate termini, filling-in of cohesive ends as appropriate, alkaline phosphatase treatment to avoid undesirable joining, and enzymatic ligation. The fusion gene can be synthesized by conventional techniques including automated DNA synthesizers. Alternatively, PCR amplification of gene fragments can be carried out using anchor primers which give rise to complementary overhangs between two consecutive gene fragments which can subsequently be annealed and reamplified to generate a chimeric gene sequence (see, for example, *Current Protocols in Molecular Biology*, eds. Ausubel et al. John Wiley & Sons: 1992).

Moreover, many expression vectors are commercially available that already encode a fusion moiety (e.g., a GST polypeptide). The polynucleotide of the invention can be cloned into such an expression vector such that the fusion moiety is linked in-frame to the encoded protein.

5

Furthermore, folding simulations and computer redesign of structural motifs of the protein of the invention can be performed using appropriate computer programs (Ol-
szewski, *Proteins* 25 (1996), 286-299; Hoffman, *Comput. Appl. Biosci.* 11 (1995), 675-
679). Computer modeling of protein folding can be used for the conformational and

10

energetic analysis of detailed peptide and protein models (Monge, *J. Mol. Biol.* 247
(1995), 995-1012; Renouf, *Adv. Exp. Med. Biol.* 376 (1995), 37-45). In particular, the
appropriate programs can be used for the identification of interactive sites of mitogenic
cyclin and its receptor, its ligand or other interacting proteins by computer assistant
searches for complementary peptide sequences (Fassina, *Immunomethods* (1994),

15

114-120. Further appropriate computer systems for the design of protein and peptides
are described in the prior art, for example in Berry, *Biochem. Soc. Trans.* 22 (1994),
1033-1036; Wodak, *Ann. N. Y. Acad. Sci.* 501 (1987), 1-13; Pabo, *Biochemistry* 25
(1986), 5987-5991. The results obtained from the above-described computer analysis

20

can be used for, e.g., the preparation of peptidomimetics of the protein of the invention
or fragments thereof. Such pseudopeptide analogues of the, natural amino acid se-
quence of the protein may very efficiently mimic the parent protein (Benkirane, *J. Biol.*
Chem. 271 (1996), 33218-33224). For example, incorporation of easily available
achiral Q-amino acid residues into a protein of the invention or a fragment thereof re-
sults in the substitution of amide bonds by polymethylene units of an aliphatic chain,

25

thereby providing a convenient strategy for constructing a peptidomimetic (Banerjee,
Biopolymers 39 (1996), 769-777).

Superactive peptidomimetic analogues of small peptide hormones in other systems are
described in the prior art (Zhang, *Biochem. Biophys. Res. Commun.* 224 (1996), 327-
331). Appropriate peptidomimetics of the protein of the present invention can also be

30

identified by the synthesis of peptidomimetic combinatorial libraries through successive
amide alkylation and testing the resulting compounds, e.g., for their binding and immu-
nological properties. Methods for the generation and use of peptidomimetic combinato-
rial libraries are described in the prior art, for example in Ostresh, *Methods in Enzymol-*
ogy 267 (1996), 220-234 and Dorner, *Bioorg. Med. Chem.* 4 (1996), 709-715.

35

Furthermore, a three-dimensional and/or crystallographic structure of the protein of
the invention can be used for the design of peptidomimetic inhibitors of the biological
activity of the protein of the invention (Rose, *Biochemistry* 35 (1996), 12933-12944;
Rutenber, *Bioorg. Med. Chem.* 4 (1996), 1545-1558).

Agrico B.V.

20030596

PF 54801

48

In a further embodiment, the present invention relates to an antibody that binds specifically to the polypeptide of the present invention or parts, i.e. specific fragments or epitopes of such a protein.

- 5 The antibodies of the invention can be used to identify and isolate Rpi-blb2 and genes in any organism, preferably plants, prepared in plants described herein. These antibodies can be monoclonal antibodies, polyclonal antibodies or synthetic antibodies as well as fragments of antibodies, such as Fab, Fv or scFv fragments etc. Monoclonal antibodies can be prepared, for example, by the techniques as originally described in Köhler and Milstein, Nature 256 (1975), 495, and Galfré, Meth. Enzymol. 73 (1981), 3,
- 10 which comprise the fusion of mouse myeloma cells to spleen cells derived from immunized mammals.

- Furthermore, antibodies or fragments thereof to the aforementioned peptides can be obtained by using methods which are described, e.g., in Harlow and Lane "Antibodies, A Laboratory Manual", CSH Press, Cold Spring Harbor, 1988. These antibodies can be
- 15 used, for example, for the immunoprecipitation and immunolocalization of proteins according to the invention as well as for the monitoring of the synthesis of such proteins, for example, in recombinant organisms, and for the identification of compounds interacting with the protein according to the invention. For example, surface plasmon resonance as employed in the BIAcore system can be used to increase the efficiency of
- 20 phage antibodies selections, yielding a high increment of affinity from a single library of phage antibodies which bind to an epitope of the protein of the invention (Schier, Human Antibodies Hybridomas 7 (1996), 97-105; Malmberg, J. Immunol. Methods 183 (1995), 7-13). In many cases, the binding phenomena of antibodies to antigens is equivalent to other ligand/anti-ligand binding.

- 25 In one embodiment, the present invention relates to an antisense nucleic acid molecule comprising the complementary sequence of the polypeptide of the present invention. Methods to modify the expression levels and/or the activity are known to persons skilled in the art and include for instance overexpression, co-suppression, the use of
- 30 ribozymes, sense and anti-sense strategies, gene silencing approaches. "Sense strand" refers to the strand of a double-stranded DNA molecule that is homologous to a mRNA transcript thereof. The "anti-sense strand" contains an inverted sequence which is complementary to that of the "sense strand".

- An "antisense" nucleic acid molecule comprises a nucleotide sequence which is complementary to a "sense" nucleic acid molecule encoding a protein, e.g., complementary
- 35 to the coding strand of a double-stranded cDNA molecule or complementary to an mRNA sequence. Accordingly, an antisense nucleic acid molecule can hydrogen bond to a sense nucleic acid molecule. The antisense nucleic acid molecule can be complementary to an entire Rpi-blb2 coding strand, or to only a portion thereof. Accordingly,
- 40 an antisense nucleic acid molecule can be antisense to a "coding region" of the coding

strand of a nucleotide sequence of a polynucleotide of the present invention. The term "coding region" refers to the region of the nucleotide sequence comprising codons which are translated into amino acid residues. Further, the antisense nucleic acid molecule is antisense to a "noncoding region" of the coding strand of a nucleotide sequence encoding Rpi-blb2. The term "noncoding region" refers to 5' and 3' sequences which flank the coding region that are not translated into a polypeptide, i.e., also referred to as 5' and 3' untranslated regions (5'-UTR or 3'-UTR).

Given the coding strand sequences encoding Rpi-blb2 disclosed herein, antisense nucleic acid molecules of the invention can be designed according to the rules of Watson and Crick base pairing. The antisense nucleic acid molecule can be complementary to the entire coding region of Rpi-blb2 mRNA, but can also be an oligonucleotide which is antisense to only a portion of the coding or noncoding region of Rpi-blb2 mRNA. For example, the antisense oligonucleotide can be complementary to the region surrounding the translation start site of Rpi-blb2 mRNA. An antisense oligonucleotide can be, for example, about 5, 10, 15, 20, 25, 30, 35, 40, 45 or 50 nucleotides in length. An antisense nucleic acid molecule of the invention can be constructed using chemical synthesis and enzymatic ligation reactions using procedures known in the art. For example, an antisense nucleic acid molecule (e.g., an antisense oligonucleotide) can be chemically synthesized using naturally occurring nucleotides or variously modified nucleotides designed to increase the biological stability of the molecules or to increase the physical stability of the duplex formed between the antisense and sense nucleic acids, e.g., phosphorothioate derivatives and acridine substituted nucleotides can be used. Examples of modified nucleotides which can be used to generate the antisense nucleic acid include 5-fluorouracil, 5-bromouracil, 5-chlorouracil, 5-iodouracil, hypoxanthine, xanthine, 4-acetylcytosine, 5-(carboxyhydroxymethyl) uracil, 5-carboxymethylaminomethyl-2-thiouridine, 5-carboxymethylaminomethyluracil, dihydrouracil, beta-D-galactosylqueosine, inosine, N6-isopentenyladenine, 1-methylguanine, 1-methylinosine, 2,2-dimethylguanine, 2-methyladenine, 2-methylguanine, 3-methylcytosine, 5-methylcytosine, N6-adenine, 7-methylguanine, 5-methylaminomethyluracil, 5-methoxyaminomethyl-2-thiouracil, beta-D-mannosylqueosine, 5'-methoxycarboxymethyluracil, 5-methoxyuracil, 2-methylthio-N6-isopentenyladenine, uracil-5-oxyacetic acid (v), wybutoxosine, pseudouracil, queosine, 2-thiocytosine, 5-methyl-2-thiouracil, 2-thiouracil, 4-thiouracil, 5-methyluracil, uracil-5-oxyacetic acid methylester, uracil-5-oxyacetic acid (v), 5-methyl-2-thiouracil, 3-(3-amino-3-N-2-carboxypropyl) uracil, (acp3)w, and 2,6-diaminopurine. Alternatively, the antisense nucleic acid can be produced biologically using an expression vector into which a polynucleotide has been subcloned in an antisense orientation (i.e., RNA transcribed from the inserted polynucleotide will be of an antisense orientation to a target polynucleotide of interest, described further in the following subsection).

The antisense nucleic acid molecules of the invention are typically administered to a cell or generated *in situ* such that they hybridize with or bind to cellular mRNA and/or genomic DNA encoding an Rpi-blb2 to thereby inhibit expression of the protein, e.g., by inhibiting transcription and/or translation. The hybridization can be by conventional nucleotide complementarity to form a stable duplex, or, for example, in the case of an antisense nucleic acid molecule which binds to DNA duplexes, through specific interactions in the major groove of the double helix. The antisense molecule can be modified such that it specifically binds to a receptor or an antigen expressed on a selected cell surface, e.g., by linking the antisense nucleic acid molecule to a peptide or an antibody which binds to a cell surface receptor or antigen. The antisense nucleic acid molecule can also be delivered to cells using the vectors described herein. To achieve sufficient intracellular concentrations of the antisense molecules, vector constructs in which the antisense nucleic acid molecule is placed under the control of a strong prokaryotic, viral, or eukaryotic including plant promoters are preferred.

In a further embodiment, the antisense nucleic acid molecule of the invention can be an α -anomeric nucleic acid molecule. An α -anomeric nucleic acid molecule forms specific double-stranded hybrids with complementary RNA in which, contrary to the usual units, the strands run parallel to each other (Gaultier et al. (1987) *Nucleic Acids Res.* 15:6625-6641). The antisense nucleic acid molecule can also comprise a 2'-o-methyl-ribonucleotide (Inoue et al. (1987) *Nucleic Acids Res.* 15:6131-6148) or a chimeric RNA-DNA analogue (Inoue et al. (1987) *FEBS Lett.* 215:327-330).

Further the antisense nucleic acid molecule of the invention can be a ribozyme. Ribozymes are catalytic RNA molecules with ribonuclease activity which are capable of cleaving a single-stranded nucleic acid, such as an mRNA, to which they have a complementary region. Thus, ribozymes (e.g., hammerhead ribozymes (described in Haselhoff and Gerlach (1988) *Nature* 334:585-591)) can be used to catalytically cleave Rpi-blb2 mRNA transcripts to thereby inhibit translation of mRNA. A ribozyme having specificity for an Rpi-blb2 -encoding nucleic acid molecule can be designed based upon the nucleotide sequence of an Rpi-blb2 cDNA disclosed herein or on the basis of a heterologous sequence to be isolated according to methods taught in this invention.

For example, a derivative of a *Tetrahymena* L-19 IVS RNA can be constructed in which the nucleotide sequence of the active site is complementary to the nucleotide sequence to be cleaved in an encoding mRNA. See, e.g., Cech et al. U.S. Patent No. 4,987,071 and Cech et al. U.S. Patent No. 5,116,742. Alternatively, Rpi-blb2 mRNA can be used to select a catalytic RNA having a specific ribonuclease activity from a pool of RNA molecules. See, e.g., Bartel, D. and Szostak, J.W. (1993) *Science* 261:1411-1418.

The antisense molecule of the present invention comprises also a polynucleotide comprising a nucleotide sequences complementary to the regulatory region of an Rpi-blb2

Agrico B.V.

20030596

PF 54801

51

nucleotide sequence, e.g., its promoter and/or enhancers, e.g. to form triple helical structures that prevent transcription of the gene in target cells. See generally, Helene, C. (1991) *Anticancer Drug Des.* 6(6):569-84; Helene, C. et al. (1992) *Ann. N.Y. Acad. Sci.* 660:27-36; and Maher, L.J. (1992) *Bioassays* 14(12):807-15.

5

In addition, in one embodiment, the present invention relates to a method for the production of transgenic plants, plant cells or plant tissue comprising the introduction of the polynucleotide or the vector of the present invention into the genome of said plant, plant tissue or plant cell. In a preferred embodiment, said vector or said polynucleotide and a vector or a polynucleotide for the expression of a further resistance gene, in particular for Rpi-blb, is also introduced into the genome of said plant, plant tissue or plant cell, before, after or together.

10

For the expression of the nucleic acid molecules according to the invention in sense or antisense orientation in plant cells, the molecules are placed under the control of regulatory elements which ensure the expression in plant cells. These regulatory elements may be heterologous or homologous with respect to the nucleic acid molecule to be expressed as well with respect to the plant species to be transformed and are described above in detail.

15

In general, such regulatory elements comprise a promoter active in plant cells. To obtain expression in all tissues of a transgenic plant, e.g. constitutive promoters are used, such as the 35 S promoter of CaMV (Odell, *Nature* 313 (1985), 810-812) or promoters of the polyubiquitin genes of maize (Christensen, *Plant Mol. Biol.* 18 (1982), 675-689). In order to achieve expression in specific tissues of a transgenic plant it is possible to use tissue specific promoters (see, e.g., Stockhaus, *EMBO J.* 8 (1989), 2245-2251). Known are also promoters which are specifically active in tubers of potatoes or in seeds of different plants species, such as maize, Vicia, wheat, barley etc. Inducible promoters may be used in order to be able to exactly control expression. Inducible promoters comprise also promoters, which are induced by infections of plants. Further embodiments are described above.

20

25

30

In one embodiment, the present invention relates to a method for producing a plant or a part thereof resistant to a pathogen of the phylum Oomyceta comprising the steps: expressing in the plant or a part thereof the polypeptide of the present invention and a further resistance protein.

35

Accordingly in one further embodiment, the present invention relates to transgenic plant or plant tissue of the invention or produced according to the method of the invention, which upon the presence of the polynucleotide or the vector is resistant to said pathogens.

Agrico B.V.

20030596

PF 54801

52

The generation of a transformed organism (or of a transformed cell or tissue) requires introducing the DNA, RNA or protein in question into the relevant host cell.

A multiplicity of methods are available for this procedure, which is termed transformation (or transduction or transfection) (Keown et al. (1990) *Methods in Enzymology*

- 5 185:527-537). For example, the DNA or RNA can be introduced directly by microinjection or by bombardment with DNA-coated microparticles. Also, the cell can be permeabilized chemically, for example using polyethylene glycol, so that DNA can enter the cell by diffusion. The DNA can also be introduced by protoplast fusion with other DNA-containing units such as minicells, cells, lysosomes or liposomes. Another suitable
- 10 method of introducing DNA is electroporation, where the cells are permeabilized reversibly by an electrical pulse. Suitable methods have been described (for example by Bilang et al. (1991) *Gene* 100:247-250; Scheid et al. (1991) *Mol Gen Genet* 228:104-112; Guerche et al. (1987) *Plant Science* 52:111-116; Neuhaus et al. (1987) *Theor Appl Genet* 75:30-36; Klein et al. (1987) *Nature* 327:70-73; Howell et al. (1980)
- 15 *Science* 208:1265; Horsch et al. (1985) *Science* 227:1229-1231; DeBlock et al. (1989) *Plant Physiology* 91:694-701; *Methods for Plant Molecular Biology* (Weissbach and Weissbach, eds.) Academic Press Inc. (1988); and *Methods in Plant Molecular Biology* (Schuler and Zielinski, eds.) Academic Press Inc. (1989)).

- 20 In plants, the above-described methods of transforming and regenerating plants from plant tissues or plant cells are exploited for transient or stable transformation. Suitable methods are especially protoplast transformation by polyethylene-glycol-induced DNA uptake, the biolistic method with the gene gun, what is known as the particle bombardment method, electroporation, incubation of dry embryos in DNA-containing solution, and microinjection.

- 25 In addition to these "direct" transformation techniques, transformation can also be effected by bacterial infection by means of *Agrobacterium tumefaciens* or *Agrobacterium rhizogenes*. The *Agrobacterium*-mediated transformation is best suited to dicotyledonous plant cells. The methods are described, for example, by Horsch et al. (1985)
- 30 *Science* 225: 1229f.

- When *agrobacteria* are used, the expression cassette must be integrated into specific plasmids, either into a shuttle or intermediate vector, or into a binary vector. If a Ti or Ri plasmid is to be used for the transformation, at least the right border, but in most cases the right and left border, of the Ti or Ri plasmid T-DNA is linked to the expression cassette to be introduced in the form of a flanking region.
- 35

Binary vectors are preferably used. Binary vectors are capable of replication both in *E. coli* and in *Agrobacterium*. As a rule, they comprise a selection marker gene and a linker or polylinker flanked by the right and left T-DNA border sequence. They can be

- 40 transferred directly into *Agrobacterium* (Holsters et al. (1978) *Mol Gen Genet* 163:181-

Agrico B.V.

20030596

PF 54801

53

187). The selection marker gene permits the selection of transformed agrobacteria and is, for example, the nptII gene, which confers resistance to kanamycin. The Agrobacterium which acts as host organism in this case should already contain a plasmid with the vir region. The latter is required for transferring the T-DNA to the plant cell. An Agrobacterium transformed in this way can be used for transforming plant cells. The use of T-DNA for transforming plant cells has been studied and described intensively (EP 120 516; Hoekema, In: The Binary Plant Vector System, Offsetdrukkerij Kanters B.V., Alblasserdam, Chapter V; An et al. (1985) EMBO J 4:277-287). Various binary vectors are known, some of which are commercially available such as, for example, pBI101.2 or pBIN19 (Clontech Laboratories, Inc. USA). Further promoters which are suitable for expression in plants have been described (Rogers et al. (1987) Meth in Enzymol 153:253-277; Schardl et al. (1987) Gene 61:1-11; Berger et al. (1989) Proc Natl Acad Sci USA 86:8402-8406).

15 Direct transformation techniques are suitable for any organism and cell type.

The plasmid used need not meet any particular requirements in the case of the injection or electroporation of DNA or RNA into plant cells. Simple plasmids such as those of the pUC series can be used. If complete plants are to be regenerated from the transformed cells, it is necessary for an additional selectable marker gene to be located on the plasmid.

Stably transformed cells, i.e. those which contain the introduced DNA integrated into the DNA of the host cell, can be selected from untransformed cells when a selectable marker is part of the DNA introduced. Examples of genes which can act as markers are all those which are capable of conferring resistance to antibiotics or herbicides (such as kanamycin, G 418, bleomycin, hygromycin or phosphinothricin) (see above). Transformed cells which express such marker genes are capable of surviving in the presence of concentrations of a corresponding antibiotic or herbicide which kill an untransformed wild type. Examples are mentioned above and preferably comprise the bar gene, which confers resistance to the herbicide phosphinothricin (Rathore KS et al. (1993) Plant Mol Biol 21(5):871-884), the nptII gene, which confers resistance to kanamycin, the hpt gene, which confers resistance to hygromycin, or the EPSP gene, which confers resistance to the herbicide Glyphosate. The selection marker permits the selection of transformed cells from untransformed cells (McCormick et al. (1986) Plant Cell Reports 5:81-84). The resulting plants can be bred and hybridized in the customary fashion. Two or more generations should be grown in order to ensure that the genomic integration is stable and hereditary.

The abovementioned methods are described, for example, in Jenes B et al. (1993) Techniques for Gene Transfer, in: Transgenic Plants, Vol. 1, Engineering and Utilization, edited by SD Kung and R Wu, Academic Press, pp. 128-143 and in Potrykus (1991) Annu Rev Plant Physiol Plant Molec Biol 42:205-225). The construct to be expressed is preferably cloned into a vector which is suitable for the transformation of *Agrobacterium tumefaciens*, for example pBin19 (Bevan et al. (1984) Nucl Acids Res 12:8711f).

As soon as a transformed plant cell has been generated, a complete plant can be obtained using methods known to the skilled worker. For example, callus cultures are used as starting material. The development of shoot and root can be induced in this as yet undifferentiated cell biomass in a known fashion. The shoots obtained can be planted out and bred.

The skilled worker is familiar with such methods of regenerating intact plants from plant cells and plant parts. Methods to do so are described, for example, by Fennell et al. (1992) Plant Cell Rep. 11: 567-570; Stoeger et al (1995) Plant Cell Rep. 14:273-278; Jahne et al. (1994) Theor Appl Genet 89:525-533.

The method according to the invention can advantageously be combined with further methods which bring about pathogen resistance (for example to insects, fungi, bacteria, nematodes and the like), stress resistance or another improvement of the plant properties. Examples are mentioned, inter alia, by Dunwell JM, Transgenic approaches to crop improvement, J Exp Bot. 2000;51 Spec No; pages 487-96.

Suitable strains of *Agrobacterium tumefaciens* and vectors as well as transformation of *Agrobacteria* and appropriate growth and selection media are well known to those skilled in the art and are described in the prior art (GV31 01 (pMK90RK), Koncz, Mol. Gen. Genet. 204 (1986), 383-396; C58C1 (pGV 3850kan), Deblaere, Nucl. Acid Res. 13 (1985), 4777; Bevan, Nucleic. Acid Res. 12(1984), 8711; Koncz, Proc. Natl. Acad. Sci. USA 86 (1989), 8467-8471; Koncz, Plant Mol. Biol. 20 (1992), 963-976; Koncz, Specialized vectors for gene tagging and expression studies. In: Plant Molecular Biology Manual Vol 2, Gelvin and Schilperoort (Eds.), Dordrecht, The Netherlands: Kluwer Academic Publ. (1994), 1-22; EP-A-120 516; Hoekema: The Binary Plant Vector System, Offsetdrukkerij Kanters B.V., Alblasserdam (1985), Chapter V, Fraley; Crit. Rev. Plant. Sci., 4, 1-46; An, EMBO J. 4 (1985), 277-287).

Although the use of *Agrobacterium tumefaciens* is preferred in the method of the invention, other *Agrobacterium* strains, such as *Agrobacterium rhizogenes*, may be used, for example if a phenotype conferred by said strain is desired.

The transformation of most dicotyledonous plants is possible with the methods described above. But also for the transformation of monocotyledonous plants several successful transformation techniques have been developed. These include the transformation using biolistic methods as, e.g., described above as well as protoplast transformation, electroporation of partially permeabilized cells, introduction of DNA using glass fibers, etc.

The term "transformation" as used herein, refers to the transfer of an exogenous polynucleotide into a host cell, irrespective of the method used for the transfer. The polynucleotide may be transiently or stably introduced into the host cell and may be maintained non-integrated, for example, as a plasmid or as chimeric links, or alternatively, may be integrated into the host genome. The resulting transformed plant cell can then be used to regenerate a transformed plant in a manner known by a skilled person.

Accordingly, in one embodiment, the present invention relates to a plant cell comprising the polynucleotide the vector of the present invention or obtainable by the method of the present invention. Preferably, the cell comprises a further resistance conferring polynucleotide or vector, more preferred is a Rpi-blb encoding vector or polynucleotide.

Thus, the present invention relates also to transgenic plant cells which contain (preferably stably integrated into the genome) a polynucleotide according to the invention linked to regulatory elements which allow expression of the polynucleotide in plant cells and wherein the polynucleotide is foreign to the transgenic plant cell. For the meaning of foreign; see supra.

Thus, the present invention also relates to transgenic plants and plant tissue comprising transgenic plant cells according to the invention. Due to the (over)expression of a polypeptide of the invention, said plant or plant tissues are resistance to plant pathogens, in particular to Oomyceta. Preferably the plants are also resistance to other pathogen, e.g. to sucking plant pathogens. Further pathogens are described herein. Preferred is that said plants or plant tissue is resistance to Phytophthora species, most preferred to P. Infestans.

For example, to obtain transgenic plants expressing the Rpi-blb2 gene, its coding region can be cloned, e.g., into the pBinAR vector (Höfgen und Willmitzer, Plant-Science, 66, 1990, 221-230). For example, following a polymerase chain reaction (PCR) technology the coding region of Rpi-blb2 can be amplified using Primers as shown in the examples and figures, e.g., in Table 3a in particular ARF1F and ARF1R. The obtained PCR fragment can be purified and subsequently the fragment can be cloned into a vector. The resulted vector can be transferred into Agrobacterium tumefaciens. This strain can be used to transform and transgenic plants can then be selected in another embodiment, the present invention relates to a transgenic plant or plant tissue comprising the plant cell of the present invention.

Agrico B.V.

20030596

PF 54801

56

"Transgenic". for example regarding a nucleic acid sequence, an expression cassette or a vector comprising said nucleic acid sequence or an organism transformed with said nucleic acid sequence, expression cassette or vector, refers to all those constructs originating by recombinant methods in which either

- a) the Rpi-blb2 nucleic acid sequence, or
- b) a genetic control sequence linked operably to the RacB nucleic acid sequence, for example a promoter, or
- c) (a) and (b)

are not located in their natural genetic environment or have been modified by recombinant methods, an example of a modification being a substitution, addition, deletion, inversion or insertion of one or more nucleotide residues. Natural genetic environment refers to the natural chromosomal locus in the organism of origin, or to the presence in a genomic library. In the case of a genomic library, the natural genetic environment of the nucleic acid sequence is preferably retained, at least in part. The environment flanks the nucleic acid sequence at least at one side and has a sequence of at least 50 bp, preferably at least 500 bp, especially preferably at least 1000 bp, very especially preferably at least 5000 bp, in length. A naturally occurring expression cassette - for example the naturally occurring combination of the Rpi-blb2 promoter with the corresponding Rpi-blb2 gene - becomes a transgenic expression cassette when it is modified by non-natural, synthetic "artificial" methods such as, for example, mutagenization. Such methods have been described (US 5,565,350; WO 00/15815; also see above).

Further, the plant cell, plant tissue or plant can also be transformed such that further enzymes and proteins are (over)expressed which expression supports an increase of the plant's or the plant tissue's resistance, for example Rpi-blb, R1, R-ber, Rpi1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11, Ph-1, Ph-2 and/or Ph-3-proteins. Preferred is the coexpression of Rpi-blb and Rpi-blb2.

The present invention also relates to cultured plant tissues comprising transgenic plant cells as described above which show expression of a protein according to the invention.

Host or starting organisms which are preferred as transgenic organisms are mainly plants in accordance with the above definition. Included within the scope of the invention are all genera and species of higher and lower plants of the Plant Kingdom.

Furthermore included are the mature plants, seed, shoots and seedlings, and parts, propagation material and cultures derived therefrom, for example cell cultures. Mature

plants refers to plants at any developmental stage beyond that of the seedling. The term seedling refers to a young immature plant in an early developmental stage.

- Any transformed plant obtained according to the invention can be used in a conventional breeding scheme or in in vitro plant propagation to produce more transformed plants with the same characteristics and/or can be used to introduce the same characteristic in other varieties of the same or related species. Such plants are also part of the invention. Seeds obtained from the transformed plants genetically also contain the same characteristic and are part of the invention. As mentioned before, the present invention is in principle applicable to any plant and crop that can be transformed with any of the transformation method known to those skilled in the art.

- In general, the plants which can be modified according to the invention and which either show overexpression of a protein according to the invention or a reduction of the synthesis of such a protein can be derived from any desired plant species. They can be monocotyledonous plants or dicotyledonous plants, preferably they belong to plant species of interest in agriculture, wood culture or horticulture interest, such as crop plants (e.g. maize, rice, barley, wheat, rye, oats etc.), potatoes, oil producing plants (e.g. oilseed rape, sunflower, pea nut, soy bean, etc.), cotton, sugar beet, sugar cane, leguminous plants (e.g. beans, peas etc.), wood producing plants, preferably trees, etc. However, plants which can be infected by Phytophthora species are preferred.

- Accordingly, in one embodiment the plant, plant cell or plant tissue of the invention or produced according to the method of the invention is selected from the group consisting of Menyanthaceae, Solanaceae, Sclerophyllaceae, Duckeodendraceae, Goetzeaceae, Convolvulaceae, Cuscutaceae, Polemoniaceae, and Hydrophyllaceae according to the Systeme Naturae 2000, Brands, S.J., Amsterdam or has its origin thereof. Preferably said plant, plant cell or plant tissue of the invention or produced according to the method of the invention is a Solanaceae, preferably selected from the group of Atropa, Browallia, Brunfelsia, Capsicum, Cestrum, Cyphomandra, Datura, Fabiana, Franciscea, Hyoscyamus, Lycium, Mandragora, Nicandra, Nicotiana, Petunia, Physalis, Schizanthus and Solanum according to the Systema Naturae 2000, Brands, S.J., Amsterdam or has its origin thereof.

- More preferred, the plant, plant cell or plant tissue of the invention or produced according to the method of the present invention is a *S. bulbocastanum*, *S. tuberosum* (potato), *S. lycopersicum*, petunia, *S. betaceum* (tree tomato), *S. muricatum* (pear melon) or *S. melongena* (eggplant). Even more preferred, the plant, plant tissue or plant cell is a *S. tuberosum* or *S. lycopersicum*. Most preferred is *S. tuberosum*. In other systems, the classification will be similar. The person skilled in the art knows the differences, e.g.

more common, tomato is named systematically *Lycopersicon Lycopersicum* (L.) Karsten ex Farwell.

5 In yet another aspect, the invention also relates to harvestable parts and to propagation material of the transgenic plants according to the invention which either contain transgenic plant cells expressing a nucleic acid molecule according to the invention or which contains cells which show a reduced level of the described protein.

10 Harvestable parts can be in principle any useful parts of a plant, for example, flowers, pollen, seedlings, tubers, leaves, stems, fruit, seeds, roots etc. Propagation material includes, for example, seeds, fruits, cuttings, seedlings, tubers, rootstocks etc. Preferred are potatoes, tomatoes, egg fruits or pear melons as harvestable or propagation material. In case, the plant of the invention is petunia, the present invention relates in one embodiment to the flowers of petunia as harvestable part.

15 The invention furthermore relates to the use of the transgenic organisms according to the invention and of the cells, cell cultures, parts - such as, for example, roots, leaves and the like in the case of transgenic plant organisms - derived from them, and to transgenic propagation material such as seeds or fruits, for the production of foodstuffs
20 or feeding stuffs, pharmaceuticals or fine chemicals. In particular, potatoes can serve for the production of fine chemicals.

Accordingly in another embodiment, the present invention relates to the use of the polynucleotide, the plant, plant cell or plant tissue, the vector, or the polypeptide of the
25 present invention for making fatty acids, carotenoids, isoprenoids, vitamins, lipids, wax esters, (poly)saccharides and/or polyhydroxyalkanoates, and/or its metabolism products, in particular, steroid hormones, cholesterol, prostaglandin, triacylglycerols, bile acids and/or ketone bodies producing cells, tissues and/or plants. There are a number of mechanisms by which the yield, production, and/or efficiency of production of fatty
30 acids, carotenoids, isoprenoids, vitamins, wax esters, lipids, (poly)saccharides and/or polyhydroxyalkanoates, and/or its metabolism products, in particular, steroid hormones, cholesterol, triacylglycerols, prostaglandin, bile acids and/or ketone bodies or further of above defined fine chemicals incorporating such an altered protein can be affected. In the case of plants, by e.g. increasing the expression of acetyl-CoA which is the basis
35 for many products, e.g., fatty acids, carotenoids, isoprenoids, vitamins, lipids, (poly)saccharides, wax esters, and/or polyhydroxyalkanoates, and/or its metabolism products, in particular, prostaglandin, steroid hormones, cholesterol, triacylglycerols, bile acids and/or ketone bodies in a cell, it may be possible to increase the amount of the produced said compounds thus permitting greater ease of harvesting and purification
40 or in case of plants more efficient partitioning. Further, one or more of said metabo-

lism products, increased amounts of the cofactors, precursor molecules, and intermediate compounds for the appropriate biosynthetic pathways maybe required. Therefore, by increasing the number and/or activity of transporter proteins involved in the import of nutrients, such as carbon sources (i.e., sugars), nitrogen sources (i.e., amino acids, ammonium salts), phosphate, and sulfur, it may be possible to improve the production of acetyl CoA and its metabolism products as mentioned above, due to the removal of any nutrient supply limitations on the biosynthetic process. In particular, it may be possible to increase the yield, production, and/or efficiency of production of said compounds, e.g. fatty acids, carotenoids, isoprenoids, vitamins, was esters, lipids, (poly)saccharides, and/or polyhydroxyalkanoates, and/or its metabolism products, in particular, steroid hormones, cholesterol, prostaglandin, triacylglycerols, bile acids and/or ketone bodies molecules etc. in plants.

Furthermore preferred is a method for the recombinant production of pharmaceuticals or fine chemicals in host organisms, wherein a host organism is transformed with one of the above-described expression cassettes and this expression cassette comprises one or more structural genes which encode the desired fine chemical or catalyze the biosynthesis of the desired fine chemical, the transformed host organism is cultured, and the desired fine chemical is isolated from the culture medium. This method can be applied widely to fine chemicals such as enzymes, vitamins, amino acids, sugars, fatty acids, and natural and synthetic flavorings, aroma substances and colorants. Especially preferred is the production of tocopherols and tocotrienols and carotenoids. The transformed host organisms are cultured and the products are isolated from the host organisms or the culture medium by methods known to the skilled worker. The production of pharmaceuticals such as, for example, antibodies or vaccines, is described by Hood EE, Jilka JM. Curr Opin Biotechnol. 1999 Aug; 10(4):382-6; Ma JK, Vine ND. Curr Top Microbiol Immunol. 1999; 236:275-92.

In one embodiment, the present invention also relates to the use of the polynucleotide, the vector, or the polypeptide of the present invention for producing a plant or a plant tissue, plant organ, or a plant cell or a part thereof resistant to said.

Furthermore, in one embodiment, the present invention relates to a method for the identification of an compound stimulating resistance to a said plant pathogen comprising:

- a) contacting cells which express the polypeptide of the present invention or its mRNA with a candidate compound under cell cultivation conditions;
- b) assaying an increase in expression of said polypeptide or said mRNA;

c) comparing the expression level to a standard response made in the absence of said candidate compound; whereby, an increased expression over the standard indicates that the compound is stimulating resistance.

- 5 Said compound may be chemically synthesized or microbiologically produced and/or comprised in, for example, samples, e.g., cell extracts from, e.g., plants, animals or microorganisms, e.g. pathogens. Furthermore, said compound(s) may be known in the art but hitherto not known to be capable of suppressing or activating Rpi-blb2. The reaction mixture may be a cell free extract or may comprise a cell or tissue culture. Suitable set ups for the method of the invention are known to the person skilled in the art and are, for example, generally described in Alberts et al., Molecular Biology of the Cell, third edition (1994), in particular Chapter 17. The compounds may be, e.g., added to the reaction mixture, culture medium, injected into the cell or sprayed onto the plant.
- 10
- 15 If a sample containing a compound is identified in the method of the invention, then it is either possible to isolate the compound from the original sample identified as containing the compound capable of activating or increasing resistance to said pathogens, or one can further subdivide the original sample, for example, if it consists of a plurality of different compounds, so as to reduce the number of different substances per sample and repeat the method with the subdivisions of the original sample. Depending on the complexity of the samples, the steps described above can be performed several times, preferably until the sample identified according to the method of the invention only comprises a limited number of or only one substance(s). Preferably said sample comprises substances of similar chemical and/or physical properties, and most preferably
- 20
- 25 said substances are identical. Preferably, the compound identified according to the above described method or its derivative is further formulated in a form suitable for the application in plant breeding or plant cell and tissue culture.
- The compounds which can be tested and identified according to a method of the invention may be expression libraries, e.g., cDNA expression libraries, peptides, proteins, nucleic acids, antibodies, small organic compounds, hormones, peptidomimetics, PNAs or the like (Milner, Nature Medicine 1 (1995), 879-880; Hupp, Cell 83 (1995), 237-245; Gibbs, Cell 79 (1994), 193-198 and references cited supra). Said compounds can also be functional derivatives or analogues of known inhibitors or activators. Methods for the preparation of chemical derivatives and analogues are well known to those skilled in the art and are described in, for example, Beilstein, Handbook of Organic Chemistry, Springer edition New York Inc., 175 Fifth Avenue, New York, N.Y. 10010 U.S.A. and Organic Synthesis, Wiley, New York, USA. Furthermore, said derivatives and analogues can be tested for their effects according to methods known in the art. Furthermore, peptidomimetics and/or computer aided design of appropriate derivatives and
- 30
- 35
- 40 analogues can be used, for example, according to the methods described above. The

cell or tissue that may be employed in the method of the invention preferably is a host cell, plant cell or plant tissue of the invention described in the embodiments hereinbefore.

5 Determining whether a compound is capable of suppressing or activating said resistance can be done, as described in the examples, in particular via sporulation index determination. The activator identified by the above-described method may prove useful as a fungicide or crop protectants. Thus, in a further embodiment the invention relates to a compound obtained or identified according to the method of the invention said compound being an agonist of Rpi-blb2.

10

Accordingly, in one embodiment, the present invention further relates to a compound identified by the method of the present invention.

Said compound is, for example, a homologous of Rpi-blb2. Homologues of the polypeptid of the present invention can be generated by mutagenesis, e.g., discrete point mutation or truncation of Rpi-blb2. As used herein, the term "homologue" refers to a variant form of the protein which acts as an agonist of the activity of the Rpi-blb2. An agonist of said protein can retain substantially the same, or a subset, of the biological activities of Rpi-blb2

15

20 In one embodiment, the invention relates to an antibody specifically recognizing the compound of the present invention.

The invention also relates to a diagnostic composition comprising at least one of the aforementioned polynucleotides, nucleic acid molecules, vectors, proteins, antibodies or compounds of the invention and optionally suitable means for detection.

25

The diagnostic composition of the present invention is suitable for the isolation of mRNA from a cell and contacting the mRNA so obtained with a probe comprising a nucleic acid probe as described above under hybridizing conditions, detecting the presence of mRNA hybridized to the probe, and thereby detecting the expression of the protein in the cell. Further methods of detecting the presence of a protein according to the present invention comprises immunotechniques well known in the art, for example enzyme linked immunosorbent assay. Furthermore, it is possible to use the nucleic acid molecules according to the invention in particular the markers described in the examples, e.g. in table 3a or 3b as molecular markers or primer in plant breeding.

30

35 Suitable means for detection are well known to a person skilled in the art, e.g. buffers and solutions for hybridization assays, e.g. the aforementioned solutions and buffers, further and means for Southern-, Western-, Northern- etc. -blots, as e.g. described in Sambrook et al. are known.

In another embodiment, the present invention relates to a kit comprising the polynucleotide, the vector, the host cell, the polypeptide, the antisense nucleic acid, the antibody, plant cell, the plant or plant tissue, the harvestable part, the propagation material or the compound of the invention.

- 5 The compounds of the kit of the present invention may be packaged in containers such as vials, optionally with/in buffers and/or solution. If appropriate, one or more of said components may be packaged in one and the same container. Additionally or alternatively, one or more of said components may be adsorbed to a solid support as, e.g. a nitrocellulose filter, a glass plate, a chip, or a nylon membrane or to the well of a micro
- 10 titerplate. The kit can be used for any of the herein described methods and embodiments, e.g. for the production of the host cells, transgenic plants, pharmaceutical compositions, detection of homologous sequences, identification of antagonists or agonists, etc.

- Further, the kit can comprise instructions for the use of the kit for any of said embodi-
- 15 ments, in particular for its use for increasing the resistance to one or more of said pathogens of a plant cell, plant tissue or plant.

- In a preferred embodiment said kit comprises further a polynucleotide encoding one or more of the aforementioned resistance protein, preferably Rpi-blb, and/or an antibody, a vector, a host cell, an antisense nucleic acid, a plant cell or plant tissue or a plant
- 20 related to said resistance protein(s), preferably to Rpi-blb.

- In a further embodiment, the present invention relates a method for the production of a crop protectant providing the polynucleotide, the vector or the polypeptide of the invention or comprising the steps of the method of the invention; and formulating
- 25 the polynucleotide, the vector or the polypeptide of the invention or the compound identified in step (c) of said method in a form applicable as plant agricultural composition.

- In another embodiment, the present invention relates to a method for the production of a crop protectant composition comprising the steps of the method of the present inven-
- 30 tion; and

- (a) formulating the compound identified in step (c) in a form acceptable as agricultural composition.
- 35 Under "acceptable as agricultural composition" is understood, that such a composition is in agreement with the laws regulating the content of fungicides, plant nutrients, herbicides, etc. Preferably such a composition is without any harm for the protected plants and the animals (humans included) fed therewith.

The present invention also pertains to several embodiments relating to further uses and methods. The polynucleotide, polypeptide, protein homologues, fusion proteins, primers, vectors, host cells, described herein can be used in one or more of the following methods: identification of plants resistant to plant pathogens as mentioned and related organisms; mapping of genomes; identification and localization of sequences of interest; evolutionary studies; determination of regions required for function; modulation of an activity.

Accordingly, the polynucleotides of the present invention have a variety of uses. First, they may be used to identify an organism as being *S. bulbocastanum* or a close relative thereof. Also, they may be used to identify the presence of *S. bulbocastanum* or a relative thereof in a mixed population of microorganisms. By probing the extracted genomic DNA of a culture of a unique or mixed population of plants under stringent conditions with a probe spanning a region of the gene of the present invention which is unique to this *S. bulbocastanum*, one can ascertain whether the present invention has been used or whether *S. bulbocastanum* or a close relative is present.

Further, the polynucleotide of the invention may be sufficiently homologous to the sequences of related species such that these nucleic acid molecules may serve as markers for the construction of a genomic map in related organism.

The polynucleotides of the invention are also useful for evolutionary and protein structural studies. By comparing the sequences of the Rpi-blb2 of the present invention to those encoding similar enzymes from other organisms, the evolutionary relatedness of the organisms can be assessed. Similarly, such a comparison permits an assessment of which regions of the sequence are conserved and which are not, which may aid in determining those regions of the protein which are essential for the functioning of the enzyme. This type of determination is of value for protein engineering studies and may give an indication of what the protein can tolerate in terms of mutagenesis without losing function.

These and other embodiments are disclosed and encompassed by the description and examples of the present invention. Further literature concerning any one of the methods, uses and compounds to be employed in accordance with the present invention may be retrieved from public libraries, using for example electronic devices. For example the public database "Medline" may be utilized which is available on the Internet, for example under <http://www.ncbi.nlm.nih.gov/PubMed/medline.html>. Further databases and addresses, such as <http://www.ncbi.nlm.nih.gov/>, <http://www.infobiogen.fr/>, <http://www.fmi.ch/biology/research-tools.html>, <http://www.tigr.org/>, are known to the person skilled in the art and can also be obtained using, e.g., <http://www.lycos.com>. An overview of patent information in biotechnology and a survey of relevant sources of

Agrico B.V.

20030596

64

patent information useful for retrospective searching and for current awareness is given in Berks, TIBTECH 12 (1994), 352-364.

Tables:

5

Table 1: Sequences:

Table 2. Segregation of resistance in 2851 progeny clones of BC4 mapping populations ARG 95-3 and ARP 96-11 in the field trial of 2000 at Marknesse, The Netherlands. Numbers of clones classified as having a resistant, susceptible or unknown phenotype is presented with percentages in parenthesis.

10

Mapping population	No clones with suscep- tible phenotype	No clones with resistant phenotype	No clones with un- known phenotype	Totals
ARG 95-3	846 (37)	886 (39)	551 (24)	2283
ARP 96-11	256 (45)	170 (30)	142 (25)	568
Totals	1102 (39)	1056 (37)	693 (24)	2851

Table 3A. Overview of markers used for mapping *Hpiblb2*

Marker	Or ¹⁾	Sequence	Annealing temp (°C)	Restriction Enzyme ²⁾
E46M52	F	TTGTGGTTATCGATGAGAAT	56,5	SCAR (b)
	R	GAAACAACAGCAGGATAGTGAG		
E46M52e	F	TTGTGGTTATCGATGAGAAT	61	SCAR (a,b);Mbol (c)
	R	GAAACAACAGCAGGATAGTGAG		
E40M58	F	GAATTCAGCACAAATACCAA	50	Ddel (a)
	R	TTAACGTTTACTATCACGAG		
E40M58e	F	GTAGAAACAGCAGCCTCATAAGC	55	SCAR (a)
	R	TTGTGCCTAATTGCCCTGTG		
S1E00	F	GGGGTTGGGAAGACAACGACAC	50	AFLP
	R	AATTCCAAGATACAGTCAAATAC		
41L	F	AGGCAGGATTAACAGTAGAAG	58	TaqI (a)
	R	CATGCTTTTAGGAAGAAGCTC		
38L	F	TTGAGACAAAGCAGCTCCAC	59	ApoI (a,b)
	R	ACGTTTCTCACACCTACAGG		
69L	F	TGATGGCACGTTTGATCGTG	61	TaqI (a,b);HpaII (c)
	R	TAAGATCCAAACCAGCCACC		
69R	E	CCTTATCACACATGTGGCTAC	58	RsaI (a,b); ApoI (c)
	R	ATTGAAACGGAGGAAGTACAAC		
141R	F	TTCTTCATATGGCAGACCAAC	60	RsaI (a,b); Ddel (c)
	R	CTACTCTGCTGACATGCAGG		

Agrico B.V.

20030596

PF 54801

65

24L	F	GAGATTCTCAAAGGTGTCTTCC	60	SCAR (a,b,c)
	R	AACCTGTGCTTTCCCATTCG		
24R	F	CTTTCACAAGCGTCACTTTGG	58	SCAR (a,b)
	R	TAAAAAGAATCAACAGGGCAAC		
14L	F	ACGACTGCTCAAAGTTGGCC	58	SCAR (a,b,c)
	R	CCAAGAAGCCAGTTGAGAGC		
123L	F	GTAGATTACACTATGGATATGG	60	SCAR (a,b)
	R	CAGTTAGCAGCAATGTCAGC		
123L2	F	CATTCAACTAGGCCAAAAGTGG	59	SCAR (a,b); DraI (c)
	R	CCAGGTAGGTGTTTTCTTCC		
123R	F	GTTCTAAGTCAGATGCCACC	62	SCAR (a,b)
	R	AAGTGCTCCAACACGAGCC		
133R	F	TGAGTTCTCTTACCCTGCG	60	SCAR (a,b)
	R	GGATATCCAGCATCAATGCC		
133R2	F	GGTGAGCCTCCTTGCAATCC	60	SCAR (a,b)
	R	CCTGAGGGAAGATGTCACG		
99L	F	CCTAGTTTAGAGTGAGTAGAC	58	SCAR (a,b)
	R	GTGATATATTGCTCAAGGATCC		
113R	F	GTTGCTGGCTGTCACTGATC	59	SCAR (a,b)
	R	GTGATGTGCAGGGTTCAAGG		
67L	F	GATTAGTGTAGATCTTAGCTTG	62	MboI (a,b)
	R	AAATCTCTCTCACAATTATCCC		
112L	F	CTATTGACTGAACCTGCTGAG	56	HaeIII (a); HinfI (c)
	R	TGAAGTCATTTAGTCCACAGC		
CT216 (RFLP)	F	AGATCGGAGTGTGAACATGG	56	
	R	CTTCTACTTCTAGTCGACTGC		
CT216	F	CGTAGTCCATCTGAAGCTCC	65	SCAR (a,b)
	R	TCTTCTTCTGCTAGTCGTCC		
CT119	F	ACTATTCTCACGTAAGGGGACAC	60	HindIII (a,b)
	R	GTGTACATGTATGAAACTCTAGC		
CT119N	F	GTTCCCTTCAATCAGAAAGTAG (aro 120)	55	SCAR (a)
	R	CTTTGGATGAGTCAAAAGGCT (aro 121)		
14L24L	F	univ14L	60	CfoI (c)
	R	univ24L		
SPB30L	F	CAAGTTACGGCAACCAAGAG	57	HpaII (c)
	R	CTTTGACACAGTGTTAGAATGC		
SPB39L	F	CGTGATCTAGGAGTTACGAC	52	SCAR (c)
	R	CTTATTTTAAATACAAGACATCTGG		
24L9spec	F	univ. 14L	56	HhaI (c)
	R	CAGAGGAAAGTCAACCAACG		
24Lspec	F	univ. 14L	60	CfoI (c)
	R	CAGAGGAAAGTCAACCAACG		
- NptII	F	TCGGCTATGACTGGGCACAACAGA	70	
	R	AAGAAGGCGATAGAAGGCGATGCG		

Agrico B.V.

20030596

66

M13

F TGTAACGACGGCCAGT
R GGAAACAGCTATGACCATG

55

¹⁾ Ori: Orientation of the primer; F: forward primer; R: reverse primers
²⁾ a: ARG95-3, b: ARP86-11, c: B6a

Tabel 3B. Overview of primers used for mapping *Rpl-bib2*

primer	Ori	Sequence ¹⁾
ARO 73	F	TTCAGCACAAATACCAAT
ARO 74	R	GATGTTCCCTTCTTTTA
ARO 77	R	TTGTGGTTATCGATGAGAAT
ARO 79	R	ACCTGGCGTTCCTTATTTTT
ARO 94	F	NGTCASWGANAWGAA
ARO 128	F	GATGGAGCGGAAAAGCCGGTG
ARO 129	F	GGTGTGTTGTAGCATCTCCAG
ARO 295		CCATGATTACGCCAAGCTGG
ARO 296		GGTTTTCCCGAGTCACGACGT
univ14L	F	AGAAAGCTCACCAGTGGACC
univ24L	R	ATTTATGGCTGCAGAGGACC
123Mi	R	AAGTCCAATTGCTCATCCATC
14L2	R	TGCACCATGCACGAAGGTC
24L2	F	CAATWTTGGTTCCCGAAATTGG
ARF1F	F	ATGGAAAAACGAAAAGATAATGAAG
ARF1R	R	CTACTTAAATAACGGGATATCCTTC
ARO 602	F	CCCATGACTCCTTGAGTTTG
S1		GGTGGGGTTGGGAAGACAACG
EcoR1+0		GTAGACTGCGTACCAATTC
MseI+0		GATGAGTCCTGAGTAA

¹⁾ N=A+T+G+C, S=G+C, W=A+T

Agrico B.V.

20030596

PF 54801

67

Table 4. Complementation of late blight susceptibility in potato

BAC-library	Source BAC	Genotype ¹	cv Impala		cv Kondor	
			RGC-containing plants/ transformants	R plants/ RGC-containing plants	RGC-containing plants/ transformants	R plants/ RGC-containing plants
ARD 1197-16	24	R ₀ (RGC1)	12/15 ^a	0/12		
			8/10 ^b	0/8		
	24	R ₀ (RGC2)	8/11 ^a	0/8		
			5/6 ^b	0/5		
	24	R ₀ (RGC3)	11/13 ^a	0/11		
Elb 2002	211	R ₀ (RGC4)	5/7 ^b	0/5	10/12 ^a	0/10
	242	R ₀ (RGC4)	5/7 ^a	0/5	8/8 ^a	0/8
	211	R ₀ (RGC5)	5/7 ^a	4/5	12/13 ^a	12/12
	211	R ₀ (RGC6)	-	-	-	-
	211	R ₀ (RGC24L)	-	-	-	-
	SPB39	R ₀ (RGC4)	5/6 ^a	0/5	3/3 ^a	0/3
	SPB39	R ₀ (RGC5)	11/15 ^a	11/11	8/8 ^a	7/8
	SPB39	R ₀ (RGC6)	3/3 ^a	0/3	0/6 ^a	0/6
	SPB30	R ₀ (RGC7)	3/4 ^a	0/3	8/9 ^a	0/9
	SPB30	R ₀ (RGC8)	-	-	-	-
	SPB39	R ₀ (24L)	-	-	-	-
		R ₀ (pBINPLUS)	3/3	0/3	8/10	0/8

¹ R₀ genotypes are primary transformants obtained from transformation of the susceptible potato cultivars Impala or Kondor with T-DNA constructs containing the *Rpi-bib2* gene candidates RGC1 to RGC8 and RGC24L or an empty pBINPLUS vector. *Agrobacterium tumefaciens* strains UA143^a or AGLO^b were used for transformation of the *P. infestans* susceptible potato cultivars Impala and Kondor.

Table 5. Cycling conditions used for TAIL-PCR

Reaction	cycle no.	Thermal condition
Primary	1	92°C (2 min), 95°C (1 min)
	5	94°C (15s), 63°C (1 min), 72°C (2 min)
	1	94°C (15s), 30°C (3 min), ramping to 72°C over 3 min, 72°C (2 min)
	10	94°C (5s), 44°C (1 min), 72°C (2 min)
	12 ^a	94°C (5s), 63°C (1 min), 72°C (2 min)
		94°C (5s), 63°C (1 min), 72°C (2 min)
		94°C (5s), 44°C (1 min), 72°C (2 min)
	1	72°C (5 min)
Secondary	10 ^a	94°C (5s), 63°C (1 min), 72°C (2 min)
		94°C (5s), 63°C (1 min), 72°C (2 min)
		94°C (5s), 44°C (1 min), 72°C (2 min)
	1	72°C (5 min)
Tertiary	20	94°C (10s), 44°C (1 min), 72°C (2 min)
	1	72°C (5 min)

^a these are nine-segment super cycles each consisting of two high-stringency and one reduced-stringency cycle

5

The figures show:

- 10 **Figure 1.** Schematic representation of the development of the complex interspecific hybrid clones designated as 'ABPT' (1a) and the *S. tuberosum* mapping populations that were derived from two of these clones: ABPT clone 55 and ABPT clone 60 (1b to d). A; *Solanum acaule*, B; *S. bulbocastanum*, P; *S. pureja*, T; *S. tuberosum*, 2x : diploid ($2n=2x=24$), 3x; triploid, 4x; tetraploid, 6x; hexaploid, cv; cultivar. Codes in
15 italics indicate mapping populations.

- Figure 2.** Disease progress curves for clone ARF 87-601 and susceptible control cultivars (cv) Bildtstar, Eersteling and the partial resistant control cultivar Pimpernel in a field test for foliar resistance to late blight in Toluca Valley, Mexico in 1991. At eight
20 time points after planting, the percentage blighted foliage due to a natural late blight infection was scored on the 1 to 9 CIP scale (Estrada-Ramos, 1983).

Figure 3. Disease progress curves for clone ARF 87-507, ARF 87-601, ARF 87-801, the susceptible control cultivar (cv) Granola and the partial resistant breeding clone AR 85-96-13 in a field test for foliar resistance to late blight in Benguet Province, Philippines in 1992. At six time points between August 25th to November 24th, the percentage blighted foliage due to a natural late blight infection was scored on the 1 to 9 CIP scale (Estrada-Ramos, 1983).

Figure 4. Typical phenotypes in tetraploid resistant and susceptible parental clones and progeny clones segregating for *Rpi-blb2* mediated resistance to late blight in the annual field trial at Marknesse, The Netherlands, approximately 6 weeks after inoculation with isolate IPO82001 of *P. infestans*. Six plant plots with a clone showing the resistant phenotype (within black solid line) that shows no or hardly any sporulating lesions and with a clone showing the susceptible phenotype (within white dotted line) that shows completely blighted foliage.

Figure 5. Genetic map based on 109 progeny clones of *S. tuberosum* mapping population ARG 95-15 showing 7 AFLP markers that were found to cosegregate with the *Rpi-blb2* locus. Numbers left to the vertical line indicate the genetic distance between flanking markers or the *Rpi-blb2* locus in centimorgan (cM).

Figure 6. Genetic map based on 137 progeny clones of *S. tuberosum* mapping population ARG 95-3 showing 15 AFLP markers and RGA marker S1E00 that were found to cosegregate with the *Rpi-blb2* locus. Phenotypes of the progeny clones were obtained with detached leaf assays. Numbers left to the vertical line indicate the genetic distance between flanking markers or the *Rpi-blb2* locus in centimorgan (cM).

Figure 7. Genetic map based on 178 progeny clones of *S. tuberosum* mapping population ARG 95-3 showing 5 markers that were found to cosegregate with the *Rpi-blb2* locus on linkage group 6 of *S. tuberosum*. Phenotypes of the progeny clones were determined in the field trial at Marknesse, the Netherlands in 1998. Markers E40M58 and E46M52 were scored either as AFLP, CAPS, SCAR or extended (suffix: e) marker (table 3A). Partly, marker CT119 was scored as marker CT119N (table 3a). Marker CT216 was scored as SCAR marker. The number left to the vertical line indicates the genetic distance between flanking markers or the *Rpi-blb2* locus in centimorgan (cM). For each marker, the number of recombinants between marker and phenotype and the total number of progeny clones scored is given in parenthesis.

Figure 8. Genetic maps based on 886 progeny clones of *S. tuberosum* mapping population ARG 95-3 and on 170 progeny clones of *S. tuberosum* mapping population ARP 96-11, showing markers that were found to cosegregate with the *Rpi-blb2* locus

on linkage group 6 of *S. tuberosum*. Phenotypes of the progeny clones were determined in the field trial at Marknesse, the Netherlands in 2000. The number left to the vertical line indicates the genetic distance between flanking markers in centimorgan (cM). The marker interval which delimitates the position of the *Rpi-blb2* gene, based on detected recombination events in progeny clones, is indicated by double arrow headed lines.

Figure 9. Physical map of the genomic region containing *Rpi-blb2* in *S. tuberosum* (upper horizontal line) and *S. bulbocastanum* (lower horizontal line). Vertical lines indicate the relative position of markers linked to resistance. Numbers above the horizontal lines are the number of recombinants identified between the flanking markers in 1056 and 1899 progeny plants of *S. tuberosum*, derived from complex species hybrids "ABPT" (Figure 1), and *S. bulbocastanum* progeny plants respectively. ABPT-derived progeny comprises clones from both the mapping populations ARG 95-3 and ARP 96-11. Rectangles represent bacterial artificial chromosome (BAC) clones from the ARD 1197-16 BAC library except for BAC clones with prefix "Bib" which were from the *S. bulbocastanum* Bib 2002 BAC library. The marker interval which delimitates the position of the *Rpi-blb2* gene, based on detected recombination events in progeny clones, is indicated by double arrow headed lines. Small arrows indicate positions of Resistance Gene Candidates (RGC's).

Figure 10. Schematic representation of the development of the diploid, intraspecific mapping population B6 of *S. bulbocastanum*. Codes in italics indicate mapping populations.

Figure 11. Genetic map based on 1899 progeny clones of *S. bulbocastanum* mapping population B6, showing markers that were found to cosegregate with the *Rpi-blb2* locus on chromosome 6 of *S. bulbocastanum*. Phenotypes of the progeny clones were determined by detached leaf assays. The number left to the vertical line indicates the genetic distance between flanking markers in centimorgan (cM). The marker interval which delimitates the position of the *Rpi-blb2* gene, based on detected recombination events in progeny clones, is indicated by a double arrow headed line.

Figure 12. Genetic complementation for late blight susceptibility. Typical disease phenotypes of potato (*S. tuberosum*) leaves, 6 days after inoculation with a sporangiospore suspensions of *P. infestans* isolate 655-2A. Leaf derived from kanamycin resistant cv Kondor plants transformed with pBINPLUS (control; A), leaves derived from cv Kondor plants harbouring BAC SPB39 derived (B) or BAC 211 derived RGC5 (C), leaf derived from kanamycin resistant cv Impala plants transformed with pBINPLUS (control; D), leaves derived from cv Impala plants harbouring BAC SPB39 derived (E) or BAC 211

derived *RGC5* (F). Panels A and D depict typical susceptible responses with extensive sporulating lesions of *P. infestans*. Panels B, C, E and F depict typical resistance reactions observed at the sites of inoculation on transgenic potato plants harbouring *Rpi-blb2*.

5

Figure 13. Nucleic acid sequences coding for the *Rpi-blb2* gene. **A.** Coding nucleic acid sequence of the *Rpi-blb2* gene. **B.** Coding nucleic acid sequence of the *Rpi-blb2* gene including the intron sequence (position 43-128). **C.** Sequence of the 7967 bp *Sau3AI* genomic DNA fragment of ARD 1197-16 BAC 211 present in p211F-C12, one of the two the genetic constructs used for genetic complementation for late blight resistance. The genomic fragment harbours the *Rpi-blb2* gene including natural regulatory elements necessary for correct expression of the gene. The initiation codon (ATG position 1546-1548) and the termination codon (TAG position 5433-5435) are underlined. **D.** Sequence of the 9949 bp *Sau3AI* genomic DNA fragment of *S. bulbocastanum* 2002 BAC BlbSP39 present in pSP39-20, one of the two the genetic constructs used for genetic complementation for late blight resistance. The genomic fragment harbours the *Rpi-blb2* gene including natural regulatory elements necessary for correct expression of the gene. The initiation codon (ATG position 1413-1415) and the termination codon (TAG position 5300-5303) are underlined.

20

Figure 14. Putative *Rpi-blb2* gene structure and deduced *Rpi-blb2* protein sequence. **A.** Schematic representation of the *Rpi-blb2* gene structure. Horizontal lines indicate exons. Open boxes represent coding sequence. Lines angled downwards indicate the positions of intron sequences. **B.** Deduced *Rpi-blb2* protein sequence. The amino acid sequence deduced from the DNA sequence of *Rpi-blb2* is divided into three domains (LZ, NBS and LRR). Hydrophobic residues in domain A that form the first residue of heptad repeats of the potential leucine zipper (LZ) domain are underlined. Conserved motifs in R proteins are written in lowercase and in italic in the NBS domain. Residues matching the consensus of the cytoplasmic LRR are indicated in bold in the LRR domain. Dots in the sequence have been introduced to align the sequence to the consensus LRR sequence of cytoplasmic LRRs.

30

Figure 15. Alignment of the deduced protein products encoded by *Rpi-blb2*, *Mi-1.1* and *Mi-1.2*. The complete amino acid sequence of *Rpi-blb2* is shown and amino acid residues from *Mi-1.1* and *Mi-1.2* that differ from the corresponding residue in *Rpi-blb2*. Dashes indicate gaps inserted to maintain optimal alignment. Amino acid residues that are specific for *Rpi-blb2*, when compared to those at corresponding positions in *Mi-1.1* and *Mi-1.2* are highlighted in bold and red. The regions of the LRRs that correspond to the β -strand/ β -turn motif *xxLxLxxx* are underlined. Conserved motifs in the NBS domain are indicated in lowercase. A vertical line indicates the division between CC-NBS

40

and LRR region. The position of the VLDL motif which is conserved in the third LRR of many plant R proteins but not in Rpi-b1b2 is indicated by a shaded rectangle.

- 5 This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patent applications, patents, and published patent applications cited throughout this application are hereby incorporated by reference.

Examples

10

Example 1: Evaluation of resistance in ABPT derived back cross clones and populations

- 15 BC2-clones ARF 87-507 and ARF 87-801 were selected from BC2-progeny obtained after two rounds of backcrossing on complex species hybrid ABPT-clone number 55 (Figure 1a) with late blight (LB) susceptible *S. tuberosum* cultivar Oberambacher Frühe as first parent and *S. tuberosum* cultivars Arkula (Figure 1b) and Blanka (Figure 1c) respectively as second parents. Similarly, BC2-clone ARF 87-601 was obtained by successive crossing on ABPT-clone 60 with LB susceptible *S. tuberosum* cultivars
- 20 Alcmaria and Blanka (Figure 1d).

- Clone ARF 87-601 was tested as part of a field test for screening of LB-resistance in the Toluca area in Mexico in 1991. A plot of clone ARF 87-601 with seven plants was evaluated in comparison to plots with nine plants each of the control cultivars Bildtstar,
- 25 Eersteling and Pimpernel. According to the ratings for resistance to late blight in the Dutch National list of recommended potato cultivars of 1988, these control cultivars scored 3, 3 and 8 respectively on a scale from 3 to 8 of increasing resistance. Cultivar Pimpernel is considered as a source of partial resistance (Colon *et al.*, 1985). About forty days after planting, natural infection by *P. infestans* established. The development
- 30 of LB in the foliage then was monitored eight times during the period from July 16th to September 2nd (Figure 2). There was a clear difference between the disease progress curves for ARF 87-601 in comparison to the control cultivars. At 74 days after planting, foliage of the control cultivars was completely or nearly completely blighted whereas clone ARF 87-601 showed no visible symptoms (Figure 2). Clones ARF 87-507,
- 35 ARF 87-801 and again clone ARF 87-601 showed comparable results in a field test for screening of LB-resistance in the Benguet Province of the Philippines in 1992 (Figure 3). Ten plants each of the three BC2 clones, control cultivar Granola and the moderately LB resistant breeding clone AR 85-96-13, which was used as female parent to obtain AR 92-1197 (Figure 1d), were planted on August 25th. The percentage of blighted foliage was scored six times after occurrence of natural infection by *P. infestans*.
- 40

P. infestans. Disease progress curves of ABPT derived BC2-clones were markedly different when compared to cultivar Granola and clone AR 85-96-13 (Figure 3). BC2-clones showed no or little LB symptoms and no clear disease progress during the scoring period whereas cultivar Granola had almost completely blighted foliage at the third scoring date.

Clones ARF 87-601, ARF 87-507 and ARF 87-801 were used for further backcrossing with LB susceptible cultivars and breeding clones of *S. tuberosum* (Figure 1b to 1d). This breeding work resulted in four different mapping populations, tetraploid BC3-
10 population ARG 95-15, tetraploid BC4-populations ARG 95-3 and ARP 96-11 and diploid BC4-population DP1. During the successive steps of this breeding work resistant clones ARF 87-507, ARF 87-601, ARF 87-801, AR 91-1263, AR 91-1292 and AR 92-1197 were selected on the basis of agronomic performance in common practice breeding evaluations as well as by screening their parents and relevant progenies in a field
15 trial at Marknesse, the Netherlands, that was inoculated with the complex isolate IPO82001 of *P. infestans*. The diploid ($2n=2x=24$) clone ARD 1197-16 was selected among the progeny of cross AR 92-1197 x Phu 81-101 (Figure 1d), the latter parental clone being known for its capacity to induce parthenogenic seed set in the female parent (Hermesen and Verdenius, 1973). Initially, resistance to LB in ARD 1197-16 was
20 found after repeated detached leaf assays using *P. infestans* isolates IPO82001, IPO655-2A and IPO428-2 and verified in a field trial in 1998 at Marknesse. The diploid status of clone ARD 1197-16 was confirmed by flow cytometry (Plant cytometry services, Schijndel, the Netherlands).

25 Clear segregation for the LB-resistance trait in ABPT-derived progeny and mapping populations was observed during successive years of field testing at the trial site of Marknesse, approximately 6 weeks after inoculation with isolate IPO82001 of *P. infestans*. Typically, resistant clones showed no or hardly any sporulating lesions whereas susceptible clones showed completely blighted foliage (Figure 4) In 2000, a
30 total of 2851 clones from the mapping populations ARG 95-3 and ARP 96-11 were screened as single plant plots. On average, 24 percent of the clones showed phenotypes that could not unambiguously be classified as resistant or susceptible. Clones that could be classified as such showed segregation ratio's of resistant to susceptible phenotypes of 1 to 1 and 1 to 1.5 for populations ARG 95-3 and ARP 96-11, respectively
35 (Table 2).

Detached leaf assay's with ABPT-derived progeny and mapping populations where found to be less accurate for phenotyping than screening under field conditions. Nevertheless, results of detached leaf assays were considered suitable for the initial determi-

nation of the phenotype of individual clones and thus, for construction of mapping populations.

Example 2: Genetic mapping of the *Rpi-blb2* resistance locus in ABPT-derived back cross populations.

In all four mapping populations (Figure 1), resistance segregated as expected for a monogenic trait, suggesting the presence of a dominant resistance allele at a single locus (Table 2). This locus was designated the *Rpi-blb2* locus.

In order to identify markers linked to *Rpi-blb2*, an initial AFLP analysis with 14 primer combinations (pc) was carried out on DNA of 10 resistant and 10 susceptible ARG 95-15 progeny plants, based on detached leaf assay, including the parental clones. The testing of 21 potentially linked markers on an additional 89 plants identified several markers linked to resistance (Figure 5). Subsequent bulked segregant analysis (BSA) with 160 pc's on 2 resistant and 2 susceptible DNA pools, each containing genomic DNA of 8 resistant or susceptible ARG 95-15 progeny plants, respectively, identified a total of 58 AFLP markers potentially linked to resistance (Figure 5). When a number of these markers were tested on 137 progeny plants of ARG 95-3, they were also linked to resistance in this population, suggesting that the resistance in the two populations was determined by the same locus (Figure 6). These cosegregating markers mapped 3 to 28 centimorgan (cM) and 1 to 7.2 cM to one side of the locus in ARG 95-15 and ARG 95-3 respectively, suggesting that *Rpi-blb2* could be situated at a distal position on a chromosome.

To determine the position of the *Rpi-blb2* on the genetic map of potato, the two cosegregating AFLP markers E40M58 and E46M52 (Figure 6) were cloned into the pGEM-T vector (Promega, the Netherlands) and sequenced. Primers designed on the ends of the sequences of the cloned AFLP fragments (Table 3) were used to develop cleaved amplified polymorphic sequence (CAPS) marker E40M58 that was found to be cosegregating with the resistance trait in 25 resistant and 25 susceptible clones of ARG 95-3. CAPS marker E40M58 was subsequently tested on 46 progeny plants of the CxE mapping population (van Eck *et al.*, 1995). These data were added to the existing marker scores of the CxE population. Joinmap (Stam, 1993) linkage analyses mapped E40M58 8 cM distal to GP79 (Gebhardt *et al.*, 1991), positioning *Rpi-blb2* on the short arm of chromosome 6. In 178 progeny plants of population ARG 95-3 no recombination between *Rpi-blb2* and AFLP markers E40M58, E40M60 and CAPS marker CT119 was observed. AFLP marker E46M52 and sequence characterised amplified region (SCAR) marker CT216 mapped 2.2 cM proximal to the gene (Figure 7).

Example 3: Identification of a RGA marker linked to *Rpi-blb2*

In an attempt to identify functionally relevant markers linked to resistance, primers designed on the conserved motifs of the NBS domain of plant *R* genes (Leister *et al.*, 1996), were used in an adapted AFLP protocol (RGA-AFLP) to identify resistance gene analogue (RGA) specific markers.

Using the P-loop based primer S1 from Leister *et al.* (1996) in combination with the Eco00 AFLP primer, an RGA specific marker, S1E00 was developed which cosegregated with resistance and markers E40M58 and CT119 in the ARG 95-3 mapping population (Figure 6 and 7).

Example 4: Development of E40M58e and E46M52e SCAR markers for recombinant screening.

Using genomic DNA of AR 91-1263 as template, the cloned fragment of AFLP marker E46M52 was extended by TAIL-PCR. The primary TAIL-PCR was performed using primers ARO 77 (sp1) and ARO 94 (AD). Subsequently, the secondary PCR was performed using ARO 128 (sp2) and the tertiary PCR using ARO 129 (sp3) both in combination with primer AD. This resulted in an E46M52e fragment that was extended on the 5' end with approximately 500 bp. The E46M52e fragment was cloned in pGEM-T and sequenced. A new forward primer was designed on this sequence and PCR in combination with primer ARO 77 resulted in SCAR marker E46M52e that cosegregated with the resistant phenotype in the four *S. tuberosum* mapping populations and as CAPS marker also in population B6.

Using genomic DNA of ARD 1197-16 as template, the cloned fragment of AFLP marker E40M58 was also extended by TAIL-PCR. The primary TAIL-PCR was performed in both the 5' and 3' directions using sp1 primers ARO 73 (3') or 74 (5') in combination with primer AD. Subsequently, the secondary PCR was performed using as sp2 ARO 82 or 79, respectively. The fragments obtained from the secondary PCR, 750 bp from the 3' end and 400 bp from the 5' end were cloned in pGEM-T and sequenced. On the basis of both sequences, two new primers were designed resulting in a SCAR marker that cosegregated with resistance in mapping population ARG 95-3 and DP1 (Table 3). The fragment of SCAR marker E40M58e could be amplified in the resistant parents of mapping populations ARG 95-3 and DP1, which were both derived from ABPT clone 55 (Figure 1), but PCR amplification in the parents or progeny clones of mapping populations ARP 96-11 and ARG 95-15, which were both derived from ABPT clone 60, did not give any detectable PCR product. It was assumed that this could have been caused by minor differences in the genomic sequence and therefore, the AFLP

76

fragment was extended by TAIL-PCR using genomic DNA of clone AR 91-1292 as template. A fragment E40M58e2 of approximately 300 bp was obtained; cloned and sequenced. Comparison of the sequence with the original fragment of AFLP marker E40M58 showed that only the first 37 bp of the extended fragment were identical. PCR with primers designed on the sequence of E40M58e2 did not result in a polymorphic marker. Both of the extended markers E40M48e and E40M58e2 were tested on five resistant or susceptible clones of *S. bulbocastanum* (BGRC 8005 and 8006). Only the fragment of SCAR marker E40M58e could be amplified in four *S. bulbocastanum* clones, indicating that part of the sequence of E40M58e2 was not derived from *S. bulbocastanum*. This observation suggested that E40M58e was located on the border of the *S. bulbocastanum* introgression fragment in clone AR 91-1292 and that the position of the *Rpi-blb2* locus was proximal to marker E40M58e.

Example 5: Mapping of *Rpi-blb2* in a diploid mapping population derived from ABPT material

A total of 149 progeny clones of diploid mapping population DP1 were screened with markers E40M58e and E46M52e. No recombination was found between these markers suggesting suppressed recombination in the genomic region studied when compared to the tetraploid mapping population ARG 95-3 (Figure 7). A subset of 112 clones was screened for resistance to *P. infestans* isolates IPO82001, IPO655-2A and IPO428-2 in a partially repeated detached leaf assay. Eleven of the clones (11%) showed intermediate reactions and were classified as having unknown phenotypes. Another 51 and 50 clones were classified as resistant and susceptible respectively. Three progeny clones DP1-28, DP1-79 and DP1-81 were identified that were putatively recombined between the *Rpi-blb2* locus and the markers E40M58e and E46M52e. In 2000, a subset of 50 out of the 112 phenotyped clones was tested for resistance to LB in the field at the trial site of Marknesse. Conclusive results on the phenotype for LB resistance were obtained for 33 out of the 50 clones. The phenotype of clones 28 and 81 as determined with the detached leaf assay appeared to be erroneous. Thus, it was concluded that these clones did not represent recombination events between *Rpi-blb2* and the markers used. The phenotype of clone DP1-79 could not be verified conclusively under field conditions and this clone may represent the only recombination event between the *Rpi-blb2* locus and the markers E40M58e and E46M52e in 101 progeny clones of DP1 (1 cM). Since it was shown that two markers, linked to the resistance trait in ARG 95-15, ARG 95-3 and ARP 96-11, cosegregated with the same locus for LB-resistance in DP1, it was concluded that the DP1 parental clone ARD 1197-16 was suitable as a source for *Rpi-blb2* gene isolation in a map-based cloning approach.

Example 6: Physical mapping of the ABPT derived *Rpi-blb2* locus

The resistant clone ARD 1197-16, heterozygous for the *Rpi-blb2* locus, was used as source DNA for the construction of a BAC library (hereafter referred to as the
5 ARD 1197-16 BAC library). High molecular weight DNA preparation and BAC library construction were carried out as described in Rouppe van der Voort *et al.* (1999). Initially, a total of 67968 clones with an average insert size of 100 kb, which corresponds to approximately 7 genome equivalents, were individually stored in 177 384-well micro-
10 titer plates at -80°C. Marker screening of the ARD 1197-16 BAC library was carried out as described in Rouppe van der Voort *et al.* (1999). Essentially, DNA pools generated for each 384-well plate were screened by PCR with SCAR or CAPS markers linked to the *Rpi-blb2* locus in order to build a BAC contig across the *Rpi-blb2* locus.

Screening of the ARD 1197-16 BAC library with markers E40M58e, S1E00 and CT119
15 Identified several positive BAC clones, which served as seed BACs from which a chromosome walk across the *Rpi-blb2* locus was initiated. Marker E40M58e was used to isolate the BAC clones 69 and 141 whereas BAC clones 14, 24, 123 and 133 were positive for marker S1E00. Marker CT119 was used to isolate BAC 67. After sequencing the left (L) and right (R) borders of these BAC clones, a new set of markers was
20 developed; 14L, 24L, 24R, 69L, 69R, 141R, 123L, 123R, 133R and 67L. Screening of the isolated BAC clones with these markers showed that the following pairs of BAC clones shared overlap: the right side of 123 with the left side of 133, 14 completely with 24, and the left side of 69 with the right side of 141. BAC 67 did not share overlap with the other BAC clones. The finding that the S1E00 positive BAC clones 14, 24, 123, and
25 133 did not form a single contig indicated that S1E00 was a repetitive sequence. This, together with the finding that the right BAC-end sequences of BAC clones 24 and 123 showed high homology to different regions of the *Mi1* resistance gene from tomato (Milligan *et al.*, 1998, Simons *et al.*, 1998), suggested that the *Rpi-blb2* locus harboured more than one *RGA*. Screening of the initial ARD 1197-16 BAC library with markers
30 141R, 24L, 24R and 123L did not lead to contig extension. However, screening of the library with markers 123R and 133R resulted in the isolation of BAC clones 99 and 113, thereby extending the BAC 123/133 contig in one direction. BAC-end sequencing of these two BAC clones lead to the development of two new markers, 99L and 113R. Screening of the ARD 1197-16 BAC library with 69R lead to the extension of the
35 141/69 contig. Consecutive screening of the BAC library with markers derived from BAC clones that further extended this contig lead to the isolation of BAC clones 36, 41 and 112, and the development of markers 36L, 41L and 112L.

In an attempt to complete the BAC contig across the *Rpi-blb2* locus, the ARD 1197-16
40 BAC library was enlarged with an additional 38864 BAC clones of ~100kb (384-well

plate numbers 178-273). This second library was screened with markers 24L, 24R, 123L, and 141R, leading to the identification of BAC clones positive for both 24R and 123L (e.g. 191) and BAC clones positive for 24L (211, 242). In this way, the gap between BAC 24 and 123 was closed and the 24/14 contig was extended towards BAC clone 141. There were no new clones in the extended ARD 1197-16 library that were positive for marker 141R.

Example 7: Construction of additional markers in BAC 123/133 region.

10 In an attempt to develop additional polymorphic markers from BAC 123 and 133, a 10 kb sub-clone library was constructed of both BAC 123 and 133. BAC DNA was partially cleaved with *Sau3A*I and fragments of approximately 10 kbp were cloned in the *Bam*HI site of vector pBINPLUS. In order to select clones containing the original BAC-end sequence, 288 subclones of BAC 123 and 192 of BAC 133 were screened with the BAC-end markers 123L or 133R. In total 14 subclones were positive for marker 123L and 11 for marker 133R. Subsequently, the orientation of the BAC-end positive clones was determined by several PCRs using either the forward or reverse primer of the relevant BAC-end marker in combination with primers M13F or M13R (Table 3). For marker 123L three sub-clones and two sub-clones for marker 133R were selected and the ends not containing the 123L or 133R marker were sequenced (approximately 500 bp). Based on the new sequence two new primers were designed for subclone 123 resulting in marker 123L2 and two new primers were designed for subclone 133 resulting in marker 123R2. SCAR marker 123L2, which was located 10 kbp proximal to marker 123L, appeared to be polymorphic in mapping populations ARG 95-3, ARP 96-11 and as CAPS in B6. SCAR marker 133R2, which was located 10 kbp distal to marker 133R, was only polymorphic in mapping populations ARG 95-3 and ARP 96-11.

Example 8: Fine mapping of the *Rpi-blb2* locus in ABPT derived mapping populations.

30 In order to fine map the *Rpi-blb2* locus in ABPT derived mapping populations a total of 2283 new progeny clones of mapping population ARG 95-3 and 598 clones of mapping population ARP 96-11 were tested for resistance to LB in the field at the trial site of Marknesse in 2000 (Table 2). In population ARG 95-3 846 clones (37%) were scored susceptible and 886 clones resistant (39%). The phenotypes of the remaining 551 clones were unclear. In population ARP 96-11 256 clones (45%) were scored susceptible and 170 clones (30%) resistant. The phenotypes of the remaining 142 (25%) were unclear (Table 2). The 846 and 170 resistant clones from mapping populations ARG 95-3 and ARP 96-11, were selected for recombinant screening with SCAR marker CT216 and CAPS marker 41L or 36L, respectively. In total 85 (9.6 cM) and 22

(12.9 cM) recombinants were obtained in mapping populations ARG 95-3 and ARP 96-11 respectively, that were subsequently screened with CAPS marker 67L, reducing the number of recombinants to 5 (0.56 cM) in the marker interval 67L – 36L in case of mapping population ARG 95-3 and to 4 recombinants (2.35 cM) in the marker interval 67L – 41L in case of the mapping population ARP 96-11 (Figure 8). These remaining 9 recombinants were further analysed with SCAR and CAPS markers 113R, 99L, 133R, 133R2, 123R, 123L, 24R, 14L, 24L, 141R, 69L, E40M58e and 69R. The latter two markers were scored only in mapping population ARG 95-3.

10 In population ARG 95-3 two clones showed recombination between markers E40M58e and 69L, positioning the *Rpi-blb2* gene 0.23 cM proximal to marker E40M58e. Two other clones were recombined between markers 113R and 67L and one was recombined between markers 133R2 and 133R, positioning the *Rpi-blb2* gene 0.11 cM distal to marker 133R.

15 In population ARP 96-11, no recombination was detected between markers 41L and 69L, positioning the *Rpi-blb2* gene 0.58 cM proximal to marker 36L. Two progeny clones were recombined between markers 113R and 67L, and one clone was recombined between markers 99L and 133R, positioning the *Rpi-blb2* gene 0.58 cM distal to marker 99L (Figure 8; Figure 9).

20 **Example 9: Evaluation and genetic mapping of late blight resistance in a *S. bulbocastanum* intraspecific mapping population.**

25 In order to develop an intraspecific mapping population of *S. bulbocastanum*, a resistant clone Blb 2002 was obtained from an inter accession cross (Figure 10). This clone was reciprocally crossed with a susceptible clone Blb 48-5 that was selected also in progeny from an inter accession cross (Figure 10). The resulting population was designated B6 with synonyms B6a, Blb 99-229, Blb 00-7 and Blb 00-8.

30 Initially a small group of 47 progeny plants of the B6 population was screened for resistance to *P. infestans* in a partially repeated detached leaf assay using a sporangio-spore solution of isolate IPO655-2A of *P. infestans* as inoculum. Plants with leaves that clearly showed sporulating lesions 6 to 9 days after inoculation were considered to have a susceptible phenotype whereas plants with leaves showing no visible symptoms or necrosis at the side of inoculation in the absence of clear sporulation were considered to be resistant. Of the 47 seedlings, 23 scored resistant and 24 susceptible. These data indicated that the progeny of mapping population B6 gave clear segregation of the resistance trait in the detached leaf assay and that resistance could be due to a single dominant gene or a tightly linked gene cluster. In order to determine the

Agrico B.V.

20030596

PF 54801

80

chromosome position of this locus, 46 seedlings were analysed with markers 112L and E46M52e. Marker 112L was found to be linked in repulsion with the resistant phenotype, as only two recombinants were obtained between this marker and the phenotype of the 46 seedlings (4 cM). Also, marker E46M52e was found to be linked in repulsion with the resistant phenotype. Here, five recombinants were obtained between marker E46M52e and the phenotype (11 cM). Furthermore, markers 69R, 69L and 141R were used for analysis of the seven recombinants between markers 112L and E40M58e with an additional group of 6, 15 and 14 non recombined seedlings respectively, and found to be completely linked in either coupling (marker 69R) or repulsion phase (markers 69L and 141R) to resistance, indicating that the resistance gene was located at the same locus, i.e. *Rpi-blb2*, as in the ABPT-derived mapping populations.

In order to determine the position of *Rpi-blb2* more precisely relative to the available markers, another 849 seedlings of the B6 mapping population and 1054 seedlings from the reciprocal cross (Figure 10) were grown and analysed for recombination between the markers E46M52e and 112L. Thus, in addition to the initial 47 seedlings, a total of 1903 individual offspring clones of the B6 population were screened. Recombination between markers E46M52e and 112L was detected in a total of 138 of these seedlings (7.25 cM). Fine mapping of the *Rpi-blb2* locus was carried out in two steps. Firstly, the group of 138 recombinants was reduced to 19 by additional screening with markers 14Lb, 113R, 123L2, 24L, 141R and 69L (Table 3), derived from left (L) and right (R) border sequences of BAC clones isolated from the ARD 1197-16 BAC library and subsequent selection of all the seedlings that were recombined between markers 113R and 69L. Possibly due to double recombination, 4 recombinants gave patterns for the markers scored that deviated from scores expected in the case of single recombination events in the genetic interval studied and when assuming co-linearity of markers. These were withdrawn from further analyses. Secondly, the remaining 15 recombinants were analysed with markers from border sequences of BAC clones isolated from the Bib 2002 library, SPB39L and SPB30L, or with MiGA markers 24L9spec, 24Lspec and 14L24L (Table 3). Results of marker analyses of these remaining 15 recombinants, which gave clearly interpretable marker scores and phenotypes, positioned the *Rpi-blb2* locus between markers 69L and 24L, on a 0.11 cM (n=1899) genetic interval (Figure 11).

Example 10: MiGA markers

Southern analysis of BAC clones 14, 24, 123 and 133 using markers 123R, 14L, or 24L as probes showed that these BAC clones contained several resistance gene analogs (RGAs). In view of the homology between the sequences of markers 14L, 24L and 123R with the *Mi1* gene from tomato, RGAs within the *Rpi-blb2* region are hereafter

referred to as *Mi* gene analogs (*Mi*GAs). In an attempt to develop additional polymorphic markers within the *Rpi-blb2* interval, PCR fragments generated from BAC clones 24 and 123 with the primer combination 14LR and 24LF were cloned into the pGEM-T vector (Promega, the Netherlands) and partially sequenced. Based on the alignment of these partial sequences, a set of universal primers were designed, univ14L and univ24L (Table 3), with the aim to amplify the corresponding region of as many as possible *Mi*GAs within the *Rpi-blb2* interval. This universal primer set was subsequently used to develop *Mi*GA specific SCAR/CAPS markers linked to *Rpi-blb2* (e.g. markers 14L24L, 24Lspec, 24L9spec; Figure 9).

10

Example 11: Physical mapping of the *S. bulbocastanum* derived *Rpi-blb2* locus.

The resistant clone Blb 2002 heterozygous for the *Rpi-blb2* locus, was used as source DNA for the construction of the *S. bulbocastanum* BAC library, hereafter referred to as the Blb 2002 BAC library. High molecular weight DNA preparation and BAC library construction were carried out as described previously. A total of approximately 100.000 clones were generated and stored as 50 bacterial pools containing approximately 2000 white colonies. These bacterial pools were generated by scraping the colonies from the agar plates into Luria Broth medium containing 18% glycerol and 12.5 µg/ml chloramphenicol using a sterile glass spreader. For the screening of the Blb 2002 BAC library, plasmid DNA was isolated from each pool of clones using the standard alkaline lysis protocol and PCR was carried out to identify positive pools. Bacteria corresponding to positive pools were diluted and plated on Luria Broth agar plates containing chloramphenicol (12.5 µg/ml). Individual white colonies were subsequently picked into 384-well microtiter plates and single positive BAC clones subsequently identified as described previously. Names of BAC clones isolated from the Blb 2002 BAC library carry the prefix BlbSP.

In order to build a Blb 2002 derived BAC contig across the *Rpi-blb2* genetic marker interval (69L-24L) the Blb 2002 BAC library was screened with markers 141R and 24L. This lead to the isolation of BAC clones BlbSP39 and BlbSP30, which overlap with each other and span the 141R-24L marker interval. BAC end sequences of both BAC clones were used to develop the markers SPB30L and SPB39L (Figure 9).

Example 12: Complementation analyses.

For complementation purposes, all *Rpi-blb2* gene candidates, i.e. all *Mi*GAs present on BAC clones BlbSP30, BlbSP39, 24, 242 and 211, were targeted for subcloning into the binary vector pBINPLUS (van Engelen *et al.*, 1996). This was done as follows. Aliquots of approximately 1 µg BAC DNA were digested with 1U, 0.1U or 0.01U of *Sau3A*I re-

40

striction enzyme for 30 min. The partially digested BAC DNA was subjected to contour-clamped homogeneous electric field (CHEF) electrophoresis at 4°C in 0.5 X TBE using a linear increasing pulse time of 1-10 sec and a field strength of 6 V/cm for 16 hr. After electrophoresis, the agarose gel was stained with ethidium bromide to locate the region of the gel containing DNA fragments of approximately 10kbp in size. This region was excised from the gel and treated with GELASE (Epicentre Technologies, USA) according to the manufacturer. The size selected DNA was ligated to the *Bam*HI-digested and dephosphorylated binary vector pBINPLUS (van Engelen *et al.*, 1995) followed by transformation to ElectroMAX *E. coli* DH10B competent cells (Life Technologies, UK).

5 Per BAC clone a total of 384 clones were PCR screened for the presence of *MiGA* sequences using the primers univ24L and univ14L (Table 3). Positive clones were selected for further characterisation. Based on the restriction pattern of the 14L24L fragments digested with the enzymes *Rsa*I, *Taq*I, *Alu*I, *Dpn*II or *Mse*I, the different groups of *MiGA*s were identified. The *MiGA* harbouring the marker 24L, which was completely present on BAC clones BlbSP39, 211 and 242 was not detected with the universal primers univ14L and univ24L.

10

15

The relative position of the *MiGA* sequences in the 10kbp subclones was determined by PCR using internal primers 123Mi and 14L2 for the 5' end and univ14L and 24L2 for the 3' end in combination with primers derived from pBINPLUS vector sequences (ARO 295 and 296; Table 3). Two subclones per RGA of each BAC-library were selected for transformation.

20

For complementation analysis, the selected subclones were transferred to the susceptible potato cultivars Impala and Kondor through *Agrobacterium* mediated transformation using isolate UIA143 (Farrand *et al.*, 1989) or AGLO (Lazo *et al.*, 1991). Primary transformants harbouring the transgenes of interest were tested for resistance to *P. infestans* in detached leaf assays using isolate IPO655-2A and IPO82001 (Table 4). Only the genetic constructs harbouring *RGC5*, both derived from *S. tuberosum* and *S. bulbocastanum*, were able to complement the susceptible phenotype both in cultivar Impala and in Kondor; in total 18 out of 19 *RGC5* containing primary transformants were resistant (Table 4, Figure 12) whereas all *RGC1*, *RGC2*, *RGC3*, *RGC4*, *RGC6* and *RGC7* containing primary transformants were susceptible to *P. infestans*. As the *RGC5* transformants showed similar resistance phenotypes as the resistant *S. bulbocastanum* parent of mapping population B6, *RGC5* was designated the *Rpi-blb2* gene. The homologues *RGC8* and *RGC24L* can also be transferred to the described susceptible potato cultivars and tested for resistance to *P. infestans* in a detached leaf assay.

25

30

35

A selection of primary transformants containing *RG5* was analysed for copy number by Southern analysis. *EcoRI* digested genomic DNA was hybridised with a *nptII* probe (Table 3). Based on the presence of the number of *nptII* hybridising fragments, the primary transformants contained at least 1 to 11 transgene inserts. In total, 4 single copy integrations in cultivar Impala and 6 in cultivar Kondor were observed of which one cultivar Kondor transformant appeared to have a *P. infestans* susceptible phenotype.

To investigate whether *Rpi-blb2* can also complement the susceptible phenotype in tomato, primary transformants of cultivar Moneymaker harbouring the *Rpi-blb2* gene construct can be produced and tested with a tomato isolate of *P. infestans* or with the potato derived isolates IPO82001 and IPO655-2A. The disease resistance assay can reveal that *Rpi-blb2* is able to complement a susceptible tomato phenotype.

Example 13: *Rpi-blb2* gene structure and putative amino acid sequence

The inserts of the *RG5* containing binary subclones 211F/C12 and SP39-20 were sequenced by a primer walk strategy whereby consecutive rounds of sequencing were carried out using a set of nested primers which were designed as the contiguous sequence was extended. The first set of sequences was generated using the M13F and M13R primers. The complete sequences of the inserts of clones 211F/C12 and SP39-20 consisted of 7967 and 9949 nucleotides (nt), respectively (Figure 13). The sequence of clone 211F/C12 was identical to the corresponding sequence within clone SP39-20. The position and putative structure of *Rpi-blb2* was predicted using GENSCAN (Burge and Karlin, 1997), GeneMark (Lukashin and Borodovsky 1998) and through alignment to the gene sequences of *Mi1.1* and *Mi1.2*.

The *Rpi-blb2* gene putatively contains two introns. Based on homology of these intron sequences to those present in *Mi1.1* and *Mi1.2* we predict intron 1 to be positioned within the 5' untranslated region (UTR), ending 32 nucleotides upstream of the putative ATG start codon. Intron 2 is predicted to be 86 nt long starting 43 nucleotides downstream of the ATG start codon of the gene (Figure 13). The coding sequence of the *Rpi-blb* transcript is predicted to be 3804 nt. The exact length of the transcript can be determined through 5' and 3' rapid amplification of cDNA ends (RACE) using the GeneRacer™ kit (Invitrogen™, Groningen, the Netherlands). 3' RACE is carried out with a set of nested *Rpi-blb2* specific oligonucleotides, which match sequences within the last 500 nt of the coding sequence of *Rpi-blb2* in combination with the GeneRacer™ 3' primer and GeneRacer™ 3' nested primers. 5' RACE is carried out on cDNA synthesised with an oligodT primer or a primer complementary to a *Rpi-blb2* specific sequence 500-1000 nt downstream of the ATG codon, using a set of nested *Rpi-blb2* specific oligonucleotides which are complementary to sequences within first 500 nt of

the coding sequence of *Rpi-blb2* in combination with the GeneRacer™ 5' primer and GeneRacer™ 5' nested primer.

The deduced open reading frame of the *Rpi-blb2* gene encodes a predicted polypeptide of 1267 amino acids with an estimated molecular weight of 146 kDa (Figure 14).

- 5 Several functional motifs present in *R* genes of the NBS-LRR class of plant *R* genes are apparent in the encoded protein. As illustrated in Figure 14, the *Rpi-blb2* protein belongs to the leucine zipper (LZ) subset of NBS-LRR resistance proteins. The N-terminal half of the *Rpi-blb2* protein contains a potential LZ region between amino acids 413 and 434 and six conserved motifs indicative of a nucleotide-binding site (van der
- 10 Blezen and Jones, 1998). The C-terminal half of *Rpi-blb* comprises a series of 15 irregular LRRs that can be aligned according to the consensus sequence $hxxhxxLxxLxxC/N/Sx(x)LxxLPxx$ observed in other cytoplasmic *R* proteins, whereby *h* can be L, I, M, V or F, and *x* any amino acid residue (Jones and Jones, 1997).

15 Example 14: Homology to known state of the art *R* gene sequences

- To identify *in silico* homologues of the *Rpi-blb2* gene, BLAST searches (Altschul *et al.*, 1990) were carried out with the coding sequence of the *Rpi-blb2* gene. BLASTN
- 20 searches identified a number of sequences with significant homology to the *Rpi-blb2* gene. Using the alignment programme ClustalW (standard settings) in the DNASTar software package, we determined that the *Rpi-blb2* coding sequence shares the highest homology to *Mi-1.1* (89.8%) and *Mi-1.2* (89.7%) (Genbank accession numbers AF039681 and AF039682, respectively). The latter sequence corresponds to the *Mi*
- 25 gene from tomato that confers resistance to three of the most damaging species of the root knot nematodes (*Meloidogyne* spp.) (Milligan *et al.*, 1998). In addition nucleotides 2410-3461 of the *Rpi-blb2* coding sequence share 87.8% sequence homology to a partial NBS-LRR sequence from *Solanum nigrum* (Genbank accession number AY055116.1). At the amino acid level the putative *Rpi-blb2* protein sequence shares the highest homology to *Mi-1.1* (82% identity) and *Mi-1.2* (81% identity) (Genbank accession numbers AF039681 and AF039682).
- 30

- Through ClustalW alignment of the deduced amino acid sequences of *Rpi-blb2*, *Mi-1.1* and *Mi-1.2* we have identified 200 amino acid (aa) residues which are unique to *Rpi-blb2* (Figure 15). Of these, 31 are found at hypervariable positions, i.e. the residue at
- 35 this position is different in all three sequences and 11 are encoded by small insertions (one 3 aa residue insertion and one 8 aa residue insertion). The rest are *Rpi-blb2* specific in that the aa residues encountered at corresponding positions in *Mi-1.1* and *Mi-1.2* are different from the *Rpi-blb2* residue but conserved in the two *Mi* protein sequences (Figure 15). Interestingly, the VLDL motif that is conserved in the third LRR of

Agrico B.V.

20030596

PF 54801

85

many NBS-LRR proteins including Mi (Axtell *et al.*, 2001; Banerjee *et al.*, 2001), is not conserved in *Rpi-blb2* (Figure 15).

Example 15: *Rpi-blb2* allele mining in wild *Solanum* species

5

Using primers ARF1F and ARF1R (Table 3B), designed around the start and stop codon of the *Rpi-blb2* gene, it is possible to amplify by PCR, alleles of *Rpi-blb2* from any *Solanum* species. The amplification products can be cloned between transcrip-

10 tional regulatory sequences in a binary plasmid and transferred to *S. tuberosum* through *Agrobacterium* mediated transformation or any method known to those skilled in the art. The resulting primary transformants can subsequently be analysed for resis-

tance to *P. infestans* or to any pathogen for which potato is a host plant.

Example 16: Material and methods

15

Plant material and development of mapping populations in (1) Solanum tuberosum.

Complex interspecific hybrid clones, designated ABPT, were made by Hermesen and co-workers (Hermesen, 1966; Hermesen and Ramanna, 1969; Ramanna and Hermesen, 1971; Hermesen and Ramanna, 1973; Hermesen, 1983; Hermesen, 1994) (Figure 1a).

20 The chromosome doubling step with colchicines was described by Hermesen (1966) and Hermesen and De Boer (1971). The resistance in some of the ABPT clones to *P. infestans* is believed to be derived from either one or both of the accessions from *S. bulbocastanum* BGRC 8007 (CGN 21306; Pi 275196) and BGRC 8008 (CGN 17693; Pi 275198) that were used in the initial cross to produce hybrids between

25 *S. acaule* and *S. bulbocastanum*, since all other parents that were used in the breeding scheme for ABPT-clones were susceptible or only partially resistant to *P. infestans* in detached leaf assays (Hermesen and Ramanna, 1973). Tubers from 19 clones of population [(ABPT clone number 55 x cultivar (cv) Oberambacher Frühe) x cv Arkula], from 7 clones of population [(ABPT clone number 55 x cv Oberambacher Frühe) x cv

30 Blanka] and from 5 clones of population [(ABPT clone number 60 x cv Alcmaria) x cv

Blanka] were received in 1988 from the former Department of Plant Breeding of the Wageningen Agricultural University (Wageningen, the Netherlands). Clones ARF 87-507, ARF 87-801 and ARF 87-601 were selected from these populations respectively.

35 They represented offspring from a second backcross (BC2) with the complex interspecific ABPT-clones and were used for further back crosses that resulted in one tetraploid BC3 population, two tetraploid BC4 populations and one diploid BC4 population that were used for genetic mapping of the *Rpi-blb2* gene (Figure 1). The tetraploid *Solanum tuberosum* mapping population ARG 95-15 was produced by crossing *P. infestans*

40 resistant clone ARF 87-507 with the susceptible cultivar Alkon. Tetraploid population

ARG 95-3 was produced by crossing *P. infestans* resistant clone AR 91-1263 with the susceptible cultivar Cosmos. Tetraploid population ARP 96-11 was produced by crossing resistant clone AR 92-1292 with the susceptible cultivar Celeste. The diploid population DP1 was obtained by crossing the resistant clone ARD 1197-16 with the susceptible clone ARD 93-2090 (Figure 1).

Plant material and development of mapping populations in (2) Solanum bulbocastanum.

- 10 The diploid *S. bulbocastanum* mapping population, designated B6 (synonym B6a. Blb 99-229, Blb 00-7 and Blb 00-8), was developed by crossing a *P. infestans* resistant clone Blb 2002 (synonym M94-81-C) with a susceptible clone Blb 48-5. Results from reciprocal crosses of population B6 were combined. The resistant parental clone of population B6 was obtained from a cross between *S. bulbocastanum* clone Blb 93-D26-3 (accession BGRC 8002; CGN 17690; PI 275187) as female parent and
- 15 *S. bulbocastanum* clone Blb 93-60-10 (accession BGRC 8006; PI 275194) as male parent. The susceptible parental clone of population B6 was obtained from a cross between *S. bulbocastanum* clones from accessions BGRC 8005 (CGN 17692, PI 275193) and BGRC 8006 (Figure 2).

20

Disease assays; (1) Phytophthora infestans isolates

- Three different *P. infestans* isolates were obtained from Plant Research International B.V. (Wageningen, the Netherlands). Isolates had different race structures and mating types as follows: IPO82001: race structure 1.2.3.4.5.6.7.10.11, mating type A2;
- 25 IPO655-2A: race structure 1.2.3.4.5.6.7.8.9.10.11, mating type A1; IPO428-2: race structure 1.2.3.4.5.6.7.8.9.10.11, mating type A2 (Flier *et al.*, 2003).

Disease assays; (2) field trials

- Glasshouse grown seedling tubers or field grown seed potatoes were planted at trial sites in Marknesse, the Netherlands from 1985 tot 2002, in the Toluca area of Mexico in 1991, or at a site in the Benguet Province in the Philippines in 1992. For individual clones, plots were planted consisting of 1 to 10 tubers. Approximately 8 weeks after planting, the field at Marknesse was inoculated with a sporangiospore solution of *P. infestans* isolate IPO82001 and disease scores were collected 3 to 6 weeks after
- 35 inoculation. Clones that were free or nearly free from late blight were classified as having a resistant phenotype whereas clones with a complete or nearly complete blighted foliage were classified as susceptible. Clones with intermediate reactions to late blight were classified as having an unknown phenotype. At the field trials in Mexico and the Philippines, natural infection had to occur. Once this natural infection by *P. infestans*
- 40 established, the percentage of blighted foliage of plants on each plot was scored on 8

Agrico.B.V.

20030596

PF 54801

87

and 6 days respectively on a 1-9 scale were estimated percentages of blighted foliage from 1 to 9 were: 0, 3, 10, 25, 50, 75, 90, 97 and 100 (Estrada-Ramos *et al.*, 1983).

Disease assays; (3) detached leaves

- 5 For the detached leaf assay, leaves from plants grown for 6 to 12 weeks in the greenhouse were placed in pieces of water-saturated florists foam, approximately 35x4x4 cm, and put in a tray (40 cm width, 60 cm length and 6 cm height) with a perforated bottom. Each leaf was inoculated with two droplets (25 µl each) of sporangiospore solution on the abaxial side. Subsequently, the tray was placed in a plastic bag on top of
- 10 a tray, in which a water-saturated filter paper was placed, and incubated in a climate room at 17°C and a 16h/8h day/night photoperiod with fluorescent light (Philips TLD50W/84HF and OSRAM L58W/21-840). After 6 to 9 days, the leaves were evaluated for the development of *P. infestans* disease symptoms.

Evaluation:

- 15 Plants with leaves that clearly showed sporulating lesions 6 to 9 days after inoculation were considered to have a susceptible phenotype, whereas plants with leaves showing no visible symptoms or necrosis at the side of inoculation in the absence of clear sporulation were considered to be resistant.

20 *Plant DNA marker screening*

- Genomic DNA was extracted from young leaves according to Bendahmane *et al.* (1997). For PCR analysis, 15 µl reaction mixtures were prepared containing 0.5 µg DNA, 15 ng of each primer, 0.2 mM of each dNTP, 0.6 units Taq-polymerase (15 U/µl, SphaeroQ, Leiden, the Netherlands), 10 mM Tris-HCl pH 9, 1.5 mM MgCl₂, 50 mM KCl,
- 25 0.1% Triton X-100 and 0.01% (w/v) gelatin. The PCRs were performed using the following cycle profile: 25 seconds DNA denaturation at 94°C, 30 seconds annealing and 40 seconds elongation at 72°C. As a first step in PCR-amplification DNA was denatured for 5 min at 94°C and finalised by an extra 5 min elongation step at 72°C. The amplification reactions were performed in a Biometra® T-Gradient or Biometra®
- 30 Uno-II thermocycler (Westburg, Leusden, the Netherlands). Depending on the marker, the PCR product was digested with an appropriate restriction enzyme. An overview of the markers including primer sequences, annealing temperature and restriction enzymes if appropriate, is given in Table 3. Subsequently, the (cleaved) PCR products were analysed by electrophoresis in agarose or acrylamide gels. For acrylamide gel
- 35 analysis, the CleanGel DNA Analysis Kit and DNA Silver Staining Kit (Amersham Pharmacia Biotech Benelux, Roosendaal, the Netherlands) were used.

Elongation of AFLP fragments by Thermal asymmetric interlaced (TAIL)-PCR

- Elongation of the sequence of an AFLP fragment was performed by TAIL-PCR according to Liu and Whittier (1995). Shortly, elongation of AFLP fragments was performed
- 40

Agrico B.V.

20030596

PF 54801

88

using 2 or 3 nested specific primers (sp) in combination with an arbitrary degenerate (AD) primer. The first PCR was performed with primers sp1 and AD, the second with sp2 and AD and the third with sp3 and AD according to the scheme described in Table 5. The PCR was performed in 25 µl reactions containing the standard PCR mix as described before, except that 30 ng of primer AD was used. The elongated fragments were cloned in pGEM-T (Promega, the Netherlands) and sequenced.

BAC library construction and screening

The resistant clone ARD 1197-16, heterozygous for the *Rpi-blb2* locus, was used as source DNA for the construction of the *S. tuberosum* BAC library. The resistant clone Blb 2002 heterozygous for the *Rpi-blb2* locus, was used as source DNA for the construction of the *S. bulbocastanum* BAC library. High molecular weight DNA preparation and BAC library construction were carried out as described in Rouppe van der Voort *et al.* (1999). For the *S. tuberosum* BAC library, approximately 120.000 clones with an average insert size of 100 kb, which corresponds to 8 to 10 genome equivalents were finally obtained. A total of approximately 70.000 clones were individually stored in 177 384-well microtiter plates at -80°C. Another 50.000 clones were stored as 14 bacterial pools containing approximately 4000 white colonies. These were generated by scraping the colonies from the agar plates into Luria Broth medium containing 18% glycerol and 12.5 µg/ml chloramphenicol using a sterile glass spreader. These so-called super pools were also stored at -80°C. Finally, another 37.000 clones were added to the *S. tuberosum* BAC library. The *S. bulbocastanum* BAC library consisted of 48 super pools of approximately 2.000 colonies.

Marker screening of the BAC library harbouring the individually stored BAC clones was carried out as described in Rouppe van der Voort *et al.* (1999). For the screening of the BAC library stored as super pools, plasmid DNA was isolated from each pool of clones using the standard alkaline lysis protocol and PCR was carried out to identify positive pools. Bacteria corresponding to positive pools were diluted and plated on Luria Broth agar plates containing chloramphenicol (12.5 µg/ml). Individual white colonies were subsequently picked into 384-well microtiter plates and single positive BAC clones subsequently identified as described above. Names of BAC clones isolated from the super pools carry the prefix SP (e.g. SP39).

Subcloning of candidate genes

Candidate RGAs were subcloned from BAC clone 24, 211, 242, BLBSP39 and BLBSP30 as follows. Aliquots of approximately 1 µg BAC DNA were digested with 1U, 0.1U or 0.01U of *Sau3A*I restriction enzyme for 30 min. The partially cleaved BAC DNA was subjected to CHEF electrophoresis at 4°C in 0.5 X TBE using a linear increasing pulse time of 1-10 sec and a field strength of 6 V/cm for 16 hr. After electrophoresis,

Agrico B.V.

20030596

89

the agarose gel was stained with ethidium bromide to locate the region of the gel containing DNA fragments of approximately 10 kbp in size. This region was excised from the gel and treated with GELASE (Epicentre Technologies, USA) according to the manufacturer. The size selected DNA was ligated to the *Bam*HI-cleaved and dephosphorylated binary vector pBINPLUS (van Engelen *et al.*, 1995) followed by transformation to ElectroMAX *E. coli* DH10B competent cells (Life Technologies, UK). A total of 192 clones were PCR screened for the presence of RGC sequences using the primers of marker 24L14L (Table 3). Positive clones were selected for further characterisation. Identification of clones harbouring *RGC1*, *RGC2*, *RGC3*, *RG4*, *RGC5*, *RGC6*, *RGC7*, *RGC8* and *RGC24L* was carried out by sequencing 14L24L PCR fragments derived from positive clones. The relative position of the RGAs within a subclone was determined by PCR analysis using internal primers (24L2, 123Mi) in combination with pBINPLUS specific primers (Table 3).

15 *Agrobacterium tumefaciens* mediated transformation of potato
Binary plasmids harbouring the candidate genes were transformed to *A. tumefaciens* strains AGL0 (Lazo *et al.*, 1991) or UA143 (Farrand *et al.*, 1989), the latter containing the helper plasmid pCH32 (Hamilton *et al.*, 1996). Overnight cultures of the transformed *A. tumefaciens* strains were used to transform potato tuber discs (cvs Impala and Kondor) according to standard protocols (Hoekema *et al.*, 1989; Fillati *et al.*, 1987).
20 Shortly, certified seed potatoes of cultivars Impala and Kondor were peeled and surface sterilised for 30 min in a 1% sodium hypochlorate solution containing 0.1% Tween-20. Tubers were then washed thoroughly in large volumes of sterile distilled water (4 times, 10 min). Discs of approximately 2 mm thickness and 7 mm in diameter
25 were sliced from cylinders of tuber tissue prepared with a corkbore. The tuber discs were transferred into liquid MS30 medium containing *A. tumefaciens* and incubated for 15 min. After removing the *A. tumefaciens* solution, the tuber discs were transferred to regeneration medium containing MS30, 0.9 mg/l IAA, 3.6 mg/l zeatine riboside and 8 g/l agar (Hoekema *et al.*, 1989). The plates were incubated at 24°C, 16 hour day-length (Philips TLD50W/84HF). After 48 hours of co-cultivation, the tuber discs were
30 rinsed for 5 min in liquid MS medium including antibiotics, 200 mg/l vancomycin, 250 mg/l cefotaxim and 75 mg/l kanamycin, and transferred to regeneration medium supplemented with the same antibiotics. The plates were incubated at 24°C, 16 hour day-length (Philips TLD50W/84HF). Every three weeks, the tuber discs were transferred to fresh medium. Regenerating shoots were transferred to MS30 medium containing 75 mg/l kanamycin. Rooting shoots were propagated *in vitro* and tested for absence of *A. tumefaciens* cells by incubating a piece of stem in 3 ml Luria Broth medium (3 weeks, 37°C, 400 rpm). One plant of each transformed regenerant was transferred
40 to the greenhouse.

Agrobacterium tumefaciens mediated transformation of tomato

- Seeds of the susceptible tomato line Moneymaker are rinsed in 70% ethanol to dissolve the seed coat and washed with sterile water. Subsequently, the seeds are surface-sterilised in 1.5% sodium hypochlorite for 15 minutes, rinsed three times in sterile water and placed in containers containing 140 ml MS medium pH 6.0 (Murashige and Skoog, 1962) supplemented with 10 g/l sucrose (MS10) and 160 ml vermiculite. The seeds are left to germinate for 8 days at 25°C and 0.5 W/M² light.

- Eight day old cotyledon explants are pre-cultured for 24 hours in Petri dishes containing a two week old feeder layer of tobacco suspension cells plated on co-cultivation medium (MS30 pH 5.8 supplemented with Nitsch vitamins (Duchefa Biochemie BV, Haarlem, the Netherlands), 0.5 g/l MES buffer and 8 g/l Daichin agar).

- Overnight cultures of *A. tumefaciens* are centrifuged and the pellet is resuspended in cell suspension medium (MS30 pH 5.8 supplemented with Nitsch vitamins, 0.5 g/l MES buffer, pH 5.8) containing 200 µM acetosyringone to a final O.D.₆₀₀ of 0.25. The explants are then infected with the diluted overnight culture of *A. tumefaciens* UIA143 containing pBINRGCS for 25 minutes, blotted dry on sterile filter paper and co-cultured for 48 hours on the original feeder layer plates. Culture conditions are as described above.

- Following the co-cultivation, the cotyledons explants are transferred to Petri dishes with selective shoot inducing medium (MS pH 5.8 supplemented with 10 g/l glucose, including Nitsch vitamins, 0.5 g/l MES buffer, 5 g/l agargel, 2 mg/l zeatine riboside, 400 mg/l carbenicilline, 100 mg/l kanamycine, 0.1 mg/l IAA) and cultured at 25°C with 3-5 W/m² light. The explants are sub-cultured every 3 weeks onto fresh medium. Emerging shoots are dissected from the underlying callus and transferred to containers with selective root inducing medium (MS10 pH 5.8 supplemented with Nitsch vitamins, 0.5 g/l MES buffer, 5 g/l agargel, 0.25 mg/l IBA, 200 mg/l carbenicillin and 100 mg/l kanamycine).

RNA extraction

- Total RNA can be isolated using Trizol[®] according to the protocol supplied by the manufacturer (Invitrogen[™], Groningen, the Netherlands) with minor modifications. Briefly, 0.5 g of young leaf tissue is ground in liquid nitrogen and the powder suspended in 5 ml Trizol[®]. After a 5 min incubation at room temperature (RT), 0.5 ml chloroform is added, the suspension is vortexed and incubated for 2 min. After centrifugation (15 min, 11404 x g, 4°C) the supernatant is transferred to a new tube and 2.5 ml isopropanol is added. After 10 min at RT, nucleic acids is precipitated (10 min, 11404 x g, 4°C). The pellet is washed with 5 ml 70% ethanol (5 min, RT) and after cen-

trifugation (5 min, 6415 x g, 4°C), the pellet was dried and resuspended in 100 µl sterile distilled water.

PolyA RNA can be extracted from total RNA using the Oligotex™ kit (Qiagen, GmbH, Germany).

5

Rapid amplification of cDNA ends

The 5' and 3' ends of the *Rpi-blb2* cDNA and confirmation of putative intron positions can be determined by rapid amplification of cDNA ends (RACE) using the GeneRacer™ kit (Invitrogen™, Groningen, The Netherlands). 3' RACE is carried out with a set of nested *Rpi-blb2* specific oligonucleotides which match sequences within the last 500 nt of the coding sequence of *Rpi-blb2* in combination with the GeneRacer™ 3' primer and GeneRacer™ 3' nested primers. 5' RACE is carried out on cDNA synthesised with an oligodT primer or a primer complementary to a *Rpi-blb2* specific sequence 500-1000 nt downstream of the ATG codon, using a set of nested *Rpi-blb2* specific oligonucleotides which are complementary to sequences within first 500 nt of the coding sequence of *Rpi-blb2* in combination with the GeneRacer™ 5' primer and GeneRacer™ 5' nested primer.

AFLP fingerprinting and cloning of AFLP fragments

Template preparation and AFLP fingerprinting were essentially performed as described in Vos *et al.* (1995). In order to clone specific fragments ³²P-labelled AFLP fragments were excised out of the acrylamide gel by overlaying the polyacrylamide gels, dried on Whatmann 3MM paper, with autoradiogram images. The pieces of gel/paper underneath the band of interest were cut out and transferred to 200 µl of TE and incubated for 1 h at room temperature. Five microlitres of supernatant was used to re-amplify the fragment, using a PCR in which the *EcoRI*+0 in combination with *MseI*+0 were used as primers. The re-amplified AFLP fragment was subsequently cloned into the pGEM-T cloning vector (Promega, the Netherlands) and the inserts of several clones sequenced.

30

The DNA sequence of the excised AFLP band was used to design locus-specific primers. The amplification product obtained with such primers was screened for internal polymorphisms with restriction enzymes. After restriction, the fragments were separated on a 2-3% agarose gel including ethidiumbromide.

35

RGA-AFLP analysis

Template preparation was essentially performed as described in Vos *et al.* (1995). However, the second amplification step was carried out with the P-loop based primer S1 from Leister *et al.* (1996) in combination with the *EcoRI*+0 AFLP primer. A 10 µl reaction mixture [0.5 µl ³²P-labelled S1 primer (10 ng/µl); 0.5 µl *EcoRI*+0 primer

40

92

- (10 ng/ μ l); 0.8 μ l dNTPs (5mM); 2 μ l 10xGoldstarTM PCR buffer (Eurogenetc, Belgium); 1.2 μ l $MgCl_2$ (25.mM); 0.06 μ l GoldstarTM DNA polymerase (5U/ μ l) (Eurogentec, Belgium); 14.94 μ l MQ water] was added to 10 μ l diluted template (20x diluted in MQ water) and a PCR reaction performed using the following cycle profile: 45 seconds
- 5 DNA denaturation at 94°C, 45 seconds primer annealing at 49°C and 2 min elongation step at 72°C (35 cycles). Prior to the cycling the template DNA was denatured for 2 min at 94°C and the PCR was finalised by a applying an extra 5 min elongation step at 72°C. The amplification reactions were performed in a Perkin Elmer 9600 thermocycler. The labelled PCR products fragments were separated on a 6% polyacrylamide gel and
- 10 the individual bands visualized by autoradiography according to standard procedures.

References

- Altschul, S.F., Gish, W., Miller, W., Myers, E.W., and Lipman, D.J. (1990) Basic local alignment search tool. *Journal of Molecular Biology* 215, 403-410.
- 5 Axtell, M.J., McNellis, T.W., Mudgett, M.B., Hsu, C.S., and Staskawicz, B.J. (2001) Mutational analysis of the *Arabidopsis* RPS2 disease resistance gene and the corresponding *Pseudomonas syringae* avrRpt2 avirulence gene. *Molecular Plant-Microbe Interactions* 14, 181-188.
- 10 Balivora, A., Ercolano, M.R., Weis, J., Meksem, K., Börmann, C.A., Oberhagen, P., Salamini, F., and Gebhardt, C. (2002) The R1 gene for potato resistance to late blight (*Phytophthora infestans*) belongs to the leucine zipper/NBS/LRR class of plant resistance genes. *Plant Journal* 30, 361-371.
- 15 Banerjee, D., Zhang, X., and Bent, A.F. (2001) The leucine-rich repeat domain can determine effective interaction between RPS2 and other host factors in *Arabidopsis* RPS2-mediated disease resistance. *Genetics* 158, 439-450.
- 20 Bendahmane, A., Kanyuka, K., and Baulcombe, D.C. (1997) The High-resolution genetic and physical mapping of the Rx gene for extreme resistance to potato virus X in tetraploid potato. *Theoretically Applied Genetics* 95, 153-162.
- 25 Burge, C.B. and Karlin, S. (1997) Prediction of complete gene structures in human genomic DNA. *Journal of Molecular Biology* 268, 78-94.
- Colon, L.T., Turkensteen, L.J., Prummel, W., Budding, D.J. and Hoogendoorn, J. (1985) Durable resistance to late blight (*Phytophthora infestans*) in old potato cultivars. *European Journal of Plant Pathology* 101, 387-397.
- 30 Dangl, J.L. and Jones, J.D.G. (2001) Plant pathogens and integrated defence responses to infection. *Nature* 411, 826-833.
- 35 Estrada-Ramos, N., Pérez-Alvarez, O., Henfling, J. and Malamud, O. (1983) In: W.J. Hooker (ed), *Research for the potato in the year 2000*. International Potato Center, Lima, Peru, pp 78-79.
- Farrand, S.K., O'Morchoe, S.P., and McCutchan, J. (1989) Construction of an *Agrobacterium tumefaciens* C58 *recA* mutant. *Journal of Bacteriology* 171, 5314-5321.
- 40

Fillati, J.J., Kiser, J., Rose, R., and Comai, L. (1987) Efficient transfer of a glyphosate tolerance gene into tomato using a binary *Agrobacterium tumefaciens* vector. *Bio Technology* 5, 726-730.

- 5 Flier, W.G. (2001) Variation in *Phytophthora infestans*, sources and implications. PhD thesis Wageningen University, Wageningen, the Netherlands, p. 93.

Flier, W.G., vandenBosch, G.B.M., and Turkensteen, L.J. (2003) Stability and partial resistance in potato cultivars exposed to aggressive strains of *Phytophthora infestans*.

- 10 Plant Pathology 52 (3), 326-337.

Gebhardt, C., Ritter, E., Barone, A., Debener, T., Walkemeier, B., Schachtschabel, U., Kaufmann, H., Thompson, R.D., Bonierbale, M.W., Ganai, M.W., Tanksley, S.D., and Salamini, F. (1991) RFLP maps of potato and their alignment with the homoeologous tomato genome. *Theoretically Applied Genetics* 83, 49-57.

- 15

Hamilton, C.M., Frary, A., Lewis, C., and Tanksley, S.D. (1996) Stable transfer of intact high molecular weight DNA into plant chromosomes. *Proceedings of National Academy of Science USA* 93, 9975-9979.

20

Hermesen, J.G.Th. (1966) Crossability, fertility and cytogenetic studies in *Solanum acaule* x *Solanum bulbocastanum*. *Euphytica* 15, 149-155.

Hermesen, J.G.Th. and Ramanna, M.S. (1969) Meiosis in different F₁-hybrids of

- 25 *Solanum acaule* Bitt. x *S. bulbocastanum* Dun. and its bearing on genome relationship, fertility and breeding behaviour. *Euphytica* 18, 27-35.

Hermesen, J.G.Th., and De Boer, A.J.E. (1971) The effect of colchicine treatments on *Solanum acaule* and *S. bulbocastanum*; a complete analysis of ploidy chimeras in

- 30 *S. bulbocastanum*. *Euphytica* 20, 171-180

Hermesen, J.G.Th. and Ramanna, M.S. (1973) Double-bridge hybrids of *Solanum bulbocastanum* and cultivars of *Solanum tuberosum*. *Euphytica* 22, 457-466

- 35 Hermesen, J.G.Th. and Verdenius, J. (1973) Selection from *Solanum tuberosum* group Phureja of genotypes combining high-frequency haploid induction with homozygosity for embryo-spot. *Euphytica* 22, 244-259.

- Hermesen, J.G.Th. (1983) Utilization of wide crosses in potato breeding. In: *Report of a planning conference on present and future strategies for potato breeding and improvement*. International Potato Center, Lima, Peru, pp 115-132.
- 5 Hermesen, J.G.Th. (1994) Introgression of genes from wild species, including molecular and cellular approaches. In: J.E. Bradshaw and G.R. Mackay (eds), *Potato Genetics*, CAB International, Wallingford, UK. pp 515-538.
- 10 Hijmans, R.J., Forbes, G.A., and Walker, T.S. (2000) Estimating the global severity of potato late blight with GIS-linked disease forecast models. *Plant Pathology* 49, 697-705.
- 15 Hoekema, A., Hirsch, P.R., Hooykaas, P.J.J., and Schilperoort, R.A. (1983) A binary plant vector strategy based on separation of vir- and T-region of the *Agrobacterium tumefaciens* Ti-plasmid. *Nature* 303, 179-180.
- 20 Hoekema, A., Huisman, M.J., Molendijk, L., van den Elzen, P.J.M., and Cornelissen, B.J.C. (1989) The genetic engineering of two commercial potato cultivars for resistance to potato virus X. *Bio/Technology* 7, 273-278.
- Jones, D.A. and Jones, J.D.G. (1997) The role of leucine-rich repeat proteins in plant defenses. *Adv. Bot. Res.* 24, 89-167.
- 25 Lazo, G.R., Stein, P.A. and Ludwig, R.A. (1991) A DNA transformation-competent *Arabidopsis* genomic library in *Agrobacterium*. *Bio/Technology* 9, 963-967.
- 30 Leach, J.E., Vera Cruz, C.M., Bai, J., and Leung, H. (2001) Pathogen fitness penalty as a predictor of durability of disease resistance genes. *Annual Review Phytopathology* 39, 187-224.
- Leister, D., Ballvora, A., Salamini, F., and Gebhardt, C. (1996) A PCR-based approach for isolating pathogen resistance genes from potato with potential for wide application in plants. *Nature Genetics* 14, 421-429.
- 35 Liu, Y-G., and Whittier, R.F. (1995) Thermal asymmetric interlaced PCR: Automatable amplification and sequencing of insert end fragments from P1 and YAC clones for chromosome walking. *Genomics* 25, 674-682.
- 40 Lukashin A.V. and M. Borodovsky (1998) GeneMark.hmm: new solutions for gene finding. *Nucleic Acids Research* 26, 1107-1115.

5 Milligan, S.B., Bodeau, J., Yaghoobi, J., Kaloshian, I., Zabel, P. and Williamson, V.M. (1998) The root-knot nematode resistance gene *Mi* from tomato is a member of the leucine zipper nucleotide binding leucine-rich repeat family of plant genes. *Plant Cell* 10, 1307-1319.

10 Nombela, G., Williamson, V.M., and Muniz, M. (2003) The root-knot nematode resistance gene *Mi-1.2* of tomato is responsible for resistance against the whitefly *Bemisia tabaci*. *Molecular Plant Microbe Interactions* 16 (7), 645-649.

15 Ramana, M.S. and Hermesen, J.G.Th. (1971) Somatic chromosome elimination and meiotic chromosome pairing in the triple hybrid 6x-(*Solanum acaule* x *S. bulbocastanum*) x 2x-*S. phureja*. *Euphytica* 20, 470-481

20 Rossi, M., Goggin, F.L., Milligan, S.B., Kaloshian, I., Ullman, D.E., Williamson, V.M. (1998) The nematode resistance gene *Mi* of tomato confers resistance against the potato aphid. *Proceedings of the National Academy of Science USA* 95, 9750-9754.

25 Rouppe van der Voort, J., Kanyuka, K., van der Vossen, E., Bendahmane, A., Mooijman, P., Klein-Lankhorst, R., Stiekema, W., Baulcombe, D. & Bakker, J. (1999) Tight physical linkage of the nematode resistance gene *Gpa2* and the virus resistance gene *Rx* on a single segment introgressed from the wild species *Solanum tuberosum* subsp. *Andigena* CPC 1673 into cultivated potato. *Molecular Plant Microbe Interactions* 12, 197-206.

30 Schepers, H. and Wustman, R. (2003) *Phytophthora* 2003: middelen en aanpak. *Informa* 6/juni 2003.

35 Simons, G., Groenendijk, J., Wijbrandi, J., Reijans, M., Groenen, J., Diergaarde, P., Van der Lee, T., Bleeker, M., Onstenk, J., de Both, M., Haring, M., Mes, J., Cornelissen, B., Zabeau, M and Vos, P. (1998) Dissection of the *Fusarium* I2 gene cluster in tomato reveals six homologues and one active gene copy. *Plant Cell* 10, 1055-1068.

40 Stam, P. (1993) Construction of integrated genetic linkage maps by means of a new computer package: Joinmap. *Plant Journal* 3, 739-744.

Song, J., Bradeen, J.M., Naess, S.K., Raasch, J.A., Wielgus, S.M., Haberlach, G.T., Liu, J., Kuang, H., Austin-Phillips, S., Buell, C.R., Helgeson, J.P. and Jiang, J. (2003) Gene *RB* cloned from *Solanum bulbocastanum* confers broad spectrum resistance to potato late blight. *Proceedings of National Academy of Science USA* 100, 9128-9133.

van der Blezen, E.A. and Jones, J.D.G. (1998) The NB-ARC domain: a novel signalling motif shared by plant resistance gene products and regulators of cell death in animals. *Current Biology* 8, 226-227.

5

van der Vossen, E., Sikkema, A., te Lintel Hekkert, B., Gros, J., Stevens, P., Muskens, M., Wouters, D., Pereira, A., Stiekema, W., and Allefs, J. (submitted) An ancient R gene from the wild potato species *Solanum bulbocastanum* confers broad-spectrum resistance to *Phytophthora infestans* in cultivated potato and tomato.

10

van Eck, H.J., Rouppe van der Voort, J.N.A.M., Draaistra, J., van Zandvoort, P., van Enckevort, E., Segers, B., Peleman, J., Jacobsen, E., Helder, J., and Bakker, J. (1995) The inheritance and chromosomal localization of AFLP markers in a non-inbred potato offspring. *Molecular Breeding* 1: 397-410.

15

van Engelen, F.A., Molthoff, J.W., Conner, A.J., Nap, J-P., Pereira, A. and Stiekema, W.J. (1995) pBINPLUS: an improved plant transformation vector based on pBIN19. *Transgenic Research* 4, 288-290.

20

Vos, P., Hogers, R., Bleeker, M., Rijans, M., Van der Lee, T., Hornes, M., Frijters, A., Pot, J., Peleman, J., Kuiper, M. and Zabeau, M. (1995) AFLP: a new technique for DNA fingerprinting. *Nucleic Acids Research*, 23, 4407-4414.

25

Vos, P., Simons, G., Jesse, T., Wijbrandi, J., Heinen, L., Hogers, R., Frijters, A., Groenendijk, J., Diergaarde, P., Reijans, M., Fierens-Onstenk, J., de Both, M., Peleman, J., Liharska, T., Hontelez, J., and Zabeau, M. (1998) The tomato Mi-1 gene confers resistance to both root-knot nematodes and potato aphids. *Nature Biotechnology* 16 (13), 1365-1369.

Claims

1. A method for generating or increasing the resistance of a plant to a plant patho-
gen of the phylum Oomyceta comprising increasing the activity of Rpi-blb2 pro-
tein in the plant or a tissue, organ or cell of a plant or a part thereof.
2. The method of claim 1, wherein said Rpi-blb2 protein is encoded by a poly-
nucleotide comprising a nucleic acid molecule selected from the group consisting
of:
- (a) nucleic acid molecule encoding at least the mature form of the polypeptide
depicted in SEQ ID NO: 2 or 4;
 - (b) nucleic acid molecule comprising the coding sequence as depicted in SEQ
ID NO: 1 or 3 or 5 or 6 encoding at least the mature form of the polypeptide;
 - (c) nucleic acid molecules the nucleotide sequence of which is degenerate as a
result of the genetic code to a nucleotide sequence of (a) or (b);
 - (d) nucleic acid molecule encoding a polypeptide derived from the polypeptide
encoded by a polynucleotide of (a) to (c) by way of substitution, deletion
and/or addition of one or several amino acids of the amino acid sequence of
the polypeptide encoded by a polynucleotide of (a) to (c);
 - (e) nucleic acid molecule encoding a polypeptide the sequence of which has an
identity of 70% or more to the amino acid sequence of the polypeptide en-
coded by a nucleic acid molecule of (a) or (b);
 - (f) nucleic acid molecule comprising a fragment or a epitope-bearing portion of
a polypeptide encoded by a nucleic acid molecule of any one of (a) to (e);
 - (g) nucleic acid molecule comprising a polynucleotide having a sequence of a
nucleic acid molecule amplified from a nucleic acid library using a primer as
listed in Tab. 3b;
 - (h) nucleic acid molecule encoding a fragment beginning with amino acid: 1,
30, 50, 100, 200, 300, 500, or 1000 and stopping with amino acid 1276,
1000, 500, 300, 200, 50, or 1 of a polypeptide encoded by any one of (a) to
(g);
 - (i) nucleic acid molecule comprising at least 20 nucleotides of a polynucleotide
of any one of (a) or (d);
 - (j) nucleic acid molecule encoding a polypeptide being recognized by a mono-
clonal antibody that have been raised against a polypeptide encoded by a
nucleic acid molecule of any one of (a) to (h);
 - (k) nucleic acid molecule obtainable by screening an appropriate library under
stringent conditions with a probe having the sequence of the nucleic acid
molecule of any one of (a) to (j) or of a fragment thereof of at least 20; and

99

(l) nucleic acid molecule the complementary strand of which hybridises under stringent conditions with a nucleic acid molecule of any one of (a) or (k);

or the complementary strand of any one of (a) to (l);

5 or expressing a polypeptide encoded by a segment of chromosome or linkage group 6 of *Solanum bulbocastanum* or *Solanum tuberosum* which co-segregates with a marker selected from table 3a or 3b and which mediates resistance to a pathogen of the phylum Oomyceta
10 and whereby the polynucleotide has not the sequence of Mi1.1 or Mi1.2 as depicted in Seq. ID NO.: 7 or 9.

3. The method of claim 1 or 2, wherein the activity of a further resistance protein is increased.

15 4. The method of any one of claims 1 to 3, wherein activity is increased due to a de novo-expression.

5. The method of any one of claims 1 to 4, wherein the endogenous activity of a Rpi-blb2 and/or the further resistance protein is increased.

20 6. The method of any one of claim 1 to 5, comprising one or more of the following steps

- 25 a) stabilizing the resistance protein;
b) stabilizing the resistance protein encoding mRNA;
c) increasing the specific activity of the resistance protein;
d) expressing or increasing the expression of a homologous or artificial transcription factor for resistance protein expression;
e) stimulate resistance protein activity through exogenous inducing factors;
30 f) expressing a transgenic resistance protein encoding gene; and/or
g) increasing the copy number of the resistance protein encoding gene.

7. The method of any one of claims 1 to 6 which results in reduction in the sporulation index of at least 30% after infection with *P. infestans* compared to a wild
35 type.

Agrico B.V.**20030596****PF 54801****100**

8. A polynucleotide encoding a Rpi-blb2 protein comprising a nucleic acid molecule selected from the group consisting of:

- 5 (a) nucleic acid molecule encoding at least the mature form of the polypeptide depicted in SEQ ID NO: 2 or 4;
- (b) nucleic acid molecules comprising the coding sequence as depicted in SEQ ID NO: 1 or 3 or 5 or 6 encoding at least the mature form of the polypeptide;
- (c) nucleic acid molecule the nucleotide sequence of which is degenerate as a result of the genetic code to a nucleotide sequence of (a) or (b);
- 10 (d) nucleic acid molecule encoding a polypeptide derived from the polypeptide encoded by a polynucleotide of (a) to (c) by way of substitution, deletion and/or addition of one or several amino acids of the amino acid sequence of the polypeptide encoded by a polynucleotide of (a) to (c);
- 15 (e) nucleic acid molecule encoding a polypeptide the sequence of which has an identity of 70% or more to the amino acid sequence of the polypeptide encoded by a nucleic acid molecule of (a) or (b);
- (f) nucleic acid molecules comprising a fragment or a epitope-bearing portion of a polypeptide encoded by a nucleic acid molecule of any one of (a) to (e);
- 20 (g) nucleic acid molecule comprising a polynucleotide having a sequence of a nucleic acid molecule amplified from a nucleic acid library using the primers as listed in Tab.3b;
- (h) nucleic acid molecule encoding polypeptide fragment beginning with amino acid: 1, 30, 50, 100, 200, 300, 500, or 1000 and stopping with amino acid 1276, 1000, 500, 300, 200, 50, or 30 of a polypeptide encoded by any one of (a) to (g);
- 25 (i) nucleic acid molecule comprising at least 20 nucleotides of a polynucleotide of any one of (a) or (d);
- (j) nucleic acid molecule encoding a polypeptide being recognized by a monoclonal antibodies that have been raised against a polypeptide encoded by a nucleic acid molecule of any one of (a) to (h);
- 30 (k) nucleic acid molecule obtainable by screening an appropriate library under stringent conditions with a probe having the sequence of the nucleic acid molecule of any one of (a) to (j) or of a fragment thereof of at least 20; and
- 35 (l) nucleic acid molecule the complementary strand of which hybridises under stringent conditions with a nucleic acid molecule of any one of (a) or (k);

or the complementary strand of any one of (a) to (l);

or encoding a polypeptide encoded by a segment of chromosome or of linkage group 6 of *Solanum bulbocastanum* or *Solanum tuberosum* which co-segregates with a marker selected from table 3a or 3b or comprises a replication site or

40

101

hybridisation site for said marker and which mediates resistance to pathogens of the phylum Oomyceta;

and whereby the polynucleotide has not the sequence of Mi1.1 or Mi1.2 as depicted in Seq. ID NO.: 7 or 9.

- 5
9. The polynucleotide of claim 8 or the method of any one of claims 1 to 7, wherein the marker is E40M58, CT119, or CT216.
- 10
10. The polynucleotide of claim 8 to 9 which is DNA or RNA.
11. A method for making a recombinant vector comprising inserting the polynucleotide of any one of claims 8 to 10 into a vector or inserting said polynucleotide and a further resistance protein.
- 15
12. A vector containing the polynucleotide of any one of claims 8 to 10 or comprising said polynucleotide and a further resistance gene or being produced by the method of claim 11.
- 20
13. The vector of claim 12 or the method of any one of claims 1 to 7 in which a polynucleotide encoding Rpi-blb2 protein or encoding the further resistance protein is operatively linked to expression control sequences and/or is operatively linked to a nucleic acid sequence encoding a transgenic expression regulating signal allowing expression in prokaryotic or eukaryotic host cells.
- 25
14. The vector of claim 12 or 13 or the method of any one of claims 1 to 7 in which the polynucleotide encoding Rpi-blb2 protein or encoding a further resistance protein is operatively linked to expression control sequences of the same species origin as the polynucleotide encoding Rpi-blb2 protein or the further resistance protein.
- 30
15. A method of making a recombinant host cell comprising introducing the vector of any one of claims 12 to 14 or introducing said vector and a vector for expressing a further resistance protein into a host cell.
- 35
16. A host cell produced according to the method of claim 15 or genetically engineered with the polynucleotide of any one of claims 8 to 10 or the vector of any one of claims 12 to 14 or genetically engineered with said vector or polynucleotide and a vector or a polynucleotide for expressing a further resistance protein.

Agrico B.V.**20030596****PF 54801****102**

17. The host cell of claim 16, which is *E. coli*, Baculovirus, Agrobacterium, or a plant cell.
- 5 18. A process for the production of a Rpi-blb2-polypeptide comprising culturing the host cell of claim 16 or 17 and recovering the polypeptide encoded by said polynucleotide and expressed by the host cell from the culture or the host cells.
- 10 19. A polypeptide having the amino acid sequence encoded by a polynucleotide of any one of claims 8 to 10 or obtainable by the process of claim 18.
20. A polypeptide having Rpi-blb2 activity.
21. An antibody that binds specifically to the polypeptide of claim 19 or 20.
- 15 22. An antisense nucleic acid molecule comprising the complementary sequence of the polynucleotide of any one of claims 8 to 10.
- 20 23. A method for the production of a transgenic plant, plant cell or plant tissue or a part thereof comprising the introduction of the polynucleotide of any one of claims 8 to 10 or said polynucleotide and a polynucleotide encoding a further resistance protein, or the vector of any one of claims 12 to 14 into the genome of said plant, plant tissue or plant cell or a part thereof.
- 25 24. A plant cell comprising the polynucleotide of any one of claims 8 to 10, the vector of any one of claims 12 to 14 or obtainable by the method of claim 23.
- 30 25. A transgenic plant or plant tissue or a part thereof comprising the plant cell of claim 24.
- 35 26. A method for producing a plant or a part thereof resistant to a plant pathogen of the phylum Oomyceta comprising the step:
expressing in the plant or a part thereof the polypeptide of claim 19 or 20 and a further resistance protein.
27. A method for producing a plant or a part thereof with a durable resistance to a *Phytophthora* sp. comprising co-expressing in the plant or a part thereof Rpi-blb and Rpi-blb2 protein or the polypeptide of claim 19 or 20.

Agrico B.V.**20030596****PF 54801****103**

28. The transgenic plant or plant tissue of claim 25 or produced according to claim 26 or 27, which upon the presence of the polynucleotide or the vector is resistant to a plant pathogen of the phylum Oomyceta.
- 5 29. Harvestable parts of the transgenic plant or plant tissue of claim 25 comprising the plant cell of claim 24.
30. Propagation material of the transgenic plant or plant tissue of claim 25 comprising the plant cell of claim 24.
- 10 31. Use of the polynucleotide of any one of claims 8 to 10, the vector of any one of claims 12 to 14, or the polypeptide of claim 19 or 20 for producing a plant or a plant tissue, plant organ, or a plant cell or a part thereof resistant to a plant pathogen of the phylum Oomyceta.
- 15 32. A method for the identification of an compound stimulating resistance to a plant pathogen of the phylum Oomyceta comprising:
- 20 (a) contacting cells which express the polypeptide of claim 19 or 20 or its mRNA with a candidate compound under cell cultivation conditions;
- (b) assaying an increase in expression of said polypeptide or said mRNA;
- (c) comparing the expression level to a standard response made in the absence of said candidate compound; whereby, an increased expression over the standard indicates that the compound is stimulating resistance.
- 25 33. Use of the polynucleotide of any one of claims 8 to 10, the vector of any one of claims 12 to 14, the polypeptide of claim 19 or 20 or the antibody of claim 21, for identifying and/or producing compounds activating or stimulating plant resistance to a plant pathogen of the phylum Oomyceta.
- 30 34. A diagnostic composition, comprising the polynucleotide of any one of claims 8 to 10, the vector of any one of claims 12 to 14, the antibody of claim 21 or the antisense nucleic acid of claim 22 and optionally suitable means for detection.
- 35 35. A kit comprising the polynucleotide of any one of claims 8 or 12, the vector of any one of claims 12 to 14, the host cell of claim 16 or 17, the polypeptide of claim 19 or 20, the antisense nucleic acid of claim 22, the antibody of claim 21, the plant cell of claim 24, the plant or plant tissue of claim 25, the harvestable part of claim 29, or the propagation material of claim 30 and optionally a polynucleotide encoding Rpi-blb, Rpi-blb protein or an antibody against Rpi-blb.
- 40

Agrico B.V.

20030596

PF 54801

104

36. A method for the production of a plant crop protectant providing the polynucleotide of any one of claims 8 to 10, the vector of any one of claims 12 to 14 or the polypeptide of claim 19 or 20 or comprising the steps of the method of claim 32; and formulating the polynucleotide of any one of claims 8 to 10, the vector of of claims 12 or 14 or the polypeptide of claim 19 or 20 or the compound identified in step (c) of claim 32 in a form applicable as agricultural composition.
37. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 36, wherein the plant pathogen is of the order Pythiales or Peronosperales.
38. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 37, wherein the plant pathogen is of the species *P. infestans*, *Phytophthora erythroseptica*, *Phytophthora capsici*, *Phytophthora sojae*, *Phytophthora parasitica* var. *nicotianae*, *Bremia lactuca*, *Peronospera tabaci* or *Plasmopara viticola*.
39. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 38, wherein the resistance protein is characterized by a P-loop and a NBS domain.
40. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 39, wherein the further resistance gene is a gene encoding Rpi-blb, R1, R-ber, Rpi1, R2, R3, R4, R5, R6, R7, R8, R9, R10, R11, Ph-1, Ph-2 and/or Ph-3.
41. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 40, wherein the further resistance protein is the Rpi-blb protein.
42. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 41 wherein the plant, plant cell or plant tissue is selected from the group consisting of Menyanthaceae, Solanaceae, Sclerophyllaceae, Duckeodendraceae, Goetzeaceae, Convolvulaceae, Cuscutaceae, Polemoniaceae, and Hydrophyllaceae according to the Systema Naturae 2000, Brands, S.J., Amsterdam or has its origin thereof.
43. The vector, host cell, plant cell, plant tissue, plant, use, kit or method of any one of claims 1 to 42, wherein the polynucleotide, the polypeptide, the plant cell, the host cell, the plant tissue or the plant is derived from the Solanaceae family,

Agrico B.V.

20030596

PF 54801

105

preferably *S. bulbocastanum*, potato (*S. tuberosum*), tomato (*S. lycopersicum* or *Lycopersicon lycopersicum* (L.) Karsten ex Farwell), petunia, tree tomato (*S. betaceum*), pear melon (*S. muricatum*) or eggplant (*S. melongena*).

Agrico B.V.**20030596****PF 54801****106****Resistant plants and uses thereof**

5 The present invention relates to a novel method for increasing the resistance of a plant, in particular of a Solanaceae, preferably of potato and tomato, to plant pathogens of the phylum Oomycetes comprising increasing the activity of the polypeptide of the present invention. The invention further relates to polynucleotides and vectors comprising these polynucleotides. The invention furthermore relates to corresponding vectors, cells, transgenic plants and transgenic propagation material derived from them, methods to produce them and to their use for the production of foodstuffs, feeding stuffs, 10 seed, pharmaceuticals or fine chemicals.

Agrico B.V.

20030596

PF 54801




Figure 1A




Figure 1B




Figure 1C





Figure 1D

Agrico B.V.


20030596



Agrico B.V.

Figure 3

* ARF 87-507 and ARF 87-601 had identical disease progress curves



11-AUG-2003 19:36

BASF AG GUX C100

Agrico B.V.

20030596

+49 621 6021183
PF 54801


S.113/221



Figure 4


Agrico B.V.

20030596



ARG 95-15

Figure 5



ARG 95-3

Figure 6




Figure 7

Agrico B.V.

20030596

PF 54801




Figure 8

Agrico B.V.

20030596

PF 54801




Figure 9

Agrico B.V.

20030596

PF 54801





Figure 10

Agrico B.V.

20030596

PF 54801



B6

Figure 11




Figure 12

Agrico B.V.

20030596

PF 54801

1

Figure 13A

ATGGAAAAACGAAAAGATAATGAAGAAGCAAACAACCTCATTGGAGTCATT 50
TTCTGCTCTTCGCAAGGATGCTGCCAATGTTCTGGATTTCCTAGAGAGAT 100
TAAAGAATGAAGAAGATCAAAAGGCTGTTGATGTGGATCTGATTGAAAGC 150
CTGAAATTGAAGCTGACATTTATTTGTACATATGTCCAGCTTTCTTATTC 200
CGATTTGGAGAAGTTTGAAGATATAATGACTAGAAAAAGACAAGAGGTTG 250
AGAATCTGCTTCAACCAATTTTGGATGATGATGGCAAAGACGTCGGGTGT 300
AAATATGTCCTTACTAGCCTCGCCGGTAATATGGATGACTGTATAAGCTT 350
GTATCATCGTTCTAAATCAGATGCCACCATGATGGATGAGCAATTGGGCT 400
TCCTCCTCTTGAATCTCTCTCATCTATCCAAGCATCGTGCTGAAAAGATG 450
TTTCCTGGAGTGACTCAATATGAGGTTCTTCAGAATGTATGTGGCAACAT 500
AAGAGATTTCCATGGATTGATAGTGAATTGTTGCATTAAGCATGAGATGG 550
TTGAGAATGTCTTATCTCTGTTTCAACTGATGGCTGAGAGAGTAGGACGC 600
TTCCTTTGGGAGGATCAGGCTGATGAAGACTCTCAACTCTCCGAGCTAGA 650
TGAGGATGATCAGAATGATAAAGACCCTCAACTCTTCAAGCTAGCACATC 700
TACTCTTGAAGATTGTTCCAAGTGAATTGGAGGTTATGCACATATGTTAT 750
AAAACCTTTGAAAGCTTCAACTTCAACAGAAATTGGACGCTTCATTAAGAA 800
GCTCCTGGAAACCTCTCCGGACATTCTCAGAGAATATCTGATTCATCTAC 850
AAGAGCATATGATAACTGTTATTACCCCTAACACTTCAGGGGCTCGAAAC 900
ATTCATGTCATGATGGAATTCCTATTGATTATTCTTTCTGATATGCCGCC 950
CAAGGACTTTATTTCATCATGACAACTTTTGTGATCTCTTGGCTCGTGTTG 100
TAGCACTTACCAGGGAGGTATCAACTCTTGTACGCGACTTGGAAGAGAAA 1050
TTAAGGATTAAAGAGAGTACTGACGAAACAAATTGTGCAACCCTAAAGTT 1100
TCTGGAAAATATTGAACTCCTTAAGGAAGATCTCAAACATGTTTATCTGA 1150
AAGTCCCGGATTCATCTCAATATTGCTTCCCCATGAGTGATGGACCTCTC 1200
TTCATGCATCTGCTACAGAGACACTTAGATGATTTGCTGGATTCCAATGC 1250
TTATTCAATTGCTTTGATAAAGGAACAAATTGGGCTGGTGAAAGAAGACT 1300
TGGAATTCATAAGATCTTTTTTTCGCGAATATTGAGCAAGGATTGTATAAA 1350
GATCTCTGGGAACGTGTTCTAGATGTGGCATATGAGGCAAAGATGTCAT 1400
AGATTCAATTATTGTTTCGAGATAATGGTCTCTTACATCTTATTTTCTCAC 1450
TTCCCATTACCAGAAAGAAGATGATGCTTATCAAAGAAGAGGTCTCTGAT 1500
TTACATGAGAACATTTCCAAGAACAGAGGTCTCATCGTTGTGAACTCTCC 1550
CAAGAAACCAGTTGAGAGCAAGTCATTGACAACCTGATAAAATAATTGTAG 1600
GTTTTGGTGAGGAGACAAACTTGATACTTAGAAAGCTCACCAGTGGACCG 1650

Agrico B.V.

20030596

PF 54801

2

GCAGATCTAGATGTCATTTTCGATCATTTGGTATGCCGGGTTTAGGTAAAC 1700
TACTTTGGCGTACAAAGTATACAATGATAAATCAGTTTCTAGCCATTTTCG 1750
ACCTTCGTGCATGGTGCACGGTCCACCAAGTATATGACGAGAAGAAGTTG 1800
TTGGATAAAATTTTCAATCAAGTTAGTGACTCAAATTCAAAATTGAGTGA 1850
GAATATTGATGTTGCTGATAAACTACGGAAACAATTGTTTGGAAAGAGGT 1900
ATCTTATTGTCTTAGATGACGTGTGGGATACTAATACATGGGATGAGCTA 1950
ACAAGACCTTTTTCCTGATGGTATGAAAGGAAGTAGAATTATTTTGACAAC 2000
TCGAGAAAAGAAAGTTGCTTTGTCATGGAAAGCTCTACACTGATCCTCTTA 2050
ACCTTCGATTGCTAAGATCAGAAGAAAGTTGGGAGTTATTAGAGAAAAGG 2100
GCATTTGGAAACGAGAGTTGCCCTGATGAACTATTGGATGTTGGTAAAGA 2150
AATAGCCGAAAATTGTAAAGGGCTTCCTTTGGTGGTGGATCTGATTGCTG 2200
GAATCATTGCTGGGAGGGAAAAGAAAAAGAGTGTGTGGCTTGAAGTTGTA 2250
AATAATTTGCATTCCCTTTATTTTGAAGAATGAAGTGGAAGTGATGAAAGT 2300
TATAGAAATAAGTTATGACCACCTTACCTGATCACCTGAAGCCATGCTTGC 2350
TGTACTTTGCAAGTGCGCCGAAGGACTGGGTAACGACAATCCATGAGTTG 2400
AACTTATTTGGGGTTTGAAGGATTTGTGGAAAAGACAGATATGAAGAC 2450
TCTGGAAGAAGTGGTGAAAATTTATTTGGATGATTTAATTTCCAGTAGCT 2500
TGGTAATTTGTTTCAATGAGATAGGTGATTACCTTACTTGCCAACTTCAT 2550
GATCTTGTGCATGACTTTTGTTTGATAAAAGCAAGAAAGGAAAAGTTGTG 2600
TGATCGGATAAGTTCAAGTGCTCCATCAGATTTGTTGCCACGTCAAATTA 2650
GCATTGATTATGATGATGATGAAGAGCACTTTGGGCTTAATTTTGTCTTG 2700
TTCGGTTCAAATAAGAAAAGGCATTCCGGTAAACACCTCTATTCTTTGAC 2750
CATAAATGGAGATGAGCTGGACGACCATCTTTCTGATACATTTTCATCTAA 2800
GACACTTGAGGCTTCTTAGAACCTTGCACCTGGAATCCTCTTTTATCATG 2850
GTTAAAGATTCTTTGCTGAATGAAATATGCATGTTGAATCATTTGAGGTA 2900
CTTAAGCATTTGGGACAGAAGTTAAATCTCTGCCTTTGTCTTTCTCAAACC 2950
TCTGGAATCTAGAAATCTTGTTTGTGGATAACAAAGAATCAACCTTGATA 3000
CTATTACCGAGAATTTGGGATCTTGTAAGTTGCAAGTGCTGTTACAGAC 3050
TGCTTGTTCTTTCTTTGATATGGATGCAGATGAATCAATACTGATAGCAG 3100
AGGACACAAAGTTAGAGAACTTGACAGCATTAGCGGAACTCGTGCTTTCC 3150
TATTGGAAAGATACAGAGGATATTTTCAAAGGCTTCCCAATCTTCAAGT 3200
GCTTCATTTCAAACCTCAAGGAGTCATGGGATTATTCAACAGAGCAATATT 3250
GGTTCCCGAAATTGGATTTCCTAACTGAAGTACAAAACCTCACTGTAGAT 3300
TTTGAAAGATCAAACACAAATGACAGTGGGTCTCTGCAGCCATAAATCG 3350
GCCATGGGATTTTCACTTTCTTCGAGTTTGAAAAGATTGCAATTGCATG 3400

Agrico B.V.

20030596

PF 54801

3

AATTTCCCTCTGACATCCGATTCACCTATCAACAATAGCGAGACTGCTGAAC 3450
CTTGAAGAGTTGTACCTTTATCGTACAATCATCCATGGGGAAGAATGGAA 3500
CATGGGAGAAGAAGACACCTTTGAGAATCTCAAATGTTTGATGTTGAGTC 3550
AAGTGATTCTTTCCAAGTGGGAGGTTGGAGAGGAATCTTTTCCCACGCTT 3600
GAGAAATTAGAACTGTCGGACTGTCATAATCTTGAGGAGATTCCGTCTAG 3650
TTTTGGGGATATTTATTCCTTGAAAATTATCGAACTTGTAAGGAGCCCTC 3700
AACTTGAAAATTCCGCTCTCAAGATTAAGGAATATGCTGAAGATATGAGG 3750
GGAGGGGACGAGCTTCAGATCCTTGGCCAGAAGGATATCCCGTTATTTAA 3800
GTAG 3804

Agrico B.V.

20030596

PF 54801

4

Figure 13B

ATGGAAAACGAAAAGATAATGAAGAAGCAAACAACTCATTGGTATGTTA 50
TTTGATAGAGTGAACTGTAAAGTATTGAATTGTAGATATCATGTGGCTTT 100
AAAAATTTGATATGTGTTATTTTGGCAGGAGTCATTTTCTGCTCTTCGCA 150
AGGATGCTGCCAATGTTCTGGATTTCTAGAGAGATTAAAGAATGAAGAA 2001
GATCAAAAGGCTGTTGATGTGGATCTGATTGAAAGCCTGAAATTGAAGCT 2501
GACATTTATTTGTACATATGTCAGCTTCTTATTCCGATTTGGAGAAGT 3001
TTGAAGATATAATGACTAGAAAAAGACAAGAGGTTGAGAATCTGCTTCAA 3501
CCAATTTTGGATGATGATGGCAAAGACGTCGGGTGTAAATATGTCCTTAC 4001
TAGCCTCGCCGGTAATATGGATGACTGTATAAGCTTGTATCATCGTTCTA 4501
AATCAGATGCCACCATGATGGATGAGCAATTGGGCTTCCTCCTCTTGAAT 5001
CTCTCTCATCTATCCAAGCATCGTGCTGAAAAGATGTTTCTGGAGTGAC 5501
TCAATATGAGGTTCTTCAGAAATGTATGTGGCAACATAAGAGATTTCCATG 6001
GATTGATAGTGAATTGTTGCATTAAGCATGAGATGGTTGAGAATGTCTTA 6501
TCTCTGTTTCAACTGATGGCTGAGAGAGTAGGACGCTTCCTTTGGGAGGA 7001
TCAGGCTGATGAAGACTCTCAACTCTCCGAGCTAGATGAGGATGATCAGA 7501
ATGATAAAGACCCTCAACTCTTCAAGCTAGCACATCTACTCTTGAAGATT 8001
GTTCCAACCTGAATTGGAGGTTATGCACATATGTTATAAACTTTGAAAGC 8501
TTCAACTTCAACAGAAATTGGACGCTTCATTAAGAAGCTCCTGGAAACCT 9010
CTCCGGACATTCTCAGAGAATATCTGATTCATCTACAAGAGCATATGATA 9510
ACTGTTATTACCCCTAACACTTCAGGGGCTCGAAACATTCATGTCATGAT 1000
GGAATTCCTATTGATTATTCTTTCTGATATGCCGCCCAAGGACTTTATTTC 1050
ATCATGACAAACTTTTGTATCTCTTGGCTCGTGTTGTAGCACTTACCAGG 1100
GAGGTATCAACTCTTGACGCCGACTTGGAAAGAGAAATTAAGGATTAAAGA 1150
GAGTACTGACGAAACAAATTGTGCAACCCCTAAAGTTTCTGGAAAATATTG 1200
AACTCCTTAAGGAAGATCTCAAACATGTTTATCTGAAAGTCCCGGATTCA 1250
TCTCAATATTGCTTCCCCATGAGTGATGGACCTCTCTTCATGCATCTGCT 1300
ACAGAGACACTTAGATGATTTGCTGGATTCCAATGCTTATTCAATTGCTT 1350
TGATAAAGGAACAAATTGGGCTGGTGAAAAGAGACTTGGAAATTCATAAGA 1400
TCTTTTTTCGCGAATATTGAGCAAGGATTGTATAAAGATCTCTGGGAACG 1450
TGTTCTAGATGTGGCATATGAGGCAAAAGATGTCATAGATTCAATTATTG 1500
TTCGAGATAATGGTCTCTTACATCTTATTTTCTCACTTCCCATTACCAGA 1550
AAGAAGATGATGCTTATCAAGAAGAGGTCTCTGATTTACATGAGAACAT 1600
TTCCAAGAACAGAGGTCTCATCGTTGTGAACTCTCCAAGAAACCAGTTG 1650

Agrico B.V.

20030596

PF 54801

5

AGAGCAAGTCATTGACAACCTGATAAAATAATTGTAGGTTTTGGTGAGGAG 1700
ACAAACTTGATACTTAGAAAGCTCACCAGTGGACCGGCAGATCTAGATGT 1750
CATTTTCGATCATTGGTATGCCGGGTTTAGGTAAAACTACTTTGGCGTACA 1800
AAGTATACAATGATAAATCAGTTTCTAGCCATTTTCGACCTTCGTGCATCG 1850
TGCACGGTCCGACCAAGTATATGACGAGAAGAAGTTGTTGGATAAAATTTT 1900
CAATCAAGTTAGTGACTCAAATTCAAAATTGAGTGAGAATATTGATGTTG 1950
CTGATAAACTACGGAAACAATTGTTTGGAAAGAGGTATCTTATTGTCTTA 2000
GATGACGTGTGGGATACTAATACATGGGATGAGCTAACAAGACCTTTTCC 2050
TGATGGTATGAAAGGAAGTAGAATTATTTTGACAACTCGAGAAAAGAAAG 2100
TTGCTTTGCATGGAAAGCTCTACACTGATCCTCTTAACCTTCGATTGCTA 2150
AGATCAGAAGAAAGTTGGGAGTTATTAGAGAAAAGGGCATTTGGAAACGA 2200
GAGTTGCCCTGATGAACTATTGGATGTTGGTAAAGAAATAGCCGAAAATT 2250
GTAAAGGGCTTCCTTTGGTGCTGGATCTGATTGCTGGAATCATTGCTGGG 2300
AGGGAAAAGAAAAGAGTGTGTGGCTTGAAGTTGTAAATAATTTGCATTC 2350
CTTTATTTTGAAGAATGAAGTGGAAGTGATGAAAGTTATAGAAATAAGTT 2400
ATGACCACTTACCTGATCACCTGAAGCCATGCTTGCTGTACTTTGCAAGT 2450
GCGCCGAAGGACTGGGTAAACGACAATCCATGAGTTGAAACTTATTTGGGG 2500
TTTTGAAGGATTTGTGGAAAAGACAGATATGAAGAGTCTGGAAGAAGTGG 2550
TGAAAATTTATTTGGATGATTTAATTTCCAGTAGCTTGGTAATTTGTTTC 2600
AATGAGATAGGTGATTACCTTACTTGCCAACCTTCATGATCTTGTGCATGA 2650
CTTTTGTGTTGATAAAAGCAAGAAAGGAAAAGTTGTGTGATCGGATAAGTT 2700
CAAGTGCTCCATCAGATTTGTTGCCACGTCAAATTAGCATTGATTATGAT 2750
GATGATGAAGAGCACCTTTGGGCTTAATTTTGTCTGTTTCGGTTCAAATAA 2800
GAAAAGGCATTCCGGTAAACACCTCTATTCTTTGACCATAAATGGAGATG 2850
AGCTGGACGACCATCTTCTGATACATTTTCATCTAAGACACTTGAGGCTT 2900
CTTAGAACCTTGCACCTGGAATCCTCTTTTATCATGGTTAAAGATTCTTT 2950
GCTGAATGAAATATGCATGTTGAATCATTTGAGGTACTTAAGCATTGGGA 3000
CAGAAGTTAAATCTCTGCCCTTTGTCTTTCTCAAACCTCTGGAATCTAGAA 3050
ATCTTGTTTGTGGATAACAAAGAATCAACCTTGATACTATTACCGAGAAT 3100
TTGGGATCTTGTAAGTTGCAAGTGCTGTTTCAGACTGCTTGTTCTTTCT 3150
TTGATATGGATGCAGATGAATCAATACTGATAGCAGAGGACACAAAGTTA 3200
GAGAACTTGACAGCATTAGGGGAACCTCGTGCTTTTCTTATTGGAAAGATAC 3250
AGAGGATATTTTCAAAGGCTTCCCAATCTTCAAGTGCTTCATTTCAAAC 3300
TCAAGGAGTCATGGGATTATTCAACAGAGCAATATTGGTTCCCGAAATTG 3350
GATTTCTAACTGAACTAGAAAACTCACTGTAGATTTTGAAAGATCAAA 3400

Agrico B.V.

20030596

PF 54801

.6

CACAAATGACAGTGGGTCCTCTGCAGCCATAAATCGGCCATGGGATTTTC 3450
ACTTTCCTTCGAGTTTGAAAAGATTGCAATTGCATGAATTTCTCTGACA 3500
TCCGATTCACATCAACAATAGCGAGACTGCTGAACCTTGAAGAGTTGTA 3550
CCTTTATCGTACAATCATCCATGGGGAAGAATCGAACATGGGAGAAGAAG 3600
ACACCTTTGAGAATCTCAAATGTTTGATGTTGAGTCAAGTGATTCTTTCC 3650
AAGTGGGAGGTTGGAGAGGAATCTTTTCCCACGCTTGAGAAATTAGAACT 3700
GTCGGACTGTCATAATCTTGAGGAGATTCCGTCTAGTTTTGGGGATATTT 3750
ATTCCTTGAAAATTATCGAACTTGTAAGGAGCCCTCAACTTGAAAATTCC 3800
GCTCTCAAGATTAAGGAATATGCTGAAGATATGAGGGGAGGGGACGAGCT 3850
TCAGATCCTTGGCCAGAAGGATATCCCGTTATTTAAGTAG 3890

Agrico B.V.

20030596

PF 54801

7

Figure 13C

GATCTAGAATCACCGAACCTCCCCCTCGGTACAGCTCCTCCAGTTCTACCA 50
TGAATTTTCATCCACTGATTCTCTTCAATCGCCATTGCAGATTCTCTCGA 100
TCTATGCTCAAAAAATCCCGAGATAAAACCCTAGATCTGCTTCAAATGCT 150
CTGATACCATGTAATTTTCAGTGAATTCTAACTAAACAATGGAGAGAATTA 200
ACTATTTTGTAGAAAGACTGATTGAAGGAGAAGAAGAGAGAAAAATTCTATA 250
TTGAACTCATGAACCAAAATGAATGAAAAAATAATGAGAAGAACTATAC 300
TATTACAATCTATATATCTCTATTTATATTTCTAATCTGAAGCAGTTAATT 350
TAACTGACTCTAACAACCTAGACTGATAGGTGTACATTTTCTGTTAGTGCA 400
CTGCAGTGCATTTAACTAACTGCTTAACATAAAGAATGTTGTTTGAAGT 450
CATTCGAATAGCTTCAATGAGAAGCAAACATGTGTACCTGTAAAGACACA 500
CAGTAAAAGTGTTAATAATGAATAAATATGAATAAATCAAATAATAAATT 550
AAAAATAAAAAACACATCCAATTAACATTGGAGGTCTTGAAAATCGATGGT 600
AATTAACAAAGACCCCTTGTGAAATTTAAGTCTGTAATTGAAAATTTGAGT 650
ATAGGTTAGGGGACATTTGACTATTTTCTCATTTTCTTTATCTTTTTCCT 700
AATTTGTGGCAGACAAGTGAGGAGGCCCACTGTAATTGATTCATGCTTT 750
TGCTTTCTTGACTTTTTTGAACAATACTATGCATCATATTTGGTCTTAAT 800
TATTCCTCTGTTTATTTCCAGAATTTTGAGCTCTATACATCTAATAACAA 850
AGCAAGCAGAGGATATATAGTTTCATCAACTAAAAAGGTTAGTCAACTCA 900
TCTAATATTTGCTACTCTCATCTCTATTTGAAGTACAGTTATGGAAAAGTA 950
GAAGTGATGTAAGAAAAATGAAAGAACTTTTAGTAGGTTAGTTGGATCTAA 1000
CAAAGAGAAAGGGAAATAAATTTGCAGGAGAAAGAGAGAGGTTAAATACTT 1050
ACTCACACCACCGATTTACAACAAATCACTTAATTGTGGTTAGTTAATGT 1100
ATACTTTCACCTCATTAATTTATTACTTACCCATGATAAGTTGTATTAAT 1150
TTGGTATTAATATCCGGTGCGGGTGAATTTCTTACCGGGTGAGAGGGATGG 1200
GCTTGGAGAGTGTGGAGTGAACAGAAGCAGATGTTTTAGATTTTTTCTAA 1250
GATGACGAAAGATTTCCCTCACTAATGAAAATATATTACTATACGCTATT 1300
AGAGATAGAAAGGTTCCGTACCAGTTGGTCTCGTTTCTGGATGAACCCCA 1350
TTTTTACAAGTCATTTTCTTCAATTCAAATCGCAAGTGACCTTTATCAT 1400
CTTCCACTAATTAAGTCCTCTTAAGTTGCGGTGAAAATAGTGAAATTATT 1450
GATTATTCTTATCATTTTCATCTTCTTTCTCCTGATAAAGTTTTATGTACT 1500
TTTTATGCATCAGGTCTTGAGAAGTTGGAAGGAAAAGTAGAATCATGGA 1550
AAAACGAAAAGATAATGAAGAAGCAAACTCAATTGGTATGTTATTTGA 1600
TAGAGTGAAGTGTAAAGTATTGAATTGTAGATATCATGTGGCTTTAAAAA 1650

Agrico B.V.

20030596

PF 54801

8

TTTGATATGTGTTATTTTGGCAGGAGTCATTTTCTGCTCTTCGCAAGGAT 17001
GCTGCCAATGTTCTGGATTTCCTAGAGAGATTAAAGAATGAAGAAGATCA 1750
AAAGGCTGTTGATGTGGATCTGATTGAAAGCCTGAAATTGAAGCTGACAT 1800
TTATTTGTACATATGTCCAGCTTTCTTATTCCGATTTGGAGAAGTTTGAA 1850
GATATAATGACTAGAAAAAGACAAGAGGTTGAGAATCTGCTTCAACCAAT 1900
TTTGGATGATGATGGCAAAGACGTCGGGTGTAAATATGTCCTTACTAGCC 1950
TCGCCGGTAATATGGATGACTGTATAAGCTTGTATCATCGTTCTAAATCA 2000
GATGCCACCATGATGGATGAGCAATTGGGCTTCCTCCTCTTGAATCTCTC 2050
TCATCTATCCAAGCATCGTGCTGAAAAGATGTTTCCTGGAGTGAATCAAT 2100
ATGAGGTTCTTCAGAATGTATGTGGCAACATAAGAGATTTCCATGGATTG 2150
ATAGTGAATTGTTGCATTAAGCATGAGATGGTTGAGAATGTCTTATCTCT 2200
GTTTCAACTGATGGCTGAGAGAGTAGGACGCTTCCTTTGGGAGGATCAGG 2250
CTGATGAAGACTCTCAACTCTCCGAGCTAGATGAGGATGATCAGAATGAT 2300
AAAGACCCTCAACTCTTCAAGCTAGCACATCTACTCTTGAAGATTGTTCC 2350
AACTGAATTGGAGGTTATGCACATATGTTATAAACTTTGAAAGCTTCAA 2400
CTTCAACAGAAATTGGACGCTTCATTAAGAAGCTCCTGGAAACCTCTCCG 2450
GACATTCTCAGAGAATATCTGATTCTATCAAGAGCATATGATAACTGT 2500
TATTACCCCTAACACTTCAGGGCTCGAAACATTCATGTCATGATGGAAT 2550
TCCTATTGATTATTCTTTCTGATATGCCGCCCAAGGACTTTATTCATCAT 2600
GACAACTTTTTTGATCTCTTGGCTCGTGTGTAGCACTTACCAGGGAGGT 2650
ATCAACTCTTGTACGCGACTTGGGAAGAGAAATTAAGGATTAAAGAGAGTA 2700
CTGACGAAACAAATTGTGCAACCCCTAAAGTTTCTGGAAAATATTGAACTC 2750
CTTAAGGAAGATCTCAAACATGTTTATCTGAAAGTCCCGGATTCTCTCA 2800
ATATTGCTTCCCCATGAGTGATGGACCTCTCTTCATGCATCTGCTACAGA 2850
GACACTTAGATGATTTGCTGGATTCCAATGCTTATTCAATTGCTTTGATA 2900
AAGGAACAAATTGGGCTGGTGAAAGAAGACTTGAATTCATAAGATCTTT 2950
TTTCGCGAATATTGAGCAAGGATTGTATAAAGATCTCTGGGAACGTGTTT 3000
TAGATGTGGCATATGAGGCAAAGATGTCATAGATTCAATTATTGTTTCGA 3050
GATAATGGTCTCTTACATCTTATTTTCTCACCTCCCATTACCAGAAAGAA 3100
GATGATGCTTATCAAAGAAGAGGTCTCTGATTTACATGAGAACATTTCCA 3150
AGAACAGAGGTCTCATCGTTGTGAACCTCTCCAAGAAACCAGTTGAGAGC 3200
AAGTCATTGACAACCTGATAAAATAATTGTAGGTTTTGGTGAGGAGACAAA 3250
CTTGATACTTAGAAAGCTCACCAGTGGACCGGAGATCTAGATGTCATTT 3300
CGATCATTGGTATGCCGGGTTTAGGTAAAACTACTTTGGCGTACAAAGTA 3350
TACAATGATAAATCAGTTTCTAGCCATTTTCGACCTTCGTGCATGGTGCAC 3400

Agrico B.V.

20030596

PF 54801

9

GGTCGACCAAGTATATGACGAGAAGAAGTTGTTGGATAAAAATTTTCAATC 3450
AAGTTAGTGACTCAAATTCAAAATTGAGTGAGAATATTGATGTTGCTGAT 3500
AAACTACGGAAACAATTGTTTGGAAAGAGGTATCTTATTGTCTTAGATGA 3550
CGTGTGGGATACTAATACATGGGATGAGCTAACAAGACCTTTTCCTGATG 3600
GTATGAAAGGAAGTAGAATTATTTTGACAACCTCGAGAAAAGAAAGTTGCT 3650
TTGCATGGAAAGCTCTACACTGATCCTCTTAACCTTCGATTGCTAAGATC 3700
AGAAGAAAGTTGGGAGTTATTAGAGAAAAGGGCATTGGGAAACGAGAGTT 3750
GCCCTGATGAACTATTGGATGTTGGTAAAGAAATAGCCGAAAATTGTAAA 3800
GGGCTTCCTTTGGTGGTGGATCTGATTGCTGGAATCATTGCTGGGAGGGA 3850
AAAGAAAAAGAGTGTGTGGCTTGAAGTTGTAAATAATTTGCATTCCTTTA 3900
TTTTGAAGAATGAAGTGGAAGTGATGAAAGTTATAGAAATAAGTTATGAC 3950
CACTTACCTGATCACCTGAAGCCATGCTTGCTGTACTTTGCAAGTGCGCC 4000
GAAGGACTGGGTAACGACAATCCATGAGTTGAAACTTATTTGGGGTTTTG 4050
AAGGATTTGTGGAAAAGACAGATATGAAGAGTCTGGAAGAAGTGGTGAAA 4100
ATTTATTTGGATGATTTAATTTCCAGTAGCTTGGTAATTTGTTTCAATGA 4150
GATAGGTGATTACCTTACTTGCCAACTTCATGATCTTGTGCATGACTTTT 4200
GTTTGATAAAAGCAAGAAAGGAAAAGTTGTGTGATCGGATAAGTTCAAGT 4250
GCTCCATCAGATTTGTTGCCACGTCAAATTAGCATTGATTATGATGATGA 4300
TGAAGAGCACTTTGGGCTTAATTTTGTCTGTTTCGGTTCAAATAAGAAAA 4350
GGCATTCCGGTAAACACCTCTATTCTTTGACCATAAATGGAGATGAGCTG 4400
GACGACCATCTTTCTGATACATTTTCATCTAAGACACTTGAGGCTTCTTAG 4450
AACCTTGCACCTGGAATCCTCTTTTATCATGGTTAAAGATTCTTTGCTGA 4500
ATGAAATATGCATGTTGAATCATTGAGGTACTTAAGCATTGGGACAGAA 4550
GTTAAATCTCTGCCTTTGTCTTTCTCAAACCTCTGGAATCTAGAAATCTT 4600
GTTTGTGGATAACAAAGAATCAACCTTGATACTATTACCGAGAATTTGGG 4650
ATCTTGTAAGTTGCAAGTGCTGTTACGACTGCTTGTTCTTTCTTTGAT 4700
ATGGATGCAGATGAATCAATACTGATAGCAGAGGACACAAAGTTAGAGAA 4750
CTTGACAGCATTAGGGGAACCTCGTGCTTTCTTATTGGAAAGATACAGAGG 4800
ATATTTTCAAAGGCTTCCCAATCTTCAAGTGCTTCATTCAAACCTCAAG 4850
GAGTCATGGGATTATTCAACAGAGCAATATTGGTTCCCGAAATTGGATTT 4900
CCTAACTGAACTAGAAAACTCACTGTAGATTTTGAAAGATCAAACACAA 4950
ATGACAGTGGGTCCCTCTGCAGCCATAAATCGGCCATGGGATTTTCACTTT 5000
CCTTCGAGTTTCAAAGATTGCAATTGCATGAATTTCTCTGACATCCGA 5050
TTCACTATCAACAATAGCGAGACTGCTGAACCTTGAAGAGTTGTACCTTT 5100
ATCGTACAATCATCCATGGGGAAGAATGGAACATGGGAGAAGAAGACACC 5150

Agrico B.V.

20030596

PF 54801

10

TTTGAGAATCTCAAATGTTTGATGTTGAGTCAAGTGATTCTTTCCAAGTG 5200
GGAGGTTGGAGAGGAATCTTTTCCCACGCTTGAGAAATTAGAAGTGTCCG 5250
ACTGTCATAATCTTGAGGAGATTCCGTCTAGTTTTGGGGATATTTATTCC 5300
TTGAAAATTATCGAACTTGTAAGGAGCCCTCAACTTGAAAATTCCGCTCT 5350
CAAGATTAAGGAATATGCTGAAGATATGAGGGGACGGGACGAGCTTCAGA 5400
TCCTTGGCCAGAAGGATATCCCGTTATTTAAGTAGTTTTTGAGCATTATG 5450
GTTGAAAAGTAGATTGCACTTTGCTGGGTAGATTGTATATGGTTAAGAAA 5500
ATTCTGTTACAGTTGTTATGAAACATTTTATTTGACTTTTCTGAGTTTC 5550
TTTTAGAAAACCTCAGAAGTTTTTAACAAAAATTATAGTTTTTATAAATAC 5600
AATGTGGATTTGCCCTTTGGCTGTCCAACCTTGGTCTGAAGTCTCATATGCT 5650
CAGAGCACTATCGTTCAACCTCAATCAAGGTACTGATTTAAAATGACATC 5700
TATACTACTTTATCACAAACCCAACGAACCTTCATCTCAAAAGCTAGGCC 5750
AGGAAGTGAAGAGGTTGTAGAGAGCTTATAAGCACTCATGACTTCCTTTT 5800
CTCGAACATTCAACCAACGTAGGCTGAAATCCCACTCTGAACGAAAATAA 5850
GTGTTTGTTTATCAAATTAACCTCTCGTAGTAGAACACTGAAATACCTTCT 5900
TCTAAACGTTCAACAAATGGGATTTCCAGCACTCAAAGTGAATGAAAGGT 5950
TCACATTAATCTTCAAAAAGAATTACGACAATTCATGACCACAAGTACAT 6000
TGACAGCACCATTCAACAGAAGAACAAGTCAATGCTGCATCTTCATCAA 6050
TAATCCGAGTGTGGAACCTCCTTCCTGACACTGTCTGTATATGTAAAGT 6100
TTCTCAACAGGGCAACTTTCTGGTCTCGTATCTGGATGACCCCTCTCGTC 6150
TATAACTTCAACATTAAGCCCTGGCAACTTCTGGACCAACAGCTTACATG 6200
CTTCAAACTTACTGAACAATTAGACATCCAAAGGGATCGCATTTGTCTCC 6250
AGCTTTGCAGCATTAGCCAACAGAGCCTCATCGCCAAAGGGGCAGTCTCT 6300
AATCTCGAATTTGAAAAAATGTTGTTGTATGACTTTCCTCTGACATCCG 6350
ATGCACTATCAACAATAGCAAGACTGGAGGTTGGAGAGGAATCCTTTATT 6400
ATACAATCATTCAGGGAGAAGAATGGAACATGGGGGAGGAAGACACTTTT 6450
GAGAATCTGAAATGTGTTAGAGCCACAAGCTACAGAAGTATTGAATTTGT 6500
CATGAATATCAACATTCTTCATCCTAGTTAATTCTTTTTCAATTTTTAAT 6550
AGACTCTCATTTTAATCACTAATATTCTTCTATTGTGACTTCTTTTCTG 6600
CAGGTGGCAACTTTAAATTCATAAAGTATAGGATTGATGACAAACTCGAA 6650
AAATATCTTAATGAGGTGAAGTTTGAGCAGTCAGCAGATGGTGGTTCCAA 6700
CTCTAAGTTGACAAGCACATACTATCCCGAGGGCGATTTCAAGCCTGAT 6750
GCATATGGTTAGTGTGGCTAGAGCAGACAGGATGTATTACCTGGATATCT 6800
ACCAAGACGAATCCACAATCAGTTTTATGTCAAGCAATACATGAAGTAAC 6850
TCCCGATAGAACAGTAAAAGCAAGATGTGTAGGTGTATCTCGACTCTAAG 6900

Agrico B.V.

20030596

PF 54801

11

AGATTGTACATTCCTCTTTGAGATTTTTACTGCTAATACAAATTTACACC 6950
TCAGAAGCGAATCTAGAATTTCTAGAGCATGAATGCACCACCTAATGAAAG 7000
GAGAAAAAAGGAAGTATGAAGTGGAATTTGATCCTTGTTTCTAGGTATA 7050
TAAAATTTATCATTCAACTATACTTCATTTAGCAAACAACCTCTCTTGCC 7100
ATTATTTTCTCAAACAAGGGCTTCTAATATTGCTAAACTAAAGACTGTCAA 7150
AAGGTAAGTTCATCTTCAAACCTCTCTTGTTTACTTTATCTAAAGGGGAAC 7200
TATGAAAAACAAGAAACATCAGGAATGTCCCGTAAACAAAGCAGCCTCAT 7250
GCACAAAACATCCAACGTTGGTAGGATTAATGGAGGGATCGCATCCCAGG 7300
AGGATACTGTAGAAAAATTAGTGGCTTCTTTACCGCTCAAACCCATGAT 7350
CTATAGGTTACATGGAGACAACCTTTATGGTTGCTCGTAGGCTCCCGTCAA 7400
TTCTCATAAAACCACAACACCAAAGTTGCATCAGACATCATCTTCATTAC 7450
AAGCTGACAATCTCCACAAGTCTTAGTCAACTTGTAATATGAATATTAGC 7500
CAGGTAGACGTACATATTTACAAAATTGAGTTTCCTATATAATATGGTTT 7550
GAAGGAATGAAACATGATGGGGAGGGTAGATAAAATAATATATGAGGCAT 7600
AAAAATAGGAAAGATATTTGTAGTGAGAGGTTTTGACTTTTTATGCTGCT 7650
TTTGATCTTCAGTTTCTTGATTTCTTTTTCTACTGCTTTCCTCTTCTTTC 7700
TCCTGAGTAAAGTTTTATGTAGGTACTTTTTATACGTCCGATCGTGAGAA 7750
CTTGAAAGAAAGCTCTCTATAGCTATGTTAGGTGCCACATAAAAAAATG 7800
AAATATTACAAAAACCTGATAATAAAATACACTAATCTAAGATATTCAC 7850
TGCAACATACATGCAAAATATATATATATAAATTTTCATGAAAATTATAA 7900
CAAATAATAGATGTGAACATATAACTTTAAAAATAATATTACATCCATAA 7950
AGCTTAAATTCTAGATC 7967

Agrico B.V.

20030596

PF 54801

12

Figure 13D

GATCTGCTTCAAATGCTCTGATACCATGTAATTTTCAGTGAATTCTAACTA 50
AACAAATGGAGAGAATTAACATATTTTAGAAAGACTGATTGAAGGAGAAGAA 100
GAGAGAAAAATCTATATTGAACTCATGAACCAAAATGAATGAAAAAAT 150
AATGAGAAGAACTATACTATTACAATCTATATATCTCTATTTATATTCTA 200
ATCTGAAGCAGTTAATTTAACTGACTCTAACAACCTAGACTGATAGGTGTA 250
CATTTTCTGTAGTGCAGTGCAGTGCATTTAACTAACTGCTTAACATAAA 300
GAATGTTGTTGGAACCTTCATTCGAATAGCTTCAATGAGAAGCAAACATGT 350
GTACCTGTAAAGACACACAGTAAAAGTGTTAATAATGAATAAATATGAAT 400
AAATCAAATAATAAATTAATAATAAACAACATCCAATTAACATTGGAGG 450
TCTTGAAAATCGATGGTAATTAACAAAGACCCTTGTGAAATTTAAGTCTG 500
TAATTGAAAATTTGAGTATAGGTTAGGGGACATTTGACTATTTTCTCATT 550
TTCTTTATCTTTTTTCCTAATTTGTGGCAGACAAGTGAGGAGGCCCCACTG 600
TAATTGATTTCATGCTTTTGCTTTCTTGACTTTTGGGAACAATACTATGCA 650
TCATATTTGGTCTTAATTATTCCTCTGTATTATTTCCAGAATTTTGAGCTC 700
TATACATCTAATAACAAAGCAAGCAGAGGATATATAGTTTCATCAACTAA 750
AAAGGTTAGTCAACTCATCTAATATTTGCTACTCTCATCTCTATTGAAGT 800
ACAGTTATGGAAAAGTAGAAGTGATGTAAGAAAAATGAAAGAAGTTTAGT 850
AGGTTAGTTGGATCTAACAAGAGAAAGCGGAAATAAATTGCAGGAGAAAG 900
AGAGAGGTTAAATACTTACTCACACCACCGATTTACAACAAATCACTTAA 950
TTGTGGTTAGTTAATGTATACTTTACCTCATTAATTTATTACTTACCCA 1000
TGATAAGTTGTATTAATTTGGTATTAATATCCGGTGCGGGTGAATTCTTA 1005
CCGGGTGAGAGGGATGGGGTTGGAGAGTGTGGAGTGAACAGAAGCAGATG 1100
TTTTAGATTTTTTCTAAGATGACGAAAGATTCCCCCTCACTAATGAAAATA 1150
TATTACTATACGCTATTAGAGATAGAAAGGTTCCGGTACCAGTTGGTCTCG 1200
TTTCTGGATGAACCCCATTTTTTACAAGTCATTTTCTTCAATTCAAATCGC 1250
AAGTGACCTTTATCATCTTCCACTAATTAAGTCCTCTTAAGTTTCGCGTG 1300
AAAATAGTGAAATTATTGATTATTCCTTATCATTTTCATCTTCTTCTCCTG 1350
ATAAAGTTTTATGTACTTTTTATGCATCAGGTCTTGAGAACTTGGAAGG 1400
AAAAGTAGAATCATGGAAAAACGAAAAGATAATGAAGAAGCAAACAACCTC 1450
ATTGGTATGTTATTTGATAGAGTGAAGTGTAAAGTATTGAATTGTAGATA 1500
TCATGTGGCTTTAAAAATTTGATATGTGTTATTTTGGCAGGAGTCATTTT 1550
CTGCTCTTCGCAAGGATGCTGCCAATGTTCTGGATTTCCTAGAGAGATTA 1600
AAGAATGAAGAAGATCAAAAGGCTGTTGATGTGGATCTGATTGAAAGCCT 1650

Agrico B.V.

20030596

PF 54801

13

GAAATTGAAGCTGACATTTATTTGTACATATGTCCAGCTTTCTTATTCGG 1700
ATTTGGAGAAGTTTGAAGATATAATGACTAGAAAAAGACAAGAGGTTGAG 1750
AATCTGCTTCAACCAATTTTGGATGATGATGGCAAAGACGTCGGGTGTAA 1800
ATATGTCCTTACTAGCCTCGCCGGTAATATGGATGACTGTATAAGCTTGT 1850
ATCATCGTTCTAAATCAGATGCCACCATGATGGATGAGCAATTGGGCTTC 1900
CTCCTCTTGAATCTCTCTCATCTATCCAAGCATCGTGCTGAAAAGATGTT 1950
TCCTGGAGTGACTCAATATGAGGTTCTTCAGAATGTATGTGGCAACATAA 2000
GAGATTTCCATGGATTGATAGTGAATTGTTGCATTAAGCATGAGATGGTT 2050
GAGAATGTCTTATCTCTGTTTCAACTGATGGCTGAGAGAGTAGGACGCTT 2100
CCTTTGGGAGGATCAGGCTGATGAAGACTCTCAACTCTCCGAGCTAGATG 2150
AGGATGATCAGAATGATAAAGACCCTCAACTCTTCAAGCTAGCACATCTA 2200
CTCTTGAAGATTGTTCCAAGTGAATTGGAGGTTATGCACATATGTTATAA 2250
AACTTTGAAAGCTTCAACTTCAACAGAAATTGGACGCTTCATTAAGAAGC 2300
TCCTGGAAACCTCTCCGGACATTCTCAGAGAATATCTGATTCATCTACAA 2350
GAGCATATGATAACTGTTATTACCCCTAACACTTCAGGGGCTCGAAACAT 2400
TCATGTCAATGATGGAATTCCTATTGATTATTCTTTCTGATATGCCGCCCA 2450
AGGACTTTATTCATCATGACAAACTTTTTGATCTCTTGGCTCGTGTGTGA 2500
GCACTTACCAGGGAGGTATCAACTCTTGTACGCGACTTGGAAGAGAAATT 2550
AAGGATTAAAGAGAGTACTGACGAAACAAATTGTGCAACCTAAAGTTTC 2600
TGGAAAATATTGAACCTCTTAAGGAAGATCTCAAACATGTTTATCTGAAA 2650
GTCCCGGATTCATCTCAATATTGCTTCCCATGAGTGATGGACCTCTCTT 2700
CATGCATCTGCTACAGAGACACTTAGATGATTGCTGGATTCCAATGCTT 2750
ATTCAATTGCTTTGATAAAGGAACAAATTGGGCTGGTGAAAGAAGACTTG 2800
GAATTCATAAGATCTTTTTTCGCGAATATTGAGCAAGGATTGTATAAAGA 2850
TCTCTGGGAACGTGTTCTAGATGTGGCATATGAGGCAAAAGATGTCATAG 2900
ATTCAATTATTGTTTCGAGATAATGGTCTCTTACATCTTATTTTCTCACTT 2950
CCCATTACCAGAAAGAAGATGATGCTTATCAAAGAAGAGGTCTCTGATTT 3000
ACATGAGAACATTTCCAAGAACAGAGGTCTCATCGTTGTGAACCTCTCCCA 3050
AGAAACCAGTTGAGAGCAAGTCATTGACAAGTGAATAAATAATTGTAGGT 3100
TTTGGTGAGGAGACAACTTGATACTTAGAAAGCTCACCAGTGGACCGGC 3150
AGATCTAGATGTCATTTTCGATCATTGGTATGCCGGGTTTAGGTAAACTA 3200
CTTTGGCGTACAAAGTATACAATGATAAATCAGTTTCTAGCCATTTTCGAC 3250
CTTCGTGCATGGTGCACGGTCGACCAAGTATATGACGAGAAGAAGTTGTT 3300
GGATAAAATTTTCAATCAAGTTAGTGAAGTCAAATTCAAAATTGAGTGAGA 3350
ATATTGATGTTGCTGATAAACTACGGAAACAATTGTTTGGAAAGACGTAT 3400

Agrico B.V.

20030596

PF 54801

14

CTTATTGTCTTAGATGACGTGTGGGATACTAATACATGGGATGAGCTAAC 3450
AAGACCTTTTCCTGATGGTATGAAAGGAAGTAGAATTATTTTGACAACTC 3500
GAGAAAAGAAAGTTGCTTTGCGATGGAAAGCTCTACACTGATCCTCTTAAC 3550
CTTCGATTGCTAAGATCAGAAGAAAGTTGGGAGTTATTAGAGAAAAGGGC 3600
ATTTGGAAACGAGAGTTGCCCTGATGAACTATTGGATGTTGGTAAAGAAA 3650
TAGCCGAAAATTGTAAAGGGCTTCCTTTGGTGGTGGATCTGATTGCTGGA 3700
ATCATTGCTGGGAGGGAAAAGAAAAAGAGTGTGTGGCTTGAAGTTGTAAA 3750
TAATTTGCATTCCTTTATTTTGAAGAATGAAGTGAAGTGATGAAAGTTA 3800
TAGAAATAAGTTATGACCACTTACCTGATCACCTGAAGCCATGCTTGCTG 3850
TACTTTGCAAGTGCCTCCGAAGGACTGGGTAACGACAATCCATGAGTTGAA 3900
ACTTATTTGGGGTTTTTGAAGGATTTGTGGAAAAGACAGATATGAAGAGTC 3950
TGGAAGAAGTGGTGAAAATTTATTTGGATGATTTAATTTCCAGTAGCTTG 4000
GTAATTTGTTTCAATGAGATAGGTGATTACCCTACTTGCCAACTTCATGA 4050
TCTTGTGCATGACTTTTGTGTTGATAAAAAGCAAGAAAGGAAAAGTTGTGTG 4100
ATCGGATAAGTTCAAGTGCTCCATCAGATTTGTTGCCACGTCAAATTAGC 4150
ATTGATTATGATGATGATGAAGAGCACTTTGGGCTTAATTTTGTCTTGTT 4200
CGGTTCAAATAAGAAAAGGCATTCCGGTAAACACCTCTATTCTTTGACCA 4250
TAAATGGAGATGAGCTGGACGACCATCTTTCTGATACATTTTCATCTAAGA 4300
CACTTGAGGCTTCTTAGAACCTTGCACCTGGAATCCTCTTTTATCATGGT 4350
TAAAGATTCTTTGCTGAATGAAATATGCATGTTGAATCATTTGAGGTAAT 4400
TAAGCATTGGGACAGAAGTTAAATCTCTGCCTTTGTCTTTCTCAAACCTC 4450
TGGAATCTAGAAATCTTGTTGTGGATAACAAAGAATCAACCTTGATACT 4500
ATTACCGAGAATTTGGGATCTTGTAAGTTGCAAGTGCTGTTACGACTG 4550
CTTGTTCTTTCTTTGATATGGATGCAGATGAATCAATACTGATAGCAGAG 4600
GACACAAAGTTAGAGAACTTGACAGCATTAGGGGAACTCGTGCTTTCCCTA 4650
TTGGAAAGATACAGAGGATATTTTCAAAGGCTTCCCAATCTTCAAGTGC 4700
TTCATTTCAAACCTCAAGGAGTCATGGGATTATTCAACAGAGCAATATTGG 4750
TTCCCGAAATTGGATTTCTTAACCTGAAGTGAAGAACTCACTGTAGATTT 4800
TGAAAGATCAAACACAAATGACAGTGGGTCTCTGCAGCCATAAATCGGC 4850
CATGGGATTTTCACTTTCTTTCGAGTTTGAAAAGATTGCAATTGCATGAA 4900
TTTCTCTGACATCCGATTCATCAACAATAGCGAGACTGCTGAACCT 4950
TGAAGAGTTGTACCTTTATCGTACAATCATCCATGGGGAAGAATCGAACA 5000
TGGGAGAAGAAGACACCTTTGAGAATCTCAAATGTTGATGTTGAGTCAA 5050
GTGATTCTTTCCAAGTGGGAGGTTGGAGAGGAATCTTTTCCACGCTTGA 5100
GAAATTAGAACTGTCCGACTGTCATAATCTTGAGGAGATTCCGTCTAGTT 5150

Agrico B.V.

20030586

PF 54801

15

TTGGGGATATTTATTCCCTTGAAAATTATCGAACTTGTAAGGAGCCCTCAA 5200
CTTGAAAATTCGCTCTCAAGATTAAGGAATATGCTGAAGATATGAGGGG 5250
AGGGGACGAGCTTCAGATCCTTGGCCAGAAGGATATCCCGTTATTTAAGT 5300
AGTTTTTTGAGCATTATGGTTGAAAAGTAGATTGCACTTTGCTGGGTAGAT 5350
TGTATATGGTTAAGAAAATTCTGTTACAGTTGTTATGAAACATTTTTATT 5400
TGACTTTTCTGAGTTTCTTTTAGAAAACCTCAGAAGTTTTTAACAAAATT 5450
ATAGTTTTTATAAATACAATGTGGATTTGCCTTTGGCTGTCCAACCTGGT 5500
CTGAAGTCTCATATGCTCAGAGCACTATCGTTCAACCTCAATCAAGGTAC 5550
TGATTTAAATGACATCTATACTACTTTATCACAAACCCAACGAACCTTC 5600
ATCTCAAAAGCTAGGCCAGGAAGTGAAGAGGTTGTAGAGAGCTTATAAGC 5650
ACTCATGACTTCCTTTTCTCGAACATTCAACCAACGTAGGCTGAAATCCC 5700
ACTCTGAACGAAAATAAGTGTGTTTATCAAATTAACCTCTCGTAGTAGA 5750
ACACTGAAATACCTTCTTCTAAACGTTCAACAAATGGGATTTCCAGCACT 5800
CAAAGTGAATGAAAGGTTACATTAATCTTCAAAAAGAATTACGACAATT 5850
CATGACCACAAGTACATTGACAGCACCATTTCAACAGAAGAACAAGTCAA 5900
TGCTGCATCTTCATCAATAATCCGAGTGTCCAACCTCCTTCCTGACACTG 5950
TCCTGTATATGTAAAGTTTCTCAACAGGGCAACTTTCTGGTCTCGTATCT 6000
GGATGACCCCTCTCGTCTATAACTTCAACATTAAGCCCTGGCAACTTCTG 6050
GACCAACAGCTTACATGCTTCAAAACTTACTGAACAATTAGACATCCAAA 6100
GGGATCGCATTTGTCTCCAGCTTTGCAGCATTAGCCAACAGAGCCTCATCG 6150
CCAAAGGGGCAGTCTCTAATCTCGAATTTGAAAAAATTGTTGTTGTATGA 6200
CTTTCCTCTGACATCCGATGCACCTATCAACAATAGCAAGACTGGAGGTTG 6250
GAGAGGAATCCTTTATTATACAATCATTCAGGGAGAAGAATGGAACATGG 6300
GGGAGGAAGACACTTTTGAGAATCTGAAATGTGTTAGAGCCACAAGCTAC 6350
AGAAGTATTGAATTTGTCATGAATATCAACATTTCTTCATCCTAGTTAATT 6400
CTTTTTCAATTTTTTAATAGACTCTCATTTTAATCACTAATATTCTTCTAT 6450
TTGTGACTTCTTTTCTGCAGGTGGCAACTTTAAATTCATAAAGTATAGGA 6500
TTGATGACAACTCGAAAATATCTTAATGAGGTGAAGTTTGAGCAGTCA 6550
GCAGATGGTGGTTCCAACCTCTAAGTTGACAAGCACATACTATCCCGGAGG 6600
GCGATTTCAAGCCTGATGCATATGGTTAGTGTGGCTAGAGCAGACAGGAT 6650
GTATTACCTGGATATCTACCAAGACGAATCCACAATCAGTTTTATGTCAA 6700
GCAATACATGAAGTAACTCCCGATAGAACAGTAAAGCAAGATGTGTAGG 6750
TGTATCTCGACTCTAAGAGATTGTACATTCCTCTTTGAGATTTTACTGC 6800
TAATACAAATTTACACCTCAGAAGCGAATCTAGAATTTCTAGAGCATGAA 6850
TGCACCCTAATGAAAGGAGAAAAAGGAAGTATGAAGTGGGAATTTGAT 6900

Agrico B.V.

20030596

PF 54801

16

CCTTGTTTCTAGGTATATAAAATTTATCATTCAACTATACTTCATTTAGC 6950
AAACAACCTCTCTTTGCCATTATTTCTCAAACAAGGGCTTCTAATATTGCT 7000
AAACTAAAGACTGTCAAAAGGTAAGTTCATCTTCAAACCTCTCTTGTTTAC 7050
TTTATCTAAAGGGGAACATATGAAAAACAAGAAACATCAGGAATGTCCCGT 7100
AAACAAAGCAGCCTCATGCACAAAACATCCAACGTTGGTAGGATTAATGG 7150
AGGGATCGCATCCCAGGAGGATACTGTAGAAAAATTAGTGGCTTCTTTCA 7200
CCGCTCAAACCCATGATCTATAGGTTACATGGAGACAACTTTATGGTTGC 7250
TCGTAGGCTCCCGTCAATTCTCATAAACACCAACACCAAAGTTGCATCAG 7300
ACATCATCTTCATTACAAAGCTGACAATCTCCACAAGTCTTAGTCAACTT 7350
GTAATATGAATATTAGCCAGGTAGACGTACATATTTACAAAATTGAGTTT 7400
CCTATATAATATGGTTTGAAGGAATGAAACATGATGGGGAGGGTAGATAA 7450
AATAATATATGAGGCATAAAAAATAGGAAAGATATTTGTAGTGAGAGGTTT 7500
TGACTTTTTTATGCTGCTTTTGTATCTTCAGTTTCTTGATTTCTTTTTCTAC 7550
TGCTTTCCTCTTCTTTCTCCTGAGTAAAGTTTTATGTAGGTACTTTTTAT 7600
ACGTCCGATCGTGAGAACTTGAAAGAAAGCTCTCTATAGCTATGTTAGGT 7650
GCCCACATAAAAAAATGAAATATTACAAAAACCCTGATAATAAATACAC 7700
TAATCTAAGATATTTCACTGCAACATACATGCAAAATATATATATATAAAT 7750
TTTCATGAAAATTATAACAAATAATAGATGTGAACATATAACTTTAAAAA 7800
TAATATTACATCCATAAAGCTTAAATTCTAGATCCATCTATGCTTGTATG 7850
ATGCATAGCTCAGAATATCTCCATCAAGTGTTAAACTACATATTTTCAATC 7900
AAATTTATATAGAAAACGATAATTAAGGTGAAAACCTTTTATAAAGATATC 7950
GTGTGGTTGTGTGAGTGAGGTGACAAAATAAGTTGTGTGATTATTCAAAA 8000
AGTTTTAATAACGAAAATCCACATGCTTGAATTAATTGAAGCATTAATGT 8050
TGTAACGAAAATATTACATTTATTGAGTTACTGTGATGTTTTAACTGAT 8100
ATATAAAATAATATTGGTATTTCTCTTCATCTGCCACATAATATGTTTTT 8150
TCATCTTTTTTCAATATACAAAATAGAATTATTATTTTGTTCATCTTTT 8200
TAAGTACAAATTATTCATATGTATATAGTACAAAATAAAATATTTACTGT 8250
GGTAAAGTAAATGGAATAAGAGGTCATATTTGAAATAACAATATACTATA 8300
CTATGTTAAAGTATTTTTTATAGTTAAATTTCTCTAGAGTACTTGATTC 8350
TACATACAAATACTAATTTCTGTAATAAATAAATTAATTTCTTCATT 8400
GTTTCTTTATTATTAAATTAGTTTATAATAACTAACTAAGGTAATAAGA 8450
CCTTAGTTTATGTTAATGTGTGTCTCTGTGATTTCTGTTTATAGTCTAAGGG 8500
TGTAAGTTGTGCTTATCCCAAAAATGAAGGAATATCAAAGATATATTAA 8550
AATTAAATTAAATATTTGGAGGTTATGAATATAAAAAGTATCAGAGTTCT 8600
ACATATAAAGAGTAACAATTGAAATAATTAATTAATATGAGATATGAAG 8650

Agrico B.V.

20030596

PF 54801

17

GCGGACATTTAAAGAAAATAATAAATAAATAAATTAAAGGGTATAAATTT 8700
CATAATACATAATACCAATAAGCCGTAGAATATCTCCGTCATAATGCATA 8750
AACTAATAAATCACAAATGTATAACTCACATACAAATATTTTTTGATAAA 8800
GAATTTGAATGTTGTAATAGAATGGAGAATAACTTGTGTCTTATTCCATT 8850
ATGTAAGACGTATAAATACAAATACAATGAGCTCTAATTAATTAAGGAAA 8900
CTAAATAAGGAAGGAATCAAAAAATATTATGTCATATCCCTACATATCTG 8950
CTAGAGATTCATCATATCCTTACATATCTGTTAAGCTATGTCTACACCT 9000
AAAGGTGTCTACAATCATTTTGTAACTCCTCCCTCAAGTTAGAGCATAG 9050
ATATTATTCATTCCCAACTTGTTACAAAGATAATCAACTCGAGTTCCATT 9100
CAACGCTTTTGTGAACAAATCAACTAGTTGCTCTCCTGTCTTCACTTAGC 9150
TAGTGGATATCAGGTTTTCATGAATCTTCTCACGAATAAAATGACAGTCA 9200
ACCTCAATATGTTTAGTTCTTTCATGAGACACCGGATTCAAGGCAATATG 9250
GAGCGCAACTTGATTATCATACTAGAGTTTGTATGGTATATGATGCTTCA 9300
ACCCTATTTCTGTAAAAGATAATGTATCCACATGATCTCACCCATAGAC 9350
TGTAACATAACTCTGTACTTTGATTCTGCACTAGATCAAGATACAACATT 9400
TTGCTTTTTTACTCCTCCATGATACCAGGTTTCATCCAACAAAGACACAAT 9450
AACTTGATAGTAGATCTTCTATCAATTTTCGATCCAGCCCAATCGACATCT 9500
GCAAAACACTCAATATGAGTATGGTCGTGATTTTGATACTATATTCCAAG 9550
ACTAGGAGTTTCTTCAAGTAACATAGAATATGTTCCAAGCTGCCCAGT 9600
GTTTGACGTAGGTGCAAACATGAACTAGCTAACAACACTTACTGCAAAAG 9650
CAATATCAAGATGAGTCACAATAAGGTAGTTTAACTTTCCAACCTAACCTT 9700
TTGTATCTCTATGGATCATTTAAAGGATCGTCGTCATCTTTCATAAGATG 9750
CATATTGGGAACCATTTGGAGAACTTCAGGGTTTGGCTGCCATCTTTCAAT 9800
TTTCTGCAAGTAGATCGAGAGAATATATTCTCTAAGACAAAAGAATTCCC 9850
TTTTTGTCTTCTATTTACTTCTACTCCCAAATGTATTTCAATTGACCCAA 9900
GTCCTTCGTATGAAACCAAGTATGCAGGAAAGACTTGAGGGAAGAGATC 9949

Agrico B.V.

20030596

PF 54801




Figure 14

Agrico B.V.

20030596

PF 54801

Mil.1		VL	S	I	D	V	---	N	L	K	Q	V	KI	MA	57	
Mil.2	I	VL	S	I	I	---	N	L	K	Q	V	KL	MA	57		
Rpi-blb2	MEKRKDNEECANNSLESFSALRKDAANVLDFLERLKNEDQKAVDVDLIESLKLKLTFTICT													60		
Mil.1		C	F	Q			L	-----	F	TS				109		
Mil.2		Y	F	Q		N	SL	-----	TS					109		
Rpi-blb2	YVQLSYSDELKFPEDIMTRKRQOEVENLLQFILDGDDGQVGVCKYVLTSLAGNMDDCISLYHR													120		
Mil.1		Y	I		D	Y	H	I					G	169		
Mil.2		Y	I		D	Y	H	I					L G	169		
Rpi-blb2	S-KSDATMMDEQLGFLLLNLHLKSHRAEKMFPGVTQYEVLLQNVCGNIRDFHGLIVNCCI													179		
Mil.1		P		D	H	D	T	R		E	R	SR		229		
Mil.2		P			H		T	R		EH	R	SR	Q T	229		
Rpi-blb2	KHEMVENVLSLFLQMAERVGFRFLWEDQADEDSQLSELDEDDQNDKDFQLFKLAHULLKIV													239		
Mil.1		V	I	TN		A	V	L	Q		P	V	S	289		
Mil.2				TN		A	V			I	Q	L	P S L	289		
Rpi-blb2	PTELEVMHICYKTLKASTSTEIGRFIKKLETSPIILREYLIHLQEHMITVTITPNTSGAR													299		
Mil.1		L						D	GV				EP N	GNNQ	348	
Mil.2		L						H	GT				N	GNNQ	348	
Rpi-blb2	NIHVMMEFLLIILSDMPKDFIHDKLPDLLARVVALTREVSTLVRDLEEKLRKESTDE													359		
Mil.1		DL			K		AL	C				HI	N	408		
Mil.2		DL			K		A	N	C			HM	N	408		
Rpi-blb2	TNCATLKFLENIELLKEDLKHVYLKVPDSSQYCFPMSDGFLFMHLLQRHLDDLDSNAYS													419		
Mil.1		E	E	Q	K		VD-A		A					467		
Mil.2		S	E	E	SQ		GDAA		I	A				468		
Rpi-blb2	I ALIKEQIGLVKEDLEFIRSFAN-IEQGLYKDLWERVLDVAYEAKDVIDSIIVRDNGLL													478		
Mil.1		I	IK		I	A	D	P	D		R		T	E	527	
Mil.2		I	IK		I	A	D	P	D		R		I	E	528	
Rpi-blb2	HLIFSLPITRKQMLIKEEVSDELHENISKNRGLIVVNSPKKPVEKSLTDDKIIVGFGE													538		
Mil.1		S		T	S					R			GC	587		
Mil.2				T	S					R			G	D	588	
Rpi-blb2	TNLILRKLTSGPADLDVISIlgmpgigkttlaYKVYNDSVSSHFDLRAWCTVDQVYDEK													598		
Mil.1		NT	S		D					T			ESK	647		
Mil.2		T	S		G	D	N			T	L		EAK	648		
Rpi-blb2	KLLDKIFNQVSDSNSKLSSENIDVADKLRRQLFGKrylivlddvwdTWDELTRPFPDGM													658		
Mil.1		E		N	D		PD							707		
Mil.2		E		N	D		PD		D	T				708		
Rpi-blb2	KGSRIILTTRKKVALHGKLYTDFLNLRLRLRSBESWELLEKRAFGNESCPELDDVGKEI													718		
Mil.1		A		V		R		QSS	S	NS		L	H	767		
Mil.2		A		V		R		QSS	S	NS		L	H	768		
Rpi-blb2	AENCKg1plvvdliagTIAGREKKKS VWLEVVANLHSLFKNEVEVMKVIEISYDHLPDH													778		
Mil.1		F	TSL	Y	NVYF	A		G	E	N	M		Y	827		
Mil.2		H	W	TPL	YLFTVYL	A		E	GI		M			828		
Rpi-blb2	lkpcillyfasAPKDWVTTHLKLINGFEGFVEKTDKMSLEEVVKIYLDLSSSLVICF													838		
Mil.1		YALNF	I				N	F	Q	R		T	C	EE	886	
Mil.2		ILNF	I				N	F	R			T	EE	888		
Rpi-blb2	NEIGDYPTCQIhdiVhCFLIKARKEKLCORISSAPSDDLPRQISIDYDDEEHFGLNE													898		
Mil.1		M	D			R	I	Q	SV	A				LRR	946	
Mil.2		M	D			R	Q	SV	A		I		P	L	N	948
Rpi-blb2	VLFGSNKKRHSCKHLYSLTINGDELDDHLSDTFHLRLRLRLT HESSFIMVKDSLNE													958		

Figure 15

Agrico B.V.

20030596

PF 54801

2

Mil.1		D Q Y	S	STNR	V	L	R	SVD	1006
Mil.2		R R Q Y F	S	S	G I V	L	R	SVG	1008
Rpi-blb2		ICMLNHLRYLSIGTEVKSLPLSFSNLWNLEILFVDNKESTLILLPRIWDLVKLOVLETTA							1018
		4	5			6			
Mil.1			RI T	LI S	KN	F	L S E		1066
Mil.2		K	RI	LI S	MN	F	Q E		1068
Rpi-blb2		CSFFDMDADESILIAEDTKLENLTALGELVLGYWKDTEIDFKRLPNLOVLHFKLKESWDY							1078
		7	8			9			
Mil.1		H SE	T S G KS		V T		N I W R		1126
Mil.2		H C	T C G KS	HC	VVT		N E L YD		1128
Rpi-blb2		STEQYWFPKLDLFLTELEKLTVDLFRSNTNDSGSSAATNRPWDFHFPSSLKRLQLHEEPLT							1138
		10				11			
Mil.1		P S H			F NFN	SI			1186
Mil.2		P N S D Q			F N RLIT				1188
Rpi-blb2		SDSLSTIARLINLEELYLYRTIIGHGEEMNGEEDTFENLKCLMLSOVILSKWEVGEESEFP							1198
		12			13				
Mil.1	N	K R G K	P	S K I K	D				1246
Mil.2	N	K Q E G K	P	F K I K	D	K	ND		1248
Rpi-blb2		TLKLRSLSDCHNLEEIFSSFGDIYSLKIILVRSPOLENSALKIKIKEYAEDMRGGDELQIL							1258
		14	15						
Mil.1	N	1255							
Mil.2	N	1257							
Rpi-blb2		GQKDIPLEK	1267						

Figure 15

Agrico B.V.

20030596

PF 54801

SEQUENCE LISTING

5 <110> Agrico

10 <120> Resistant plants and uses thereof

<130> AE 20030596

15 <160> 94

20 <170> PatentIn version 3.1

25 <210> 1

<211> 3804

<212> DNA

30 <213> Solanum bulbocastanum

35 <220>

<221> CDS

40 <222> (1)..(3804)

<223>

45 <400> 1 48

atg gaa aaa cga aaa gat aat gaa gaa gca aac aac tca ttg gag tca
Met Glu Lys Arg Lys Asp Asn Glu Glu Ala Asn Asn Ser Leu Glu Ser
1 5 10 15

50 ttt tct gct ctt cgc aag gat gct gcc aat gtt ctg gat ttc cta gag 96

Phe Ser Ala Leu Arg Lys Asp Ala Ala Asn Val Leu Asp Phe Leu Glu
20 25 30

55 aga tta aag aat gaa gaa gat caa aag gct gtt gat gtg gat ctg att 144

Arg Leu Lys Asn Glu Glu Asp Gln Lys Ala Val Asp Val Asp Leu Ile
35 40 45

60 gaa agc ctg aaa ttg aag ctg aca ttt att tgt aca tat gtc cag ctt 192

Glu Ser Leu Lys Leu Lys Leu Thr Phe Ile Cys Thr Tyr Val Gln Leu
50 55 60

65 tot tat tcc gat ttg gag aag ttt gaa gat ata atg act aga aaa aga 240

Ser Tyr Ser Asp Leu Glu Lys Phe Glu Asp Ile Met Thr Arg Lys Arg
65 70 75 80

caa gag gtt gag aat ctg ctt caa cca att ttg gat gat gat ggc aaa 288

Gln Glu Val Glu Asn Leu Leu Gln Pro Ile Leu Asp Asp Asp Gly Lys
85 90 95

70 gac gtc ggg tgt aaa tat gtc ctt act agc ctc gcc ggt aat atg gat 336

Asp Val Gly Cys Lys Tyr Val Leu Thr Ser Leu Ala Gly Asn Met Asp
100 105 110

75 gac tgt ata agc ttg tat cat cgt tct aaa tca gat gtc acc atg atg 384

Asp Cys Ile Ser Leu Tyr His Arg Ser Lys Ser Asp Ala Thr Met Met

Agrico B.V.

20030596

PF 54801

2

		gac	tgt	ata	agc	ttg	tat	cat	cgt	tct	aaa	tca	gat	gcc	acc	atg	atg	384
		Asp	Cys	Ile	Ser	Leu	Tyr	His	Arg	Ser	Lys	Ser	Asp	Ala	Thr	Met	Met	
				115					120					125				
5		gat	gag	caa	ttg	ggc	ttc	ctc	ctc	ttg	aat	ctc	tct	cat	cta	tcc	aag	432
		Asp	Glu	Gln	Leu	Gly	Phe	Leu	Leu	Leu	Asn	Leu	Ser	His	Leu	Ser	Lys	
			130					135					140					
10		cat	cgt	gct	gaa	aag	atg	ttt	cct	gga	gtg	act	caa	tat	gag	gtt	ctt	480
		His	Arg	Ala	Glu	Lys	Met	Phe	Pro	Gly	Val	Thr	Gln	Tyr	Glu	Val	Leu	
							150					155					160	
15		cag	aat	gta	tgt	ggc	aac	ata	aga	gat	ttc	cat	gga	ttg	ata	gtg	aat	528
		Gln	Asn	Val	Cys	Gly	Asn	Ile	Arg	Asp	Phe	His	Gly	Leu	Ile	Val	Asn	
						165					170					175		
20		tgt	tgc	att	aag	cat	gag	atg	gtt	gag	aat	gtc	tta	tct	ctg	ttt	caa	576
		Cys	Cys	Ile	Lys	His	Glu	Met	Val	Glu	Asn	Val	Leu	Ser	Leu	Phe	Gln	
					180					185					190			
25		ctg	atg	gct	gag	aga	gta	gga	cgc	ttc	ctt	tgg	gag	gat	cag	gct	gat	624
		Leu	Met	Ala	Glu	Arg	Val	Gly	Arg	Phe	Leu	Trp	Glu	Asp	Gln	Ala	Asp	
				195				200					205					
30		gaa	gac	tct	caa	ctc	tcc	gag	cta	gat	gag	gat	gat	cag	aat	gat	aaa	672
		Glu	Asp	Ser	Gln	Leu	Ser	Glu	Leu	Asp	Glu	Asp	Asp	Gln	Asn	Asp	Lys	
			210					215					220					
35		gac	cct	caa	ctc	ttc	aag	cta	gca	cat	cta	ctc	ttg	aag	att	gtt	cca	720
		Asp	Pro	Gln	Leu	Phe	Lys	Leu	Ala	His	Leu	Leu	Lys	Ile	Val	Pro		
			225				230					235				240		
40		act	gaa	ttg	gag	gtt	atg	cac	ata	tgt	tat	aaa	act	ttg	aaa	gct	tca	768
		Thr	Glu	Leu	Glu	Val	Met	His	Ile	Cys	Tyr	Lys	Thr	Leu	Lys	Ala	Ser	
						245					250					255		
45		act	tca	aca	gaa	att	gga	cgc	ttc	att	aag	aag	ctc	ctg	gaa	acc	tct	816
		Thr	Ser	Thr	Glu	Ile	Gly	Arg	Phe	Ile	Lys	Lys	Leu	Leu	Glu	Thr	Ser	
					260				265						270			
50		ccg	gac	att	ctc	aga	gaa	tat	ctg	att	cat	cta	caa	gag	cat	atg	ata	864
		Pro	Asp	Ile	Leu	Arg	Glu	Tyr	Leu	Ile	His	Leu	Gln	Glu	His	Met	Ile	
				275				280					285					
55		act	gtt	att	acc	cct	aac	act	tca	ggg	gct	cga	aac	att	cat	gtc	atg	912
		Thr	Val	Ile	Thr	Pro	Asn	Thr	Ser	Gly	Ala	Arg	Asn	Ile	His	Val	Met	
			290				295					300						
60		atg	gaa	ttc	cta	ttg	att	att	ctt	tct	gat	atg	cog	coc	aag	gac	ttt	960
		Met	Glu	Phe	Leu	Leu	Ile	Ile	Leu	Ser	Asp	Met	Pro	Pro	Lys	Asp	Phe	
			305				310					315				320		
65		att	cat	cat	gac	aaa	ctt	ttt	gat	ctc	ttg	gct	cgt	gtt	gta	gca	ctt	1008
		Ile	His	His	Asp	Lys	Leu	Phe	Asp	Leu	Leu	Ala	Arg	Val	Val	Ala	Leu	
						325					330				335			
70		acc	agg	gag	gta	tca	act	ctt	gta	cgc	gac	ttg	gaa	gag	aaa	tta	agg	1056
		Thr	Arg	Glu	Val	Ser	Thr	Leu	Val	Arg	Asp	Leu	Glu	Glu	Lys	Leu	Arg	
					340				345					350				
75		att	aaa	gag	agt	act	gac	gaa	aca	aat	tgt	gca	acc	cta	aag	ttt	ctg	1104
		Ile	Lys	Glu	Ser	Thr	Asp	Glu	Thr	Asn	Cys	Ala	Thr	Leu	Lys	Phe	Leu	
				355				360						365				
80		gaa	aat	att	gaa	ctc	ctt	aag	gaa	gat	ctc	aaa	cat	gtt	tat	ctg	aaa	1152
		Glu	Asn	Ile	Glu	Leu	Leu	Lys	Glu	Asp	Leu	Lys	His	Val	Tyr	Leu	Lys	
			370				375					380						
85		gtc	ccg	gat	tca	tct	caa	tat	tgc	ttc	ccc	atg	agt	gat	gga	cct	ctc	1200
		Val	Pro	Asp	Ser	Ser	Gln	Tyr	Cys	Phe	Pro	Met	Ser	Asp	Gly	Pro	Leu	
							390					395					400	
90		ttc	atg	cat	ctg	cta	cag	aga	cac	tta	gat	gat	ttg	ctg	gat	tcc	aat	1248

Agrico B.V.

20030596

PF 54801

3

	Phe	Met	His	Leu	Leu	Gln	Arg	His	Leu	Asp	Asp	Leu	Leu	Asp	Ser	Asn	
				405						410					415		
5	gct	tat	tca	att	gct	ttg	aca	aag	gaa	caa	att	ggg	ctg	gtg	aaa	gaa	1296
	Ala	Tyr	Ser	Ile	Ala	Leu	Ile	Lys	Glu	Gln	Ile	Gly	Leu	Val	Lys	Glu	
				420					425					430			
10	gac	ttg	gaa	ttc	ata	aga	tct	ttt	ttc	gcg	aat	att	gag	caa	gga	ttg	1344
	Asp	Leu	Glu	Phe	Ile	Arg	Ser	Phe	Phe	Ala	Asn	Ile	Glu	Gln	Gly	Leu	
				435				440					445				
15	tat	aaa	gat	ctc	tgg	gaa	cgt	gtt	cta	gat	gtg	gca	tat	gag	gca	aaa	1392
	Tyr	Lys	Asp	Leu	Trp	Glu	Arg	Val	Leu	Asp	Val	Ala	Tyr	Glu	Ala	Lys	
				450			455					460					
20	gat	gtc	ata	gat	tca	att	att	gtt	cga	gat	aat	ggg	ctc	tta	cat	ctt	1440
	Asp	Val	Ile	Asp	Ser	Ile	Ile	Val	Arg	Asp	Asn	Gly	Leu	Leu	His	Leu	
						470					475					480	
25	att	ttc	tca	ctt	ccc	att	acc	aga	aag	aag	atg	atg	ctt	atc	aaa	gaa	1488
	Ile	Phe	Ser	Leu	Pro	Ile	Thr	Arg	Lys	Lys	Met	Met	Leu	Ile	Lys	Glu	
					485					490					495		
30	gag	gtc	tct	gat	tta	cat	gag	aac	att	tcc	aag	aac	aga	ggt	ctc	atc	1536
	Glu	Val	Ser	Asp	Leu	His	Glu	Asn	Ile	Ser	Lys	Asn	Arg	Gly	Leu	Ile	
				500					505					510			
35	gtt	gtg	aac	tct	ccc	aag	aaa	cca	gtt	gag	agc	aag	tca	ttg	aca	act	1584
	Val	Val	Asn	Ser	Pro	Lys	Lys	Pro	Val	Glu	Ser	Lys	Ser	Leu	Thr	Thr	
				515				520					525				
40	gat	aaa	ata	att	gta	ggg	ttt	ggg	gag	gag	aca	aac	ttg	ata	ctt	aga	1632
	Asp	Lys	Ile	Ile	Val	Gly	Phe	Gly	Glu	Glu	Thr	Asn	Leu	Ile	Leu	Arg	
							535					540					
45	aag	ctc	acc	agt	gga	ccg	gca	gat	cta	gat	gtc	att	tcg	atc	att	ggg	1680
	Lys	Leu	Thr	Ser	Gly	Pro	Ala	Asp	Leu	Asp	Val	Ile	Ser	Ile	Ile	Gly	
						550					555					560	
50	atg	ccg	ggg	tta	ggg	aaa	act	act	ttg	gcg	tac	aaa	gta	tac	aat	gat	1728
	Met	Pro	Gly	Leu	Gly	Lys	Thr	Thr	Leu	Ala	Tyr	Lys	Val	Tyr	Asn	Asp	
					565				570						575		
55	aaa	tca	gtt	tct	agc	cat	ttc	gac	ctt	cgt	gca	tgg	tgc	acg	gtc	gac	1776
	Lys	Ser	Val	Ser	Ser	His	Phe	Asp	Leu	Arg	Ala	Trp	Cys	Thr	Val	Asp	
				580					585					590			
60	caa	gta	tat	gac	gag	aag	aag	ttg	ttg	gat	aaa	att	ttc	aat	caa	gtt	1824
	Gln	Val	Tyr	Asp	Glu	Lys	Lys	Leu	Leu	Asp	Lys	Ile	Phe	Asn	Gln	Val	
				595				600					605				
65	agt	gac	tca	aat	tca	aaa	ttg	agt	gag	aat	att	gat	gtt	gct	gat	aaa	1872
	Ser	Asp	Ser	Asn	Ser	Lys	Leu	Ser	Glu	Asn	Ile	Asp	Val	Ala	Asp	Lys	
				610			615					620					
70	cta	cgg	aaa	caa	ttg	ttt	gga	aag	agg	cat	ctt	att	gtc	tta	gat	gac	1920
	Leu	Arg	Lys	Gln	Leu	Phe	Gly	Lys	Arg	Tyr	Leu	Ile	Val	Leu	Asp	Asp	
						630					635					640	
75	gtg	tgg	gat	act	aat	aca	tgg	gat	gag	cta	aca	aga	cct	ttt	cct	gat	1968
	Val	Trp	Asp	Thr	Asn	Thr	Trp	Asp	Glu	Leu	Thr	Arg	Pro	Phe	Pro	Asp	
					645					650					655		
80	ggg	atg	aaa	gga	agt	aga	att	att	ttg	aca	act	cga	gaa	aag	aaa	gtt	2016
	Gly	Met	Lys	Gly	Ser	Arg	Ile	Ile	Leu	Thr	Thr	Arg	Glu	Lys	Lys	Val	
				660					665					670			
85	gct	ttg	cat	gga	aag	ctc	tac	act	gat	cct	ctt	aac	ctt	cga	ttg	cta	2064
	Ala	Leu	His	Gly	Lys	Leu	Tyr	Thr	Asp	Pro	Leu	Asn	Leu	Arg	Leu	Leu	
				675				680					685				
90	aga	tca	gaa	gaa	agt	tgg	gag	tta	tta	gag	aaa	agg	gca	ttt	gga	aac	2112
	Arg	Ser	Glu	Glu	Ser	Trp	Glu	Leu	Leu	Glu	Lys	Arg	Ala	Phe	Gly	Asn	

Agrico B.V.

20030596

PF 54801

4

	690	695	700	
5	gag agt tgc cct gat gaa cta ttg gat gtt ggt aaa gaa ata gcc gaa Glu Ser Cys Pro Asp Glu Leu Leu Asp Val Gly Lys Glu Ile Ala Glu 705 710 715 720			2160
10	aat tgt aaa ggg ctt cct ttg gtg gtg gat ctg att gct gga atc att Asn Cys Lys Gly Leu Pro Leu Val Val Asp Leu Ile Ala Gly Ile Ile 725 730 735			2208
15	gct ggg agg gaa aag aaa aag agt gtg tgg ctt gaa gtt gta aat aat Ala Gly Arg Glu Lys Lys Lys Ser Val Trp Leu Glu Val Val Asn Asn 740 745 750			2256
20	ttg cat tcc ttt att ttg aag aat gaa gtg gaa gtg atg aaa gtt ata Leu His Ser Phe Ile Leu Lys Asn Glu Val Glu Val Met Lys Val Ile 755 760 765			2304
25	gaa ata agt tat gac cac tta cct gat cac ctg aag cca tgc ttg ctg Glu Ile Ser Tyr Asp His Leu Pro Asp His Leu Lys Pro Cys Leu Leu 770 775 780			2352
30	tac ttt gca agt gcg ccg aag gac tgg gta acg aca atc cat gag ttg Tyr Phe Ala Ser Ala Pro Lys Asp Trp Val Thr Thr Ile His Glu Leu 785 790 795 800			2400
35	aaa ctt att tgg ggt ttt gaa gga ttt gtg gaa aag aca gat atg aag Lys Leu Ile Trp Gly Phe Glu Gly Phe Val Glu Lys Thr Asp Met Lys 805 810 815			2448
40	agt ctg gaa gaa gtg gtg aaa att tat ttg gat gat tta att tcc agt Ser Leu Glu Glu Val Val Lys Ile Tyr Leu Asp Asp Leu Ile Ser Ser 820 825 830			2496
45	agc ttg gta att tgt ttc aat gag ata ggt gat tac cct act tgc caa Ser Leu Val Ile Cys Phe Asn Glu Ile Gly Asp Tyr Pro Thr Cys Gln 835 840 845			2544
50	ctt cat gat ctt gtg cat gac ttt tgt ttg ata aaa gca aga aag gaa Leu His Asp Leu Val His Asp Phe Cys Leu Ile Lys Ala Arg Lys Glu 850 855 860			2592
55	aag ttg tgt gat cgg ata agt tca agt gct cca tca gat ttg ttg cca Lys Leu Cys Asp Arg Ile Ser Ser Ser Ala Pro Ser Asp Leu Leu Pro 865 870 875 880			2640
60	cgt caa att agc att gat tat gat gat gat gaa gag cac ttt ggg ctt Arg Gln Ile Ser Ile Asp Tyr Asp Asp Asp Glu Glu His Phe Gly Leu 885 890 895			2688
65	aat ttt gtc ctg ttc ggt tca aat aag aaa agg cat tcc ggt aaa cac Asn Phe Val Leu Phe Gly Ser Asn Lys Lys Arg His Ser Gly Lys His 900 905 910			2736
70	ctc tat tct ttg acc ata aat gga gat gag ctg gac gao cat ctt tct Leu Tyr Ser Leu Thr Ile Asn Gly Asp Glu Leu Asp Asp His Leu Ser 915 920 925			2784
75	gat aca ttt cat cta aga cao ttg agg ctt ctt aga acc ttg cac ctg Asp Thr Phe His Leu Arg His Leu Arg Leu Leu Arg Thr Leu His Leu 930 935 940			2832
80	gaa tcc tct ttt atc atg gtt aaa gat tct ttg ctg aat gaa ata tgc Glu Ser Ser Phe Ile Met Val Lys Asp Ser Leu Leu Asn Glu Ile Cys 945 950 955 960			2880
85	atg ttg aat cat ttg agg tac tta agc att ggg aca gaa gtt aaa tct Met Leu Asn His Leu Arg Tyr Leu Ser Ile Gly Thr Glu Val Lys Ser 965 970 975			2928
90	ctg cct ttg tct ttc tca aac ctc tgg aat cta gaa atc ttg ttt gtg Leu Pro Leu Ser Phe Ser Asn Leu Trp Asn Leu Glu Ile Leu Phe Val 980 985 990			2976

Agrico B.V.

20030596

PF 54801

5

5	gat aac aaa gaa tca acc ttg ata cta tta ccg aga att tgg gat ctt 3024
	Asp Asn Lys Glu Ser Thr Leu Ile Leu Leu Pro Arg Ile Trp Asp Leu 995 1000 1005
10	gta aag ttg caa gtg ctg ttc acg act get tgt tct ttc ttt gat 3069
	Val Lys Leu Gln Val Leu Phe Thr Thr Ala Cys Ser Phe Phe Asp 1010 1015 1020
15	atg gat gca gat gaa tca ata ctg ata gca gag gac aca aag tta 3114
	Met Asp Ala Asp Glu Ser Ile Leu Ile Ala Glu Asp Thr Lys Leu 1025 1030 1035
20	gag aac ttg aca gca tta ggg gaa ctc gtg ctt tcc tat tgg aaa 3159
	Glu Asn Leu Thr Ala Leu Gly Glu Leu Val Leu Ser Tyr Trp Lys 1040 1045 1050
25	gat aca gag gat att ttc aaa agg ctt ccc aat ctt caa gtg ctt 3204
	Asp Thr Glu Asp Ile Phe Lys Arg Leu Pro Asn Leu Gln Val Leu 1055 1060 1065
30	cat ttc aaa ctc aag gag tca tgg gat tat tca aca gag caa tat 3249
	His Phe Lys Leu Lys Glu Ser Trp Asp Tyr Ser Thr Glu Gln Tyr 1070 1075 1080
35	tgg ttc ccg aaa ttg gat ttc cta act gaa cta gaa aaa ctc act 3294
	Trp Phe Pro Lys Leu Asp Phe Leu Thr Glu Leu Glu Lys Leu Thr 1085 1090 1095
40	gta gat ttt gaa aga tca aac acm aat gac agt ggg tcc tct gca 3339
	Val Asp Phe Glu Arg Ser Asn Thr Asn Asp Ser Gly Ser Ser Ala 1100 1105 1110
45	gcc ata aat cgg cca tgg gat ttt cac ttt cct tcg agt ttg aaa 3384
	Ala Ile Asn Arg Pro Trp Asp Phe His Phe Pro Ser Ser Leu Lys 1115 1120 1125
50	aga ttg caa ttg cat gaa ttt cct ctg aca tcc gat tca cta tca 3429
	Arg Leu Gln Leu His Glu Phe Pro Leu Thr Ser Asp Ser Leu Ser 1130 1135 1140
55	aca ata gcg aga ctg ctg aac ctt gaa gag ttg tac ctt tat cgt 3474
	Thr Ile Ala Arg Leu Leu Asn Leu Glu Glu Leu Tyr Leu Tyr Arg 1145 1150 1155
60	aca atc atc cat ggg gaa gaa tgg aac atg gga gaa gaa gac acc 3519
	Thr Ile Ile His Gly Glu Glu Trp Asn Met Gly Glu Glu Asp Thr 1160 1165 1170
65	ttt gag aat ctc aaa tgt ttg atg ttg agt caa gtg att ctt tcc 3564
	Phe Glu Asn Leu Lys Cys Leu Met Leu Ser Gln Val Ile Leu Ser 1175 1180 1185
70	aag tgg gag gtt gga gag gaa tct ttt ccc acg ctt gag aaa tta 3609
	Lys Trp Glu Val Gly Glu Glu Ser Phe Pro Thr Leu Glu Lys Leu 1190 1195 1200
75	gaa ctg tcg gac tgt cat aat ctt gag gag att ccg tct agt ttt 3654
	Glu Leu Ser Asp Cys His Asn Leu Glu Glu Ile Pro Ser Ser Phe 1205 1210 1215
80	ggg gat att tat tcc ttg aaa att atc gaa ctt gta agg agc cct 3699
	Gly Asp Ile Tyr Ser Leu Lys Ile Ile Glu Leu Val Arg Ser Pro 1220 1225 1230
85	caa ctt gaa aat tcc gct ctc aag att aag gaa tat gct gaa gat 3744
	Gln Leu Glu Asn Ser Ala Leu Lys Ile Lys Glu Tyr Ala Glu Asp 1235 1240 1245
90	atg agg gga ggg gac gag ctt cag atc ctt ggc cag aag gat atc 3789
	Met Arg Gly Gly Asp Glu Leu Gln Ile Leu Gly Gln Lys Asp Ile 1250 1255 1260

Agrico B.V.

20030596

PF 54801

6

ccg tta ttt aag tag
Pro Leu Phe Lys
1265

3804

5

<210> 2

<211> 1267

10

<212> PRT

<213> Solanum bulbocastanum

15

<400> 2

20

Met Glu Lys Arg Lys Asp Asn Glu Glu Ala Asn Asn Ser Leu Glu Ser
1 3 10 15

Phe Ser Ala Leu Arg Lys Asp Ala Ala Asn Val Leu Asp Phe Leu Glu
20 25 30

25

Arg Leu Lys Asn Glu Glu Asp Gln Lys Ala Val Asp Val Asp Leu Ile
35 40 45

30

Glu Ser Leu Lys Leu Lys Leu Thr Phe Ile Cys Thr Tyr Val Gln Leu
50 55 60

35

Ser Tyr Ser Asp Leu Glu Lys Phe Glu Asp Ile Met Thr Arg Lys Arg
65 70 75 80

40

Gln Glu Val Glu Asn Leu Leu Gln Pro Ile Leu Asp Asp Asp Gly Lys
85 90 95

Asp Val Gly Cys Lys Tyr Val Leu Thr Ser Leu Ala Gly Asn Met Asp
100 105 110

45

Asp Cys Ile Ser Leu Tyr His Arg Ser Lys Ser Asp Ala Thr Met Met
115 120 125

50

Asp Glu Gln Leu Gly Phe Leu Leu Leu Asn Leu Ser His Leu Ser Lys
130 135 140

55

His Arg Ala Glu Lys Met Phe Pro Gly Val Thr Gln Tyr Glu Val Leu
145 150 155 160

Gln Asn Val Cys Gly Asn Ile Arg Asp Phe His Gly Leu Ile Val Asn
165 170 175

60

Cys Cys Ile Lys His Glu Met Val Glu Asn Val Leu Ser Leu Phe Gln
180 185 190

65

Leu Met Ala Glu Arg Val Gly Arg Phe Leu Trp Glu Asp Gln Ala Asp
195 200 205

70

Glu Asp Ser Gln Leu Ser Glu Leu Asp Glu Asp Asp Gln Asn Asp Lys
210 215 220

Agrico B.V.

20030596

PF 54801

7

	Asp	Pro	Gln	Leu	Phe	Lys	Leu	Ala	His	Leu	Leu	Leu	Lys	Ile	Val	Pro	225	230	235	240
5	Thr	Glu	Leu	Glu	Val	Met	His	Ile	Cys	Tyr	Lys	Thr	Leu	Lys	Ala	Ser	245	250	255	
10	Thr	Ser	Thr	Glu	Ile	Gly	Arg	Phe	Ile	Lys	Lys	Leu	Leu	Glu	Thr	Ser	260	265	270	
15	Pro	Asp	Ile	Leu	Arg	Glu	Tyr	Leu	Ile	His	Leu	Gln	Glu	His	Met	Ile	275	280	285	
	Thr	Val	Ile	Thr	Pro	Asn	Thr	Ser	Gly	Ala	Arg	Asn	Ile	His	Val	Met	290	295	300	
20	Met	Glu	Phe	Leu	Leu	Ile	Ile	Leu	Ser	Asp	Met	Pro	Pro	Lys	Asp	Phe	305	310	315	320
25	Ile	His	His	Asp	Lys	Leu	Phe	Asp	Leu	Leu	Ala	Arg	Val	Val	Ala	Leu	325	330	335	
30	Thr	Arg	Glu	Val	Ser	Thr	Leu	Val	Arg	Asp	Leu	Glu	Glu	Lys	Leu	Arg	340	345	350	
35	Ile	Lys	Glu	Ser	Thr	Asp	Glu	Thr	Asn	Cys	Ala	Thr	Leu	Lys	Phe	Leu	355	360	365	
	Glu	Asn	Ile	Glu	Leu	Leu	Lys	Glu	Asp	Leu	Lys	His	Val	Tyr	Leu	Lys	370	375	380	
40	Val	Pro	Asp	Ser	Ser	Gln	Tyr	Cys	Phe	Pro	Met	Ser	Asp	Gly	Pro	Leu	385	390	395	400
45	Phe	Met	His	Leu	Leu	Gln	Arg	His	Leu	Asp	Asp	Leu	Leu	Asp	Ser	Asn	405	410	415	
50	Ala	Tyr	Ser	Ile	Ala	Leu	Ile	Lys	Glu	Gln	Ile	Gly	Leu	Val	Lys	Glu	420	425	430	
55	Asp	Leu	Glu	Phe	Ile	Arg	Ser	Phe	Phe	Ala	Asn	Ile	Glu	Gln	Gly	Leu	435	440	445	
	Tyr	Lys	Asp	Leu	Trp	Glu	Arg	Val	Leu	Asp	Val	Ala	Tyr	Glu	Ala	Lys	450	455	460	
60	Asp	Val	Ile	Asp	Ser	Ile	Ile	Val	Arg	Asp	Asn	Gly	Leu	Leu	His	Leu	465	470	475	480
65	Ile	Phe	Ser	Leu	Pro	Ile	Thr	Arg	Lys	Lys	Met	Met	Leu	Ile	Lys	Glu	485	490	495	
70	Glu	Val	Ser	Asp	Leu	His	Glu	Asn	Ile	Ser	Lys	Asn	Arg	Gly	Leu	Ile	500	505	510	
	Val	Val	Asn	Ser	Pro	Lys	Lys	Pro	Val	Glu	Ser	Lys	Ser	Leu	Thr	Thr				

Agrico B.V.

20030596

PF 54801

8

515

520

525

5 Asp Lys Ile Ile Val Gly Phe Gly Glu Glu Thr Asn Leu Ile Leu Arg
530 535 540

10 Lys Leu Thr Ser Gly Pro Ala Asp Leu Asp Val Ile Ser Ile Ile Gly
545 550 555 560

Met Pro Gly Leu Gly Lys Thr Thr Leu Ala Tyr Lys Val Tyr Asn Asp
565 570 575

15 Lys Ser Val Ser Ser His Phe Asp Leu Arg Ala Trp Cys Thr Val Asp
580 585 590

20 Gln Val Tyr Asp Glu Lys Lys Leu Leu Asp Lys Ile Phe Asn Gln Val
595 600 605

25 Ser Asp Ser Asn Ser Lys Leu Ser Glu Asn Ile Asp Val Ala Asp Lys
610 615 620

30 Leu Arg Lys Gln Leu Phe Gly Lys Arg Tyr Leu Ile Val Leu Asp Asp
625 630 635 640

Val Trp Asp Thr Asn Thr Trp Asp Glu Leu Thr Arg Pro Phe Pro Asp
645 650 655

35 Gly Met Lys Gly Ser Arg Ile Ile Leu Thr Thr Arg Glu Lys Lys Val
660 665 670

40 Ala Leu His Gly Lys Leu Tyr Thr Asp Pro Leu Asn Leu Arg Leu Leu
675 680 685

45 Arg Ser Glu Glu Ser Trp Glu Leu Leu Glu Lys Arg Ala Phe Gly Asn
690 695 700

50 Glu Ser Cys Pro Asp Glu Leu Leu Asp Val Gly Lys Glu Ile Ala Glu
705 710 715 720

Asn Cys Lys Gly Leu Pro Leu Val Val Asp Leu Ile Ala Gly Ile Ile
725 730 735

55 Ala Gly Arg Glu Lys Lys Lys Ser Val Trp Leu Glu Val Val Asn Asn
740 745 750

60 Leu His Ser Phe Ile Leu Lys Asn Glu Val Glu Val Met Lys Val Ile
755 760 765

65 Glu Ile Ser Tyr Asp His Leu Pro Asp His Leu Lys Pro Cys Leu Leu
770 775 780

Tyr Phe Ala Ser Ala Pro Lys Asp Trp Val Thr Thr Ile His Glu Leu
785 790 795 800

70 Lys Leu Ile Trp Gly Phe Glu Gly Phe Val Glu Lys Thr Asp Met Lys
805 810 815

Agrico B.V.

20030596

PF 54801

9

5 Ser Leu Glu Glu Val Val Lys Ile Tyr Leu Asp Asp Leu Ile Ser Ser
820 825 830

10 Ser Leu Val Ile Cys Phe Asn Glu Ile Gly Asp Tyr Pro Thr Cys Gln
835 840 845

15 Leu His Asp Leu Val His Asp Phe Cys Leu Ile Lys Ala Arg Lys Glu
850 855 860

20 Lys Leu Cys Asp Arg Ile Ser Ser Ser Ala Pro Ser Asp Leu Leu Pro
865 870 875 880

25 Arg Gln Ile Ser Ile Asp Tyr Asp Asp Asp Glu Glu His Phe Gly Leu
885 890 895

30 Asn Phe Val Leu Phe Gly Ser Asn Lys Lys Arg His Ser Gly Lys His
900 905 910

35 Leu Tyr Ser Leu Thr Ile Asn Gly Asp Glu Leu Asp Asp His Leu Ser
915 920 925

40 Asp Thr Phe His Leu Arg His Leu Arg Leu Leu Arg Thr Leu His Leu
930 935 940

45 Glu Ser Ser Phe Ile Met Val Lys Asp Ser Leu Leu Asn Glu Ile Cys
945 950 955 960

50 Met Leu Asn His Leu Arg Tyr Leu Ser Ile Gly Thr Glu Val Lys Ser
965 970 975

55 Leu Pro Leu Ser Phe Ser Asn Leu Trp Asn Leu Glu Ile Leu Phe Val
980 985 990

60 Asp Asn Lys Glu Ser Thr Leu Ile Leu Leu Pro Arg Ile Trp Asp Leu
995 1000 1005

65 Val Lys Leu Gln Val Leu Phe Thr Thr Ala Cys Ser Phe Phe Asp
1010 1015 1020

70 Met Asp Ala Asp Glu Ser Ile Leu Ile Ala Glu Asp Thr Lys Leu
1025 1030 1035

Glu Asn Leu Thr Ala Leu Gly Glu Leu Val Leu Ser Tyr Trp Lys
1040 1045 1050

Asp Thr Glu Asp Ile Phe Lys Arg Leu Pro Asn Leu Gln Val Leu
1055 1060 1065

His Phe Lys Leu Lys Glu Ser Trp Asp Tyr Ser Thr Glu Gln Tyr
1070 1075 1080

Trp Phe Pro Lys Leu Asp Phe Leu Thr Glu Leu Glu Lys Leu Thr
1085 1090 1095

Agrico B.V.

20030596

PF 54801

10

Val Asp Phe Glu Arg Ser Asn Thr Asn Asp Ser Gly Ser Ser Ala
1100 1105 1110

5 Ala Ile Asn Arg Pro Trp Asp Phe His Phe Pro Ser Ser Leu Lys
1115 1120 1125

10 Arg Leu Gln Leu His Glu Phe Pro Leu Thr Ser Asp Ser Leu Ser
1130 1135 1140

15 Thr Ile Ala Arg Leu Leu Asn Leu Glu Glu Leu Tyr Leu Tyr Arg
1145 1150 1155

20 Thr Ile Ile His Gly Glu Glu Trp Asn Met Gly Glu Glu Asp Thr
1160 1165 1170

Phe Glu Asn Leu Lys Cys Leu Met Leu Ser Gln Val Ile Leu Ser
1175 1180 1185

25 Lys Trp Glu Val Gly Glu Glu Ser Phe Pro Thr Leu Glu Lys Leu
1190 1195 1200

30 Glu Leu Ser Asp Cys His Asn Leu Glu Glu Ile Pro Ser Ser Phe
1205 1210 1215

35 Gly Asp Ile Tyr Ser Leu Lys Ile Ile Glu Leu Val Arg Ser Pro
1220 1225 1230

Gln Leu Glu Asn Ser Ala Leu Lys Ile Lys Glu Tyr Ala Glu Asp
1235 1240 1245

40 Met Arg Gly Gly Asp Glu Leu Gln Ile Leu Gly Gln Lys Asp Ile
1250 1255 1260

45 Pro Leu Phe Lys
1265

50 <210> 3
<211> 3890

55 <212> DNA
<213> Solanum bulbocastanum

60 <220>
<221> gene

65 <222> (1)..(3890)
<223> Coding nucleic acid sequence of the Rpi-blb2 gene including the i
ntron sequence (position 43-128).

70 <220>
<221> Intron

Agrico B.V.

20030596

PF 54801

11

<222> (43)..(128)

5 <223> Coding nucleic acid sequence of the Rpi-blb2 gene including the intron sequence (position 43-128).

```

10 <400> 3
   atggaaaaac gaaaagataa tgaagaagca aacaactcat tggatatgta ttgatagag 60
   tgaactgtaa agtattgaat tglagatata atgtggcctt aaaaatttga tatgtgttat 120
15 tttggcagga gtcattttct gctcttcgca aggatgctgc caatgttctg gatctcctag 180
   agagattaaa gaatgaagaa gatcaaaagg ctgttgatgt ggaatcgatt gaaagcctga 240
   aattgaaget gacatttatt tgtacatatg tccagctttc ttattccgat ttggagaagt 300
20 ttgaagatat aatgactaga aaaagacaag aggttgagaa tctgcttcaa ccaatttttg 360
   atgatgatgg caaagacgtc ggggtgtaaat atgtccttac tagcctcgcc ggtaatatgg 420
25 atgactgtat aagcttgtat catcgttcta aatcagatgc caccatgatg gatgagcaat 480
   tgggcttccct cctcttgaat ctctctcctc tatccaagca tctgtctgaa aagatgttct 540
   ctggagtgc tcaatatgag gttcttcaga atgtatgtgg caacataaga gatttccatg 600
30 gattgatagt gaattgttgc attaagcatg agatggttga gaatgtctca tctctgttto 660
   aactgatggc tgagagagta ggacgcttcc tttgggagga tcaggctgat gaagactctc 720
35 aactctccga gctagatgag gatgatcaga atgataaaga cctcdaactc ttcaagctag 780
   cacatctact cttgaagatt gttccaactg aattggaggt tatgcacata tgtcataaaa 840
   ctttgaaagc ttcaacttca acagaaattg gacgcttcat taagaagctc ctggaaacct 900
40 ctccggacat tctcagagaa tatctgattc atctacaaga gcatatgata actgttatta 960
   cccetaacac ttcaggggtc cgaaacatc atgtcatgat ggaattccta ttgattatcc 1020
45 tttctgatat gccgccaag gactttatc atcatgacaa actttctgat ctcttggtc 1080
   gtgttgttagc acttaccagg gaggtatcaa ctcttgtacg cgacttgga gagaaattaa 1140
   ggattaaaga gagtactgac gaaacaaatt gtgcaaccct aaagtctctg gaaaatattg 1200
50 aactccctaa ggaagatctc aaacatgttt atctgaaagt cccggattca tctcaatatt 1260
   gcttccccat gagtcatgga cctctottca tgcattctgt acagagacac ttagatgatt 1320
55 tgctggattc caatgcttat tcaattgctt tgataaagga acaaattggg ctggtgaaag 1380
   aagacttggg attcataaga tctttttctg cgaatattga gcaaggattg tataaagatc 1440
   tctgggaacg tgtcttagat gtggcatatg aggcataaga tgtcatagat tcaattattg 1500
60 ttcgagataa tggctcttta catcttattt tctcacttcc cattaccaga aagaagatga 1560
   tgettattca agaagaggtc tctgatttac atgagaacat ttccaagaac agaggtctca 1620
65 tcgttgtgaa ctctcccaag aaaccagttg agagcaagtc attgacaact gataaaatca 1680
   ttgttaggtt tggtagaggag acaaacttga tacttagaaa gtcaccagt ggatcggcag 1740
   atctagatgt catctcgatc atttggtatgc cgggtttagg taaaactact ctggcgtaaa 1800
70 aagtatacaa tgataaatca gtttctagcc atttcgacct tctgtcatgg tgcacggtcg 1860
   accaagtata tgacgagaag aagctgttgg ataaaatctt caatcaagtt agtgactcaa 1920

```

Agrico B.V.

20030596

PF 54801

12

attcaaaatt gagtgagaat attgatgttg ctgataaact acggaaacaa ttgtcttgaa 1980
agaggtatct tattgtctta gatgacgtgt gggatactaa tacatgggat gagctaacaa 2040
5 gaccccccc tgatgggtatg aaagggaagta gaattatttt gacaactcga gaaaagaaag 2100
ttgtcttgca tggaaagctc tacactgac cctttaacct tcgattgcta agatcagaag 2160
10 aaagtctgga gttattagag aaaagggcat ttggaaacga gagttgccct gatgaactat 2220
tggatgttgg taaagaaata gccgaaaatt gtaaagggct tcctttggtg gtggatctga 2280
ttgctggaat cattgctggg agggaaaaga aaaagagtgt gtggcttgaa gttgtaata 2340
15 atttgcattc ctttattttg aagaatgaag tggaaagtgt gaaagtata gaaataagtc 2400
atgaccactt acctgatcac ctgaagccat gcttgctgta ctttgcaagt gcgccgaagg 2460
actgggtaac gacaatccat gagttgaaac ttatttgggg ttttgaagga tttgtggaaa 2520
20 agacagatat gaagagtctg gaagaagtgg tgaatactta tttggatgat ttaatttcca 2580
gtagcttggg aatttgttcc aatgagatag gtgattacc cacttgccaa ctccatgatc 2640
25 ttgtgcatga cttttgtttg ataaaagcaa gaaagggaaa gttgtgtgat cggataagtt 2700
caagtgtccc atcagatttg ttgccacgtc aaattagcat tgattatgat gatgatgaag 2760
agcacttttg gcttaatttt gtccctgttc gtccaaataa gaaaaggcat tccggtaaac 2820
30 acctctattc ttgaccata aatggagatg agctggacga ccattcttct gatacatttc 2880
atctaagaca cttgaggctt cttagaacct tgcacctga atcctctttt atcatggtta 2940
35 aagattcttt gctgaatgaa atatgcatgt tgaatcattt gaggtactta agcattggga 3000
cagaagttaa atctctgccc ttgtcttctc caaacctctg gaatctagaa atcttgtctg 3060
40 tggataacaa agaatacaac ttgatactat taccgagaat ttgggatctt gtaaagttgc 3120
aagtgtgttt caogactgtc tgttcttctc ttgatattga tgcagatgaa tcaatactga 3180
tagcagagga cacaagttta gagaacttga cagcattagg ggaactcgtg ctttcctatt 3240
45 ggaaagatac agaggatatt ttcaaaaggc tccccaatct tcaagtgtt catttcaaac 3300
tcaaggagtc atgggattat tcaacagagc aatatgggtt cccgaaattg gatttcctaa 3360
ctgaactaga aaaactcaot gtagattttt aaagatcaaa cacaatgac agtgggtcct 3420
50 ctgcagccat aaatcggtca tgggatttct actttccttc gagtttgaaa agattgcaat 3480
tgcataaatt tcctctgaca tccgattcac tatcaacaat agcgagactg ctgaaccttg 3540
55 aagagttgta cttttatctg acaatcatcc atggggaaga atggaacatg ggagaagaaag 3600
acacotttga gaatcccaaa tgtttgatgt tgagtcaagt gattcttctc aagtgggagg 3660
60 ttggagagga atcttttccc acgcttgaga aattagaact gtcggactgt cataatcttg 3720
aggagattcc gtctagtctt ggggatattt attccttgaa aattatcgaa cttgtaagga 3780
gccctcaact tgaataattc gctctcaaga ttaaggaata tgctgaagat atgaggggag 3840
65 gggacgagct tcagattctt ggccagaagg atatcccggt atttaagtag 3890
<210> 4
70 <211> 1267
<212> PRT

Agrico B.V.

20030596

PF 54801

13

<213> Solanum bulbocastanum

5 <220>

<221> protein

10 <222> (1)..(1267)

<223> Deduced Rpi-b1b2 protein sequence

15 <400> 4

Met Glu Lys Arg Lys Asp Asn Glu Glu Ala Asn Asn Ser Leu Glu Ser
1 5 10 15

20

Phe Ser Ala Leu Arg Lys Asp Ala Ala Asn Val Leu Asp Phe Leu Glu
20 25 30

25

Arg Leu Lys Asn Glu Glu Asp Gln Lys Ala Val Asp Val Asp Leu Ile
35 40 45

30

Glu Ser Leu Lys Leu Lys Leu Thr Phe Ile Cys Thr Tyr Val Gln Leu
50 55 60

35

Ser Tyr Ser Asp Leu Glu Lys Phe Glu Asp Ile Met Thr Arg Lys Arg
65 70 75 80

40

Gln Glu Val Glu Asn Leu Leu Gln Pro Ile Leu Asp Asp Asp Gly Lys
85 90 95

45

Asp Val Gly Cys Lys Tyr Val Leu Thr Ser Leu Ala Gly Asn Met Asp
100 105 110

50

Asp Cys Ile Ser Leu Tyr His Arg Ser Lys Ser Asp Ala Thr Met Met
115 120 125

55

Asp Glu Gln Leu Gly Phe Leu Leu Asn Leu Ser His Leu Ser Lys
130 135 140

His Arg Ala Glu Lys Met Phe Pro Gly Val Thr Gln Tyr Glu Val Leu
145 150 155 160

Gln Asn Val Cys Gly Asn Ile Arg Asp Phe His Gly Leu Ile Val Asn
165 170 175

60

Cys Cys Ile Lys His Glu Met Val Glu Asn Val Leu Ser Leu Phe Gln
180 185 190

65

Leu Met Ala Glu Arg Val Gly Arg Phe Leu Trp Glu Asp Gln Ala Asp
195 200 205

70

Glu Asp Ser Gln Leu Ser Glu Leu Asp Glu Asp Asp Gln Asn Asp Lys
210 215 220

Asp Pro Gln Leu Phe Lys Leu Ala His Leu Leu Leu Lys Ile Val Pro

Agrico B.V.

20030596

PF 54801

14

	225		230		235		240
5	Thr Glu Leu Glu Val Met His Ile Cys Tyr Lys Thr Leu Lys Ala Ser	245		250		255	
10	Thr Ser Thr Glu Ile Gly Arg Phe Ile Lys Lys Leu Leu Glu Thr Ser	260		265		270	
15	Pro Asp Ile Leu Arg Glu Tyr Leu Ile His Leu Gln Glu His Met Ile	275		280		285	
20	Thr Val Ile Thr Pro Asn Thr Ser Gly Ala Arg Asn Ile His Val Met	290		295		300	
25	Met Glu Phe Leu Leu Ile Ile Leu Ser Asp Met Pro Pro Lys Asp Phe	305		310		315	320
30	Ile His His Asp Lys Leu Phe Asp Leu Leu Ala Arg Val Val Ala Leu	325		330		335	
35	Thr Arg Glu Val Ser Thr Leu Val Arg Asp Leu Glu Glu Lys Leu Arg	340		345		350	
40	Ile Lys Glu Ser Thr Asp Glu Thr Asn Cys Ala Thr Leu Lys Phe Leu	355		360		365	
45	Glu Asn Ile Glu Leu Leu Lys Glu Asp Leu Lys His Val Tyr Leu Lys	370		375		380	
50	Val Pro Asp Ser Ser Gln Tyr Cys Phe Pro Met Ser Asp Gly Pro Leu	385		390		395	400
55	Phe Met His Leu Leu Gln Arg His Leu Asp Asp Leu Leu Asp Ser Asn	405		410		415	
60	Ala Tyr Ser Ile Ala Leu Ile Lys Glu Gln Ile Gly Leu Val Lys Glu	420		425		430	
65	Asp Leu Glu Phe Ile Arg Ser Phe Phe Ala Asn Ile Glu Gln Gly Leu	435		440		445	
70	Tyr Lys Asp Leu Trp Glu Arg Val Leu Asp Val Ala Tyr Glu Ala Lys	450		455		460	
	Asp Val Ile Asp Ser Ile Ile Val Arg Asp Asn Gly Leu Leu His Leu	465		470		475	480
	Ile Phe Ser Leu Pro Ile Thr Arg Lys Lys Met Met Leu Ile Lys Glu	485		490		495	
	Glu Val Ser Asp Leu His Glu Asn Ile Ser Lys Asn Arg Gly Leu Ile	500		505		510	
	Val Val Asn Ser Pro Lys Lys Pro Val Glu Ser Lys Ser Leu Thr Thr	515		520		525	

Agrico B.V.

20030596

PF 54801

15

5 Asp Lys Ile Ile Val Gly Phe Gly Glu Glu Thr Asn Leu Ile Leu Arg
530 535 540

10 Lys Leu Thr Ser Gly Pro Ala Asp Leu Asp Val Ile Ser Ile Ile Gly
545 550 555 560

15 Met Pro Gly Leu Gly Lys Thr Thr Leu Ala Tyr Lys Val Tyr Asn Asp
565 570 575

20 Lys Ser Val Ser Ser His Phe Asp Leu Arg Ala Trp Cys Thr Val Asp
580 585 590

25 Gln Val Tyr Asp Glu Lys Lys Leu Leu Asp Lys Ile Phe Asn Gln Val
595 600 605

30 Ser Asp Ser Asn Ser Lys Leu Ser Glu Asn Ile Asp Val Ala Asp Lys
610 615 620

35 Leu Arg Lys Gln Leu Phe Gly Lys Arg Tyr Leu Ile Val Leu Asp Asp
625 630 635 640

40 Val Trp Asp Thr Asn Thr Trp Asp Glu Leu Thr Arg Pro Phe Pro Asp
645 650 655

45 Gly Met Lys Gly Ser Arg Ile Ile Leu Thr Thr Arg Glu Lys Lys Val
660 665 670

50 Ala Leu His Gly Lys Leu Tyr Thr Asp Pro Leu Asn Leu Arg Leu Leu
675 680 685

55 Arg Ser Glu Glu Ser Trp Glu Leu Leu Glu Lys Arg Ala Phe Gly Asn
690 695 700

60 Glu Ser Cys Pro Asp Glu Leu Leu Asp Val Gly Lys Glu Ile Ala Glu
705 710 715 720

65 Asn Cys Lys Gly Leu Pro Leu Val Val Asp Leu Ile Ala Gly Ile Ile
725 730 735

70 Ala Gly Arg Glu Lys Lys Lys Ser Val Trp Leu Glu Val Val Asn Asn
740 745 750

75 Leu His Ser Phe Ile Leu Lys Asn Glu Val Glu Val Met Lys Val Ile
755 760 765

80 Glu Ile Ser Tyr Asp His Leu Pro Asp His Leu Lys Pro Cys Leu Leu
770 775 780

85 Tyr Phe Ala Ser Ala Pro Lys Asp Trp Val Thr Thr Ile His Glu Leu
785 790 795 800

90 Lys Leu Ile Trp Gly Phe Glu Gly Phe Val Glu Lys Thr Asp Met Lys
805 810 815

Agrico B.V.

20030596

PF 54801

16

Ser Leu Glu Glu Val Val Lys Ile Tyr Leu Asp Asp Leu Ile Ser Ser
 820 825 830
 5 Ser Leu Val Ile Cys Phe Asn Glu Ile Gly Asp Tyr Pro Thr Cys Gln
 835 840 845
 10 Leu His Asp Leu Val His Asp Phe Cys Leu Ile Lys Ala Arg Lys Glu
 850 855 860
 15 Lys Leu Cys Asp Arg Ile Ser Ser Ser Ala Pro Ser Asp Leu Leu Pro
 865 870 875 880
 20 Arg Gln Ile Ser Ile Asp Tyr Asp Asp Asp Glu Glu His Phe Gly Leu
 885 890 895
 Asn Phe Val Leu Phe Gly Ser Asn Lys Lys Arg His Ser Gly Lys His
 900 905 910
 25 Leu Tyr Ser Leu Thr Ile Asn Gly Asp Glu Leu Asp Asp His Leu Ser
 915 920 925
 30 Asp Thr Phe His Leu Arg His Leu Arg Leu Leu Arg Thr Leu His Leu
 930 935 940
 35 Glu Ser Ser Phe Ile Met Val Lys Asp Ser Leu Leu Asn Glu Ile Cys
 945 950 955 960
 40 Met Leu Asn His Leu Arg Tyr Leu Ser Ile Gly Thr Glu Val Lys Ser
 965 970 975
 Leu Pro Leu Ser Phe Ser Asn Leu Trp Asn Leu Glu Ile Leu Phe Val
 980 985 990
 45 Asp Asn Lys Glu Ser Thr Leu Ile Leu Leu Pro Arg Ile Trp Asp Leu
 995 1000 1005
 50 Val Lys Leu Gln Val Leu Phe Thr Thr Ala Cys Ser Phe Phe Asp
 1010 1015 1020
 55 Met Asp Ala Asp Glu Ser Ile Leu Ile Ala Glu Asp Thr Lys Leu
 1025 1030 1035
 60 Glu Asn Leu Thr Ala Leu Gly Glu Leu Val Leu Ser Tyr Trp Lys
 1040 1045 1050
 Asp Thr Glu Asp Ile Phe Lys Arg Leu Pro Asn Leu Gln Val Leu
 1055 1060 1065
 65 His Phe Lys Leu Lys Glu Ser Trp Asp Tyr Ser Thr Glu Gln Tyr
 1070 1075 1080
 70 Trp Phe Pro Lys Leu Asp Phe Leu Thr Glu Leu Glu Lys Leu Thr
 1085 1090 1095

Agrico B.V.

20030596

PF 54801

17

Val Asp Phe Glu Arg Ser Asn Thr Asn Asp Ser Gly Ser Ser Ala
1100 1105 1110

5 Ala Ile Asn Arg Pro Trp Asp Phe His Phe Pro Ser Ser Leu Lys
1115 1120 1125

10 Arg Leu Gln Leu His Glu Phe Pro Leu Thr Ser Asp Ser Leu Ser
1130 1135 1140

15 Thr Ile Ala Arg Leu Leu Asn Leu Glu Glu Leu Tyr Leu Tyr Arg
1145 1150 1155

Thr Ile Ile His Gly Glu Glu Trp Asn Met Gly Glu Glu Asp Thr
1160 1165 1170

20 Phe Glu Asn Leu Lys Cys Leu Met Leu Ser Gln Val Ile Leu Ser
1175 1180 1185

25 Lys Trp Glu Val Gly Glu Glu Ser Phe Pro Thr Leu Glu Lys Leu
1190 1195 1200

30 Glu Leu Ser Asp Cys His Asn Leu Glu Glu Ile Pro Ser Ser Phe
1205 1210 1215

35 Gly Asp Ile Tyr Ser Leu Lys Ile Ile Glu Leu Val Arg Ser Pro
1220 1225 1230

Gln Leu Glu Asn Ser Ala Leu Lys Ile Lys Glu Tyr Ala Glu Asp
1235 1240 1245

40 Met Arg Gly Gly Asp Glu Leu Gln Ile Leu Gly Gln Lys Asp Ile
1250 1255 1260

45 Pro Leu Phe Lys
1265

50 <210> 5
<211> 7967
<212> DNA

55 <213> Solanum bulbocastanum

60 <220>
<221> genomic_DNA_fragment
<222> (1)..(7967)

65 <223> Sequence of the 7967 bp Sau3AI genomic DNA fragment of ARD 1197-1
6 BAC 211 present in p211F-C12, Rpi-blb2 gene including natural
regulatory elements necessary for correct expression of the gene.
The initiation codon (ATG position 1546-1548) and the termination
codon (TAG position 5433-5435)

70 <220>

Agrico B.V.

20030596

PF 54801

18

<221> stop_codon

<222> (5433)..(5435)

5

<223> Sequence of the 7967 bp Sau3AI genomic DNA fragment of ARD 1197-1
6 BAC 211 present in p211F-C12, Rpi-blb2 gene including natural r
egulatory elements necessary for correct expression of the gene.
The initiation codon (ATG position 1546-1548) and the termination
codon (TAG position 5433-5435)

10

<220>

15

<221> start_codon

<222> (1546)..(1548)

20

<223> Sequence of the 7967 bp Sau3AI genomic DNA fragment of ARD 1197-1
6 BAC 211 present in p211F-C12, Rpi-blb2 gene including natural r
egulatory elements necessary for correct expression of the gene.
The initiation codon (ATG position 1546-1548) and the termination
codon (TAG position 5433-5435)

25

<400> 5

30	gatctagaat caccgaacct cccctcggtta cagctcctcc agttctacca tgaatttcat	60
	ccactgattc ctcttcaatg gccattgcag attctctcga tctatgctca aaaaatcccg	120
	agataaaacc ctagatctgc ttcaaatgct ctgataccat gtaatttcag tgaattctaa	180
35	ctaaacaatg gagagaatta actattcttag aaagactgat tgaaggagaa gaagagagaa	240
	aaattctata ttgaactcat gaacccaaat gaatgaaaaa aataatgaga agaactatac	300
40	tattacaatc tatatatctc tatttatatt ctaatttgaa gcagtttaatt taactgactc	360
	taacaactag actgataggt gtacattttc tgttagtgca ctgcagtgca ttttaactaac	420
	tgcttaacat aaagaatgtt gttcgaactt cattcgaata gcttcaatga gaagcaaaaca	480
45	tgtgtacctg taaagacaca cagtaaaagt gtttaataatg aataaatatg aataaatcaa	540
	ataataaact aaaaataaaa acacatccaa ttaacattgg aggtcttgaa aatcgatggc	600
50	aattaacaaa gaccttctg aaatttaagt ctgtaattga aaatttgagt atagggttagy	660
	ggacatttga ctattttctc attttcttta tctttctctt aatttctggtc agacaagtga	720
	ggaggcccca ctgtaattga ttcattgctt tgctttcttg actttttgga acaatactat	780
55	gcatacatat tgggtctaat tattctctcg tttatttcca gaatttttag ctctatacat	840
	ctaataacaa agcaagcaga ggatatatag tttcatcaac taaaaagggt agtcaactca	900
60	tctaataattt getactctca tctctattga agtacagtta tggaaaagta gaagtgatgt	960
	aagaaaaatg aaagaacttt agtagggttag ttggatctaa caaagagaaa gggaaataaa	1020
	ttgcaggaga aagagagagg cttaaatactt actcacacca ccgatttaca acaaatcact	1080
65	taattgtggt tagttaatgt atactttcac ctcatataat tattacttac ccattgataag	1140
	ttgtattaat ttggtattaa tatccgggtg ggggtgaattc ttaccgggtg agaggggatgg	1200
70	ggttggagag tgtggagtga acagaagcag atgtttttaga ttttttctaa gatgacgaaa	1260
	gattccccct actaatgaaa atatattact atacgctatt agagatagaa aggttcggta	1320
	ccagttggct cegttttctg atgaacccca tttttacaag tcatcttctt caattcaaat	1380

Agrico B.V.

20030596

PF 54801

19

	cgcaagtgtgta cctttatcat cttccactaa ttaagtccto ttaagttcgc gtgaaaatag	1440
5	tgaaattatt gattattcctt atcatttcat cttctttctc ctgataaagt tttatgtact	1500
	ttttatgcat caggtcttga gaacttggaa aggaaaaagta gaatcatgga aaaacgaaaa	1560
	gataatgaag aagcaaacaa ctcatggta tgttatttga tagagtgaac tgtaaagtat	1620
10	tgaattgtag atatcatgtg gctttaaaaa tttgatatgt gttatttttg caggagtcac	1680
	tttctgctct tgcgaaggat gctgccaatg ttctggattt cctagagaga ttaaagaatg	1740
15	aagaagatca aaaggctgtt gatgtggatc tgattgaaag cctgaaattg aagctgacac	1800
	ttatttgtac atatgtccag ctttcttatt ccgatttgga gaagtttgaa gatataatga	1860
	ctagaaaaag acaagagggt gagaatctgc ttcaaccaat tttggatgat gatggcaag	1920
20	acgtcgggtg taaatatgtc cttactagcc tcgcccgtta tatggatgac tgtataagot	1980
	tgtatcatcg ttctaaatca gatgccacca tgatggatga gcaattgggc ttctctctct	2040
25	tgaatctctc tcactatoc aagcactgtg ctgaaaagat gtttcttga gtgactcaat	2100
	atgaggttct tcagaatgta tgtggcaaca taagagattt ccatggattg atagtgaatt	2160
	gttgcattaa gcatgagatg gttgagaatg tcttatctct gtttcaactg atggctgaga	2220
30	gagtaggacg cttccttttg gaggatcagg ctgatgaaga ctctcaactc tccgagctag	2280
	atgaggatga tcagaatgat aaagaccctc aactcttcaa gctagcacat ctactcttga	2340
35	agattgttcc aactgaattg gaggttatgc acatatgtta taaaactttg aaagcttcaa	2400
	cttcaacaga aattggacgc ttcatthaaga agctcttga aacctctcgc gacattctca	2460
	gagaatatct gattcatcta caagagcata tgataactgt tattaacctt aacacttcag	2520
40	gggctcgaaa cattcatgtc atgatggaat tectattgat tattctttct gatatgccgc	2580
	ccaaggactt cattcatcat gacaaacttt ttgatctctt ggtctgtgtt gtgacactta	2640
45	ccagggaggt atcaactctt gtacgcgact tggaagagaa attaaggatt aaagagagta	2700
	ctgacgaaac aaattgtgca accctaaagt ttctggaaaa cattgaaactc cttaaaggaag	2760
	atctcaaaaca tgttatctg aaagtcccg attcatctca atattgcttc cccatgagtg	2820
50	atggacctct cttcatgcat ctgctacaga gacacttaga tgatttgtgt gattccaatg	2880
	cttattcaat tgccttgata aaggaaacaaa ttgggtctgt gaaagaagac ttggaattca	2940
55	taagatcttt tttcggaat attgagcaag gattgtataa agatctcttg gaactgttc	3000
	tagatgtggc atatgaggca aaagatgtca tagattcaat tattgttcga gataatggtc	3060
	tcttacatct tatttcttca ctccccatta ccagaaagaa gatgatgott atcaaagaag	3120
60	aggtctctga tttacatgag aacatttcca agaacagagg tctcatcgtt gtgaactctc	3180
	ccaagaaacc agttgagago aagtcattga caactgataa aataattgta ggttttggctg	3240
65	aggagacaaa cttgatactt agaaagctca ccagtggacc ggcagatcta gatgtcattt	3300
	cgatcatttg tatgccgggt ttaggtaaaa ctactttggc gtacaaagta tacaatgata	3360
	aatcagtttc tagccatttc gaccttcgtg catggtgcac ggtcgaccaaa gtatatgacg	3420
70	agaagaagtt gttggataaa attttcaatc aagttagtga ctcaaattca aaattgagtg	3480
	agaatattga tgttgctgat aaactacgga aacaattggt tggaaagagg tatcttattg	3540

Agrico B.V.

20030596

PF 54801

20

	tcttagatga	cgtgtgggat	actaatacat	gggatgagct	aacaagacct	tttccctgatg	3600
	gtatgaaagg	aagtagaatt	attttgacaa	ctcgagaaaa	gaaagtgtgt	ttgcatggaa	3660
5	agctctacac	tgatcctctt	aaccttcgat	tgctaagatc	agaagaaagt	tgggagttat	3720
	tagagaaaag	ggcatttgga	aacgagaggt	gccttgatga	actattggat	gttggttaaag	3780
10	aaatagccga	aaattgtaaa	gggcttcctt	tgggtgtgga	tctgattgtt	ggaatcattg	3840
	ctgggaggga	aaagaaaaag	agtgtgtggc	ttgaagtgtt	aaataatttg	catlccttta	3900
	ttttgaagaa	tgaagtggaa	gtgatgaaag	ttatagaaat	aagttatgac	cacttacctg	3960
15	atcacctgaa	gccatgcttg	ctgtaqcttg	caagtgcgcc	gaaggactgg	gtaacgacaa	4020
	tccatgagtt	gaaacttact	tggggttttg	aaggatttgt	ggaaaagaca	gatatgaaga	4080
20	gtctggaaga	agtgtgaaa	atttattttg	atgatttaat	ttccagtagc	ttggtaattt	4140
	gtttcaatga	gataggtgat	tacctactt	gccaaacttca	tgatcttgtg	catgactttt	4200
	gttctgataa	agcaagaaag	gaaaagttgt	gtgatcggat	aagttcaagt	gtcccatcag	4260
25	atttgttgcc	acgtcaaatt	agcatttgatt	atgatgatga	tgaagagcac	tttgggctta	4320
	attttgtcct	gttcggttca	aataagaaaa	ggcattccgg	taaacacctc	tattctttga	4380
30	ccataaatgg	agatgagctg	gacgaccatc	tttctgatac	atctcatcta	agacacttga	4440
	ggcttcttag	aaccttgcac	ctggaatcct	cttttatcat	ggtaaaagat	tctttgotga	4500
	atgaaatatz	catgttgaat	catttgaggt	acttaagcat	tgggacagaa	gttaaatctc	4560
35	tgcctttgtc	tttctcaaac	ctctggaatc	tagaaatctt	gtttgtggat	aacaaagaat	4620
	caaccttgat	actattaccg	agaatttggg	atcttgtaaa	gttgcaagtg	ctgttcacga	4680
40	ctgcttgttc	tttctttgat	atggatgcag	atgaatcaat	actgatagca	gaggacacaa	4740
	agttagagaa	cttgacagca	ttaggggaac	tcgtgttttc	ctattggaaa	gatacagagg	4800
	atattttcaa	aaggcttccc	aatcttcaag	tgcttcattt	caaaactcaag	gagtcattgg	4860
45	attattcaac	agagcaatat	tggttccoga	aattggattt	cctaactgaa	ctagaaaaac	4920
	tcactgtaga	ttttgaaaga	tcaaacacaa	atgacagtgg	gtcctctgca	gccataaatc	4980
50	ggccatggga	ttttcacttt	ccttcgagtt	tgaaaagatt	gcaattgcac	gaatttcctc	5040
	tgacatccga	ttcactatca	acaatagcga	gactgctgaa	ccttgaagag	ttgtaccttt	5100
	atogtaoaat	catccatggg	gaagaatgga	acatggggaga	agaagacacc	tttgagaatc	5160
55	tcaaatgttc	gatgttgagt	caagtgtatc	tttccaagtg	ggagggttga	gaggaatctt	5220
	ttcccacgct	tgagaaatta	gaactgtcgg	actgtcataa	tcttgaggag	altccgtcta	5280
60	gttttgggga	tatttatccc	ttgaaaatta	tcgaacttgt	aaggagccct	caacttgaaa	5340
	attccgctct	caagattaag	gaatatgctg	aagatatgag	gggaggggac	gagcttcaga	5400
	tccttggcca	gaaggatata	ccgttattta	agtagttttt	gagcattatg	gttgaaaagt	5460
65	agattgcact	ttgctgggta	gattgtatat	ggtaagaaa	attctgttac	agttgtctatg	5520
	aaacattttt	atttgacttt	tctgagtttc	ttttagaaaa	ctcagaagtt	tttaacaaaa	5580
70	attatagttt	ttataaatac	aatgtggatt	tgcttttggc	tgtccaactt	ggtctgaagt	5640
	ctcatatgct	cagagcacta	tcgttcaacc	tcaatcaagg	tactgattta	aaatgacatc	5700
	tatactactt	tatcacaaac	ccaacgaact	ttcatctcaa	aagctaggcc	aggaagtga	5760

Agrico B.V.

20030596

PF 54801

21

	gagggtttag agagcttata agcactcatg acttcccttt ctggaacatt caaccaacgt	5820
5	aggctgaaat cccactctga acgaaaataa gtgtttgttt atcaaattaa ctctcgtagt	5880
	agaacactga aataccttct tctaaacgtt caacaaatgg gatttccagc actcaaagtg	5940
	aatgaaaggt tcacattaat cttcaaaaag aattacgaca attcatgacc acaagtacat	6000
10	tgacagcacc atttcaacag aagaacaagt caatgctgca tcttcatcaa taatccgagt	6060
	gtcgaacctc ettcctgaca ctgtcctgta tatgtaaagt tcttcaacag ggcaacttcc	6120
15	tggtctcgta tctggatgac cctctcgtc tataacttca acattaagcc ctggcaactt	6180
	ctggaccaac agcttacatg cttcaaaact tactgaacaa ttagacatcc aaagggatcg	6240
	cattgtctcc agctttgcag cattagocaa cagagcctca tcgccaaagg ggcagtctct	6300
20	aatctcgaa ttgaaaaaat tgtctgtgta tgactttcct ctgacatccg atgcactatc	6360
	aacaatagca agactggagg ttggagagga atcctttatt atacaatcat tcagggagaa	6420
25	gaatggaaca tgggggagga agacactttt gagaatctga aatgtgttag agccacaagc	6480
	tacagaagta ttgaatttgt catgaataac aacattcttc atcctagtta attctttttc	6540
	aatttttaat agactctcat tttaatcact aatattcttc tattgtgac tctttttctg	6600
30	cagggtggcaa ctttaaattc ataaaagtata ggattgatga caaactcgaa aaatatctta	6660
	atgaggtgaa gtttgagcag tcagcagatg gtggttccaa ctctaagttg acaagcacat	6720
35	actatcccgg agggcgattt caagcotgat gcatacgtt agtgtggcta gagcagacag	6780
	gatgtattac ctggatatct accaagacga atccacaatc agttttatgt caagcaatac	6840
	atgaagtaac tcccgataga acagtaaaag caagatgtgt aggtgtatct cgaacttaag	6900
40	agattgtaca ttctctcttg agatttttac tgctaataca aatttacacc tcagaagcga	6960
	atctagaatt tctagagcat gaatgcacca ctaatgaaag gagaaaaaag gaagtatgaa	7020
45	gtgggaattt gatccttgtt tctaggata taaaatttat cattcaacta tacttcattt	7080
	agcaaacac tctctttgac attatttctc aaacaagggg tctctaattt gctaaactaa	7140
	agactgtcaa aaggtaagtt catcttcaca ctctcttgtt tactttatct aaaggggaac	7200
50	catgaaaaac aagaacatc agaatgtcc cgtaaaacaaa gcagcctcat gcacaaaaca	7260
	tccaacgttg gtaggattaa tggagggatc gcattcccag aggatactgt agaaaaatta	7320
55	gtggcttctt tcaccgctca aaccatgat ctataggta catggagaca acttcatggt	7380
	tgctcgtagg ctcccgtaaa ttctcataaa ccacaaacac aaagttgcat cagacatcat	7440
	cttcattcac aagctgacaa tctccacaag tottagtcaa cttgtaatat gaatattagc	7500
60	caggtagacg tacatattta caaaattgag tttcctatat aatatggttt gaaggaaatga	7560
	aacatgatgg ggagggtaga taaaataata tatgaggcat aaaaatagga aagatatttg	7620
65	tagtgagagg ttttgacttt ttatgctgt tttgatcttc agttctctgt attctttttc	7680
	tactgcttct ctctcttttc tctgagtaa agttttatgt aggtatcttt tatagctccg	7740
	atcgtgagaa cttgaaagaa agctctctat agctatgcta ggtgcccaca taaaaaatg	7800
70	aaatattaca aaaaccctga taataaaata cactaatcta agatattcac tgcaacatac	7860
	atgcaaaaata tatatatata aattttcatg aaattataa caaataatag atgtgaacat	7920

Agrico B.V.

20030596

PF 54801

22

ataacttttaa aaataatatt acatocataa agcttaaatt ctagatc

7967

- 5 <210> 6
 <211> 9949
 <212> DNA
- 10 <213> Solanum bulbocastanum
- 15 <220>
 <221> genomic_DNA_fragment
 <222> (1)..(9949)
- 20 <223> Sequence of 9949 bp Sau3AI genomic DNA fragment of S. bulbocastanum 2002 BAC BlbSP39 present in pSP39-20. The genomic fragment harbours the Rpi-blb2 gene including natural elements necessary for expression. Initiation codon (ATG position 1413-1415), the termination codon (TAG position 5300-5303)
- 25
- 30 <220>
 <221> start_codon
 <222> (1413)..(1415)
- 35 <223> Sequence of 9949 bp Sau3AI genomic DNA fragment of S. bulbocastanum 2002 BAC BlbSP39 present in pSP39-20. The genomic fragment harbours the Rpi-blb2 gene including natural elements necessary for expression. Initiation codon (ATG position 1413-1415), the termination codon (TAG position 5300-5303)
- 40
- 45 <220>
 <221> stop_codon
 <222> (5300)..(5303)
- 50 <223> Sequence of 9949 bp Sau3AI genomic DNA fragment of S. bulbocastanum 2002 BAC BlbSP39 present in pSP39-20. The genomic fragment harbours the Rpi-blb2 gene including natural elements necessary for expression. Initiation codon (ATG position 1413-1415), the termination codon (TAG position 5300-5303)
- 55
- 60 <400> 6
 gatctgcttc aaatgctctg ataccatgta atttcagtga attctaacta aacaatggag 60
 agaatttaact attcttagaaa gactgattga aggagaagaa gagagaaaaa ttcttatattg 120
 aactcatgaa ccaaaatgaa tgaaaaaaat aatgagaaga actataactat tacaatctat 180
 atatctctat ttatatctta atctgaagca gttaatttaa ctgactctaa caactagact 240
- 65 gataggtgta cattttctgt tagtgacttg cagtgcattt aactaactgc ttaacataaa 300
 gaatgttggt cgaacttcatt tgaatagct tcaatgagaa gcaaacatgt gtacctgtaa 360
- 70 agacacacag taaaagtgtt aataatgaat aaatatgaat aaatcaaata ataaattaaa 420
 aataaaaaca catccaatta acattggagg tcttgaaaat cgatggtaat taacaaagac 480
 ccttgtagaa ttttaagtctg taattgaaaa tttagagtata ggtagggga catttgacta 540

Agrico B.V.

20030596

PF 54801

23

	ttttctcatt ttctttatct ttttctaat ttgtggcaga caagtgagga ggccccactg	600
5	taattgatcc atgcttttgc tttcttgact ttttggaaca atactatgca tcataatttg	660
	tccttaattat tccctctgtt atttccagaa ttttgagctc tatacatcta ataacaagc	720
	aagcagagga catatagttt catcaactaa aaaggttagt caactcatct aatatttgct	780
10	actctcatct ctattgaagt acagttatgg aaaagtagaa gtgatgtaag aaaaatgaaa	840
	gaacttttagt aggttagttg gatctaacaa agagaaaggg aaataaattg caggagaaag	900
15	agagagggtta aatacttact cacaccaccg atttacaaca aatcacttaa ttgtggctag	960
	ttaattgtata ctttcacctc attaaattat tacttaccca tgataagttg tattaatttg	1020
	gtattaatat ccggtgcggg tgaattctta ccgggtgaga gggatggggg tggagagtgt	1080
20	ggagtgaaca gaagcagatg ttttagattt tttctaagat gacgaaagat tccccctact	1140
	aatgaaaata tattactata cgotattaga gatagaaagg ttcggtacca gttggtctcg	1200
25	tttctggatg aaccccatth ttacaagtca ttttcttcaa ttcaaactgc aagtgtacct	1260
	ttatcatctt ccactaatta agtcctctta agttcggtg aaaaatagtg aattattgat	1320
	tactcttact atttcatctt ctttctcttg ataaagtctt atgtactttt catgcatcag	1380
30	gtcttgagaa cttggaaagg aaaagtagaa tcatggaaaa ccgaaaagat aatgaugaag	1440
	caaacaactc attggtatgt tatttgatag agtgaactgt aaagtattga attgtagata	1500
35	tcattgtggct ttaaaaaatt gatcatgctt attttggcag gagtattttt ctgctcttcg	1560
	caaggatgct gccaatgttc tggatttctt agagagatta aagaatgaag aagatcaaaa	1620
	ggctgttgat gtggatctga ttgaaagcct gaaattgaag ctgacattta tctgtacata	1680
40	tgtccagctt tcttattcog atttggagaa gtttgaagat ataattgacta gaaaaagaca	1740
	agaggttgag aatctgcttc aaccaatttt ggatgatgat ggcaaaagac tggggtgtaa	1800
45	atatgtcctt actagcctcg ccggtaatat ggatgactgt ataagcttgc atcatcgctt	1860
	caaactagat gccaccatga tggatgagca attgggcttc ctctcttga atctctctca	1920
	tctatccaag catcgtgctg aaaagatgtt tcctggagtg actcaatatg aggttcttca	1980
50	gaatgtatgt ggcaacataa gagatttcca tggattgata gtgaattgtt gcattaagca	2040
	tgagatgggt gagaatgtct tatctctgtt tcaactgatg gctgagagag taggacgctt	2100
55	cctttgggag gatcagggtg atgaagactc tcaactctcc gagctagatg aggatgatca	2160
	gaatgataaa gaccctcaac tcttcaagct agcaatcta ctcttgaaga ttgttccaac	2220
	tgaattggag gttatgcaca tatgttataa aactttgaaa gcttcaactc caacagaaat	2280
60	tggacgcttc attaagaagc tcctggaaac ctctccggac attctcagag aatatctgat	2340
	tcactctaaa gagcatatga taactgttat taccctaac acttcagggg ctcgaaacat	2400
65	tcattgtcatg atggaattcc tattgattat tctttctgat atgccgccca aggactttat	2460
	tcattcatgac aaactttttg atctcttggc tegtgtgtga gcacttacca gggaggatc	2520
	aactcttgta cgcgacttgg aagagaaatt aaggattaaa gagagtactg acgaaacaaa	2580
70	tcgtgcaacc cttaaagttc tggaaaatat tgaactcctt aaggaagatc tcaaacatgt	2640
	ttatctgaaa gtcccggatt catctcaata ttgcttcccc atgagtgatg gacctctctt	2700

Agrico B.V.

20030596

PF 54801

24

	catgcacctg ctacagagac acttagatga ttctgtggat tccaatgctt attcaattgo	2760
	tttgataaag gaacaaattg ggctgggtgaa agaagacttg gaattcataa gatctctctt	2820
5	cgcgaaatatt gagcaaggat tgtataaaga tctctgggaa cgtgttctag atgtggcata	2880
	tgaggcaaaa gatgtcatag attcaattat tgttcgagat aatggctctt tacatcttat	2940
10	cttctcactt cccattacca gaaagaagat gatgcttatt aaagaagagg tctctgattt	3000
	acatgagAAC atttccaaga acagaggtct catcgttggt aactctccca agaaaccagt	3060
	tgagagcaag tcattgacaa ctgataaaat aattgtagggt ttgggtgagg agacaaactt	3120
15	gatacttaga aagctcacca gtggaccggc agatctagat gtcattctga tcattggtat	3180
	gccgggttta ggtaaaaata ctttggcgta aaggtatata aatgataaat cagtttctag	3240
20	ccatttctgac cttcgtgcat ggtgcacggc cgaccaagta tatgacgaga agaagtgtt	3300
	ggataaaatt ttcaatcaag ttagtgactc aaattcaaaa ttgagtgaga atattgatgt	3360
	tgctgataaa ctacggaaac aattgtttgg aaagagggtat cttattgtct tagatgacgt	3420
25	gtgggatact aatcacatgg atgagctaac aagacctttt cctgatggta tgaaaggaag	3480
	tagaattatt ttgacaactc gagaaaagaa agttgctttg catggaaagc tctacactga	3540
30	tcctcttaac cttcgattgc taagatcaga agaaagtggg gatttattag agaaaagggc	3600
	atttggaaac gagagtggcc ctgatgaact attggatgtt ggtaaagaaa tagccgaaa	3660
	ttgtaaaggg cttcctttgg tgggtggatct gattgctgga atcattgctg ggagggaaaa	3720
35	gaaaaagagt gtgtggcttg aagtgtgaaa taatttgcat tctcttattt tgaagaatga	3780
	agtggaaagt atgaaagtta tagaaataag ttatgaccac ttacctgac acctgaagcc	3840
40	atgotttctg taotttgcaa gtgcgccgaa ggactgggta acgacaatcc atgagttgaa	3900
	acttatttgg ggttttgaag gatttgtgga aaagacagat atgaagagtc tggaagaagt	3960
	ggtgaaaatt tatttggatg atttaatttc cagtagcttg gtaatttgtt tcaatgagat	4020
45	aggtgattac cctacttgcc aacttcattg tcttgtgcat gacttttgtt tgataaaagc	4080
	aagaaaggaa aagtgtgtgt atcgggataag ttcaagtgtt ccatcagatt tgttgcoacg	4140
50	tcaaattagc attgattatg atgatgatga agagcacttt gggcttaatt ctgtcctgtc	4200
	cggttcaaat aagaaaaggc attccggtta acacctctat tctttgacca taaatggaga	4260
	tgagctggac gaccatcttt ctgatacatt tcactaaga cacttgaggc ttcttagaac	4320
55	cttgacactg gaatctctct ttatcatggt taaagattct ttgctgaatg aaatatgcat	4380
	gttgaatcat ttgaggtact taagcattgg gacagaagtt aaatctctgc ctttgtctct	4440
60	ctcaaacctc tggaatctag aaatcttgtt tgtggataac aaagaatcaa ccttgatact	4500
	attaccgaga atttgggac ttgtaaagtt gcaagtgtg ttacagactg cttgttcttt	4560
	cttctgatag gatgcagatg aatcaatact gatagcagag gacacaaagt tagagaactt	4620
65	gacagcatta ggggaactcg tgctttctta ttggaaagat acagaggata ttttcaaaag	4680
	gttctcgaat ottcaagtgc ttcatctcaa actcaaggag tcatgggatt attcaacaga	4740
70	gaaatattgg tccccgaaat tggatttctt aactgaacta gaaaaactca ctgtagattt	4800
	tgaaagatca aacacaaatg acagtgggtc ctctgcagcc ataaatcggc catgggattt	4860
	tcactttctt tcgagtttga aaagattgca attgcatgaa tttctctctg catccgattc	4920

Agrico B.V.

20030596

PF 54801

25

	actatcaaca atagggagac tgetgaacct tgaagagttg tacctttatc gtacaatcat	4980
5	ccatggggaa gaatggaaca tgggagaaga agacaccttc gagaatctca aatgtttgat	5040
	gtttagtcaa gtgattcttc ccaagtggga gggtggagag gaatcttttc ccacgcttga	5100
	gaaattagaa ctgtcggact gtcataatct tgaggagatt ccgtctagct ttggggatat	5160
10	ttattctctg aaaattatcg aacttgtaag gagccctcaa cttgaaaatt ccgcrctcaa	5220
	gattaaggaa tatgctgaag atatgagggg aggggaagag cttcagatcc ttggccagaa	5280
15	ggatatcccg ttatttaagt agtttttgag cattatggct gaaaagtaga ttgcactctg	5340
	ctgggtagat tgtatatggt taagaaaatt ctgttacagt tgttatgaaa catctttact	5400
	tgacctttct gagtttcttc tagaaaaactc agaagttttt aacaaaaatt atagtcttta	5460
20	taaacacaat gtggatttgc ctttggtctgt ccaacttggc ctgaagcttc atatgctcag	5520
	agcactatcg ttcaacctca atcaagggtac tgatttaaaa tgacatctat actactttat	5580
25	cacaaaccca acgaactttc atctcaaaag ctaggccagg aagtgaagag gttgtagaga	5640
	gcttataaagc actcatgact tccctttctc gaacattcaa ccaacgtagg ctgaaatccc	5700
	actctgaacg aaaataagtg tctgtttatc aaactaactc tcgtagtaga acactgaaat	5760
30	accttcttct aaacgttcaa caaatgggat ttccagcact caaagtgaat gaaagggtca	5820
	cattaatctt caaaaagaat tacgacaatt catgaccaca agtacattga cagcaccatt	5880
35	tcaacagaag aacaagtcaa tgetgcatct tcatcaataa tccgagtgtc gaancctctt	5940
	cctgcactcg tctgtatct gtaaaagttc tcaacagggc aactttctcg tctcgtatct	6000
	ggatgacccc tctcgtctat aacttcaaca ttaagccctg gcaacttctg gaccaacagc	6060
40	ttacatgctt caaaacttac tgaacaatta gacatccaaa gggatcgcat tgcctccagc	6120
	ttcgcagcat tagccaacag agcctcatcg ccaaaggggc agtctctaatt ctgcaatttg	6180
45	aaaaaattgt tgttgatga ctttctctcg acatccgatg cactatcaac aatagcaaga	6240
	ctggagggtg gagaggaatc ctttattata caatcattca gggagaagaa tggaaatggg	6300
	gggaggaaga cacttttgag aatctgaaat gtgttagagc cacaagctac agaagtattg	6360
50	aatttgctcat gaatatcaac attcttctat ctagttaatt ctttttcaat ttttaataga	6420
	ctctcatttt aatcactaat attcttctat ttgtgacttc ttttctgcag gtggcaactt	6480
55	taaattcata aagtatagga ttgatgacaa actcgaaaaa tatcttaatg aggtgaagtt	6540
	tgagcagtoa gcagatggtg gtcccaactc taagttgaca agcacatact atcccggagg	6600
	gcgatttcaa gcctgatgca tatgggttagt gtggctagag cagacaggat gtattacctg	6660
60	gatatctacc aagacgaatc cacaatcagt tttatgtcaa gcaatcacatg aagtaactcc	6720
	cgatagaaca gtaaaagcaa gatgtgtagg tgtatctcga ctctaagaga ttgtacattc	6780
65	ctctttgaga tttttaactg caatacaaat ttacacctca gaagcgaatc tggaaattctt	6840
	agagcatgaa tgcaccacta atgaaaggag aaaaaaggaa gtatgaagtg ggaatttgat	6900
	ccttggtttct aggtatataa aatttatcat tcaactatac ttcatattago aaacaactct	6960
70	ctttgcoatt atttctcaaa caagggcttc taatattgct aaactaaaga ctgtcaaaag	7020
	gtaagttcat cttcaaaactc tcttggtttac tttatctaaa ggggaactat gaaaaacaag	7080

Agrico B.V.

20030596

PF 54801

26

	aaacatcagg	aatgtcccg	aaacaaagca	gcctcatgca	caaaacatcc	aaogttggta	7140
	ggattaatgg	agggatcgca	ccccaggagg	atacctgtaga	aaaattagtg	gcttctttca	7200
5	ccgtcaaac	ccatgatcta	taggttacat	ggagacaact	ttatggttgc	tcgtaggctc	7260
	cogtcaattc	tcataaacca	caacacccaa	gttgcacag	acatcatctt	cattcacaag	7320
10	ctgacaatct	ccacaagtct	tagtcaactt	gtaatatgaa	tattagccag	gtagacgtac	7380
	atatttacaa	aattgagttt	cctatataat	atggcttgaa	ggaatgaaac	atgatgggga	7440
	gggtagataa	aataatata	gaggcataaa	aataggaaag	atattttag	tgagagggtt	7500
15	tgacttttta	tgetgctttt	gatcttcagt	ttcttgtatt	ctttttctac	tgctttctct	7560
	ttctttctcc	tgagtaagt	tttatgtagg	tactttttat	acgtccgac	gtgagaactt	7620
20	gaaagaaagc	tctctatagc	tatgttaggt	gcccacataa	aaaaatgaa	tattacaaa	7680
	acctgataa	taaaatacac	taactaaga	tattcactgc	aacatacatg	caaaatata	7740
	atatataaat	ttcatgaaa	attataacaa	ataatagatg	tgaacatata	actttaaaaa	7800
25	taatattaca	tcataaagc	ttaaattcta	gatccatcta	tgctgtatg	atgcatagct	7860
	cagaatatct	ccatcaagt	ttaaactaca	tatttcattc	aaatttatat	agaaaacgat	7920
30	aattaagggtg	aaaactttta	taaagatata	gtgtggttgt	gtgagtggag	tgacaaaata	7980
	agttgtgtga	ttattcaaaa	agttttaata	acgaaaatcc	acatgcttga	attaattgaa	8040
	gcattaatgt	tgraacgaaa	aatattacat	ttattgagtt	actgtgatgt	tttaactgat	8100
35	atataaaata	atattggtat	ttctcttcac	ctgcgacata	atctgttttt	tcactttttt	8160
	tcaatataca	aaatagaatt	attattttgt	tgcatctttt	taagtacaaa	ttattcatat	8220
40	gtatatagta	caaaataaaa	tatttactgt	ggtaaagtaa	atggaaataag	aggtcatatt	8280
	tgaaataaca	atatactata	ctatgttaaa	gtatttttta	tagtcaaaat	ttctctagag	8340
	tacttgattc	tacatacaaa	tactaatttc	gtaaaaaaat	caalattgaa	ttctttcatt	8400
45	gtttctttat	tattaaatta	gtttataata	actaaactaa	ggtaataaga	ccttagttta	8460
	gttaatgtgt	gtctctgtga	tttcgttcac	agtotaaggg	tgtacttgtg	ccttatccca	8520
50	aaaatgaagg	aatatcaaaa	gatataattaa	aattaaatta	aatatttggg	ggttatgaat	8580
	ataaaaagta	tcagagttct	acatataaag	agtaacaatt	gaaataatta	attaatatg	8640
	agatatgaag	goggacattt	aaagaaaata	ataaataaat	aaattaaagg	gtataaattt	8700
55	cataatacat	aataccaata	agccgtagaa	tatctcogtc	ataatgcata	aactaataaa	8760
	tcacaaatgt	ataactcaca	tacaaatatt	tttgcataaa	gaatttgaat	gttgtaatag	8820
60	aatggagaat	aacttgtgtc	ttattccatt	atgtaagacg	tataaataca	aatacaatga	8880
	gctctaatta	attaaggaaa	ctaaataagg	aaggaatcaa	aaaatattat	gtcatatccc	8940
	tacatatctg	ctagagattc	tatcatatcc	ttacatatct	gttaagctat	gtctacacct	9000
65	aaagggtgtct	acaatcattt	tgtaacactc	cccccaagt	tagagcatag	atattattcca	9060
	ttcccaactt	gttacaaaag	taatcaactc	gagttccatt	caacgctttt	gtgaacaaat	9120
70	caactagttg	ctctccgtgc	ttcacttagc	tagtggatat	caggttttta	tgaactctct	9180
	cacgaataaa	atgacagtc	acctcaatat	gttttagttc	ttcatgagac	accggattca	9240
	aggcaatatg	gagcgcaact	tgattatcat	actagagctt	tgatggtata	tgatgcttca	9300

Agrico B.V.

20030596

PF 54801

27

accctatttc tgttaaaaga taatgtatcc acatgatctc acccatagac tgtaacataa 9360
 5 ctctgtactt tgattctgca ctagatcaag atacaacatt ttgcttttta ctctccatg 9420
 ataccaggtt tcatacaaca aagacacaat aacttgtagt agatcttcta tcaattttctg 9480
 atccagccca atcgacatct gcaaaacact caatatgagt atggctcgtga ttttgatact 9540
 10 atattccaag actaggagtt ttcttcaagt aacatagaat atgttccaaa gctgcccagt 9600
 gtttgacgta ggtgcaaaaca tgaactagct aacaacactt actgcaaaaag caatatcaag 9660
 atgagtcaca ataaggtagt ttaactttcc aactaacctt ttgtatctct atggatcatt 9720
 15 aaaaggatcg tcgtcatctc tcataagatg catattggga accattggag aaottcaggg 9780
 tttggctgcc atctttcaat tttctgcaag tagatcgaga gaatatattc tctaagacaa 9840
 20 aagaattccc tttttgttcc tatttacttc tactcccaaa atgtatttca attgacccaa 9900
 gtccttcgta tgaaaccaag tatgcaggaa agacttgagg gaagagatc 9949
 25 <210> 7
 <211> 3768
 <212> DNA
 30 <213> Lycopersicon lycopersicum
 35 <220>
 <221> CDS
 <222> (1)..(3768)
 40 <223> Mil.1 from tomato
 45 <400> 7
 atg gaa aaa cga aaa gat aat gaa gaa gca aac aac tca ttg gtg cta 48
 Met Glu Lys Arg Lys Asp Asn Glu Glu Ala Asn Asn Ser Leu Val Leu
 1 5 10 15
 50 ttt tct gct ctt agc aag gac att gcc gat gtt ctg gtt ttc cta gag 96
 Phe Ser Ala Leu Ser Lys Asp Ile Ala Asp Val Leu Val Phe Leu Glu
 20 25 30
 55 aat gag gaa aat caa aaa gct ctt gac aaa gat caa gtt gaa aag ata 144
 Asn Glu Glu Asn Gln Lys Ala Leu Asp Lys Asp Gln Val Glu Lys Ile
 35 40 45
 60 aaa ttg aaa atg gca ttt att tgt aca tat gtt cag ctt tct tgt tcc 192
 Lys Leu Lys Met Ala Phe Ile Cys Thr Tyr Val Gln Leu Ser Cys Ser
 50 55 60
 gat ttt gag cag ttt gaa gat ata atg act aga aaa aga caa gag gtt 240
 Asp Phe Glu Gln Phe Glu Asp Ile Met Thr Arg Lys Arg Gln Glu Val
 65 70 75 80
 65 gag aat ctg ctt caa cca ctt ttg gat gat gat gtc ttt act agc ctc 288
 Glu Asn Leu Leu Gln Pro Leu Leu Asp Asp Val Phe Thr Ser Leu
 85 90 95
 70 acc agt aat atg gat gac tgt atc agc ttg tat cat cgt tct tat aaa 336
 Thr Ser Asn Met Asp Asp Cys Ile Ser Leu Tyr His Arg Ser Tyr Lys
 100 105 110

Agrico B.V.

20030596

PF 54801

28

	tca	gat	gcc	atc	atg	atg	gat	gag	caa	ttg	gac	ttc	ctc	ctc	ttg	aat	384
	Ser	Asp	Ala	Ile	Met	Met	Asp	Glu	Gln	Leu	Asp	Phe	Leu	Leu	Leu	Asn	
			115					120					125				
5	ctc	tat	cat	cta	tcc	aag	cat	cac	gct	gaa	aag	ata	ttt	cct	gga	gtg	432
	Leu	Tyr	His	Leu	Ser	Lys	His	His	Ala	Glu	Lys	Ile	Phe	Pro	Gly	Val	
		130					135					140					
10	act	caa	tat	gaa	gtt	ctt	cag	aat	ata	tgt	ggc	aac	ata	aga	gat	ttc	480
	Thr	Gln	Tyr	Glu	Val	Leu	Gln	Asn	Ile	Cys	Gly	Asn	Ile	Arg	Asp	Phe	
	145					150					155					160	
15	cat	ggg	ttg	ata	gtg	aat	ggg	tgc	att	aag	cat	gag	atg	gtt	gag	aat	528
	His	Gly	Leu	Ile	Val	Asn	Gly	Cys	Ile	Lys	His	Glu	Met	Val	Glu	Asn	
					165					170					175		
20	gtc	ttr	cct	ctg	ttt	caa	ctc	atg	gct	gac	aga	gta	gga	cac	ttc	ctt	576
	Val	Xaa	Pro	Leu	Phe	Gln	Leu	Met	Ala	Asp	Arg	Val	Gly	His	Phe	Leu	
				180					185					190			
25	tgg	gat	gat	cag	act	gat	gaa	gac	tct	cga	ctc	tcc	gag	cta	gat	gag	624
	Trp	Asp	Asp	Gln	Thr	Asp	Glu	Asp	Ser	Arg	Leu	Ser	Glu	Leu	Asp	Glu	
			195					200					205				
30	gat	gaa	caa	aat	gat	aga	gac	tct	cga	ctt	ttc	aag	cta	gca	cat	cta	672
	Asp	Glu	Gln	Asn	Asp	Arg	Asp	Ser	Arg	Leu	Phe	Lys	Leu	Ala	His	Leu	
							215					220					
35	ctc	ttg	aag	atc	gtt	ccg	gtt	gaa	ctg	gag	gtt	ata	cac	ata	tgt	tat	720
	Leu	Leu	Lys	Ile	Val	Pro	Val	Glu	Leu	Glu	Val	Ile	His	Ile	Cys	Tyr	
						230					235					240	
40	aca	aac	ttg	aaa	gct	tca	act	tca	gct	gaa	gtt	gga	ctc	ttc	att	aag	768
	Thr	Asn	Leu	Lys	Ala	Ser	Thr	Ser	Ala	Glu	Val	Gly	Leu	Phe	Ile	Lys	
					245					250					255		
45	cag	ctt	cta	gaa	acc	tct	cca	gat	att	ctg	agg	gaa	tat	cta	att	cct	816
	Gln	Leu	Leu	Glu	Thr	Ser	Pro	Asp	Ile	Leu	Arg	Glu	Tyr	Leu	Ile	Pro	
				260					265					270			
50	ctg	caa	gag	cac	atg	gta	act	gtt	att	acc	cct	agc	act	tca	ggg	gct	864
	Leu	Gln	Glu	His	Met	Val	Thr	Val	Ile	Thr	Pro	Ser	Thr	Ser	Gly	Ala	
			275					280					285				
55	cga	aac	att	cat	gtc	atg	atg	gaa	ttc	cta	tta	ctt	att	ctt	tct	gat	912
	Arg	Asn	Ile	His	Val	Met	Met	Glu	Phe	Leu	Leu	Leu	Ile	Leu	Ser	Asp	
						295						300					
60	atg	ccc	aag	gac	ttt	att	cat	cat	gac	aaa	ctt	ttt	gat	ctc	ttg	gat	960
	Met	Pro	Lys	Asp	Phe	Ile	His	His	Asp	Lys	Leu	Phe	Asp	Leu	Leu	Asp	
						310					315					320	
65	cgt	gtc	gga	gta	ctt	acc	agg	gag	gta	tca	act	ctt	gta	cgt	gac	ttg	1008
	Arg	Val	Gly	Val	Leu	Thr	Arg	Glu	Val	Ser	Thr	Leu	Val	Arg	Asp	Leu	
					325					330					335		
70	gaa	gag	gaa	cca	agg	aat	aaa	gag	ggg	aat	aac	caa	aca	aat	tgt	gca	1056
	Glu	Glu	Glu	Pro	Arg	Asn	Lys	Glu	Gly	Asn	Asn	Gln	Thr	Asn	Cys	Ala	
				340					345					350			
75	acc	cta	gac	ttg	ctg	gaa	aat	att	gaa	ctc	ctc	aag	aaa	gat	ctc	aaa	1104
	Thr	Leu	Asp	Leu	Leu	Glu	Asn	Ile	Glu	Leu	Leu	Lys	Lys	Asp	Leu	Lys	
								360					365				
80	cat	gtt	tat	ctg	aaa	gcc	ctg	gat	tca	tct	caa	tgt	tgc	ttc	ccc	atg	1152
	His	Val	Tyr	Leu	Lys	Ala	Leu	Asp	Ser	Ser	Gln	Cys	Cys	Phe	Pro	Met	
							375					380					
85	agt	gat	gga	cca	ctc	ttc	atg	cat	ctt	cta	cac	ata	cac	tta	aat	gat	1200
	Ser	Asp	Gly	Pro	Leu	Phe	Met	His	Leu	Leu	His	Ile	His	Leu	Asn	Asp	
						390					395					400	
90	ttg	tta	gat	tct	aat	gct	tat	tca	att	gct	ttg	ata	aag	gaa	gaa	atc	1248

Agrico B.V.

20030596

PF 54801

29

	Leu	Leu	Asp	Ser	Asn	Ala	Tyr	Ser	Ile	Ala	Leu	Ile	Lys	Glu	Glu	Ile	
					405					410						415	
5	gag	ctg	gtg	aag	caa	gac	ctg	aaa	ttc	ata	aga	tca	ttc	ttt	gtg	gat	1296
	Glu	Leu	Val	Lys	Gln	Asp	Leu	Lys	Phe	Ile	Arg	Ser	Phe	Phe	Val	Asp	
				420					425					430			
10	gct	gag	caa	gga	ttg	tat	aaa	gat	ctc	tgg	gca	cgt	gtr	cta	gat	gtg	1344
	Ala	Glu	Gln	Gly	Leu	Tyr	Lys	Asp	Leu	Trp	Ala	Arg	Val	Leu	Asp	Val	
				435				440					445				
15	gct	tat	gag	gca	aaa	gat	gtc	ata	gat	tca	att	att	gtr	cga	gat	aat	1392
	Ala	Tyr	Glu	Ala	Lys	Asp	Val	Ile	Asp	Ser	Ile	Ile	Val	Arg	Asp	Asn	
				450			455					460					
20	ggt	ctc	tta	cat	ctt	att	ttc	tca	ctt	ccc	att	acc	ata	aag	aag	atc	1440
	Gly	Leu	Leu	His	Leu	Ile	Phe	Ser	Leu	Pro	Ile	Thr	Ile	Lys	Lys	Ile	
						470				475						480	
25	aaa	ctt	atc	aaa	gaa	gag	atc	tct	gct	tta	gat	gag	aac	att	ccc	aag	1488
	Lys	Leu	Ile	Lys	Glu	Glu	Ile	Ser	Ala	Leu	Asp	Glu	Asn	Ile	Pro	Lys	
					485					490					495		
30	gac	aga	ggt	cta	atc	gtt	gtg	aac	tct	ccc	aag	aaa	cca	gtr	gag	aga	1536
	Asp	Arg	Gly	Leu	Ile	Val	Val	Asn	Ser	Pro	Lys	Lys	Pro	Val	Glu	Arg	
				500					505					510			
35	aag	tca	ttg	aca	act	gat	aaa	ata	act	gta	ggt	ttt	gag	gag	gaa	aca	1584
	Lys	Ser	Leu	Thr	Thr	Asp	Lys	Ile	Thr	Val	Gly	Phe	Glu	Glu	Glu	Thr	
				515				520					525				
40	aac	ttg	ata	ctt	aga	aag	ctc	acc	agt	gga	tgc	gca	gat	cta	gat	gtc	1632
	Asn	Leu	Ile	Leu	Arg	Lys	Leu	Thr	Ser	Gly	Ser	Ala	Asp	Leu	Asp	Val	
				530			535					540					
45	att	tgc	atc	act	ggt	atg	ccg	ggt	tca	ggt	aaa	act	act	ttg	gca	tac	1680
	Ile	Ser	Ile	Thr	Gly	Met	Pro	Gly	Ser	Gly	Lys	Thr	Thr	Leu	Ala	Tyr	
						550					555					560	
50	aaa	gta	tac	aat	gat	aag	tca	gtt	tct	agc	cgt	ttc	gac	ctt	cgt	gca	1728
	Lys	Val	Tyr	Asn	Asp	Lys	Ser	Val	Ser	Ser	Arg	Phe	Asp	Leu	Arg	Ala	
					565				570						575		
55	tgg	tgc	acg	gtc	gac	caa	gga	tgt	gat	gag	aag	aag	ttg	ttg	aat	aca	1776
	Trp	Cys	Thr	Val	Asp	Gln	Gly	Cys	Asp	Glu	Lys	Lys	Leu	Leu	Asn	Thr	
				580					585					590			
60	att	ttc	agt	caa	gtt	agt	gac	tca	gat	tca	aaa	ttg	agt	gag	aat	att	1824
	Ile	Phe	Ser	Gln	Val	Ser	Asp	Ser	Asp	Ser	Lys	Leu	Ser	Glu	Asn	Ile	
				595				600					605				
65	gat	gtt	gct	gat	aaa	tta	cgg	aaa	caa	ctg	ttt	gga	aag	agg	tat	ctt	1872
	Asp	Val	Ala	Asp	Lys	Leu	Arg	Lys	Gln	Leu	Phe	Gly	Lys	Arg	Tyr	Leu	
				610			615					620					
70	att	gtc	tta	gat	gac	gtg	tgg	gat	act	act	aca	tgg	gat	gag	tta	aca	1920
	Ile	Val	Leu	Asp	Asp	Val	Trp	Asp	Thr	Thr	Thr	Trp	Asp	Glu	Leu	Thr	
						630					635					640	
75	aga	cct	ttt	cct	gaa	tct	aag	aaa	gga	agt	agg	att	att	ttg	aca	act	1968
	Arg	Pro	Phe	Pro	Glu	Ser	Lys	Lys	Gly	Ser	Arg	Ile	Ile	Leu	Thr	Thr	
					645				650						655		
80	cgg	gaa	aag	gaa	gtg	gct	ttg	cat	gga	aag	ctg	aac	act	gat	cct	ctt	2016
	Arg	Glu	Lys	Glu	Val	Ala	Leu	His	Gly	Lys	Leu	Asn	Thr	Asp	Pro	Leu	
				660					665					670			
85	gac	ctt	cga	ttg	cta	aga	cca	gat	gaa	agt	tgg	gaa	cta	tta	gag	aaa	2064
	Asp	Leu	Arg	Leu	Leu	Arg	Pro	Asp	Glu	Ser	Trp	Glu	Leu	Leu	Glu	Lys	
				675				680					685				
90	agg	gca	ttt	ggg	aat	gag	agt	tgc	cct	gat	gaa	cta	tta	gat	gtc	ggt	2112
	Arg	Ala	Phe	Gly	Asn	Glu	Ser	Cys	Pro	Asp	Glu	Leu	Leu	Asp	Val	Gly	

Agrico B.V.

20030596

PF 54801

30

	690	695	700	
5	aaa gaa ata gcc gaa aat tgt aaa ggg ctt cct ttg gtg gct gat ctg Lys Glu Ile Ala Glu Asn Cys Lys Gly Leu Pro Leu Val Ala Asp Leu 705 710 715 720			2160
10	att gct gga gtc att gct ggg agg gaa aag aaa agg agt gtg tgg ctt Ile Ala Gly Val Ile Ala Gly Arg Glu Lys Lys Arg Ser Val Trp Leu 725 730 735			2208
15	gaa gtt caa agt agt ttg agt tct ttt att ttg aac agt gaa gtg gaa Glu Val Gln Ser Ser Leu Ser Ser Phe Ile Leu Asn Ser Glu Val Glu 740 745 750			2256
20	gtg atg aaa gtt ata gaa tta agt tat gac cat tta cca cat cac ctc Val Met Lys Val Ile Glu Leu Ser Tyr Asp His Leu Pro His His Leu 755 760 765			2304
25	aag cca tgc ttg ctg tat ttt gca agt ttt ccg aag gac act tca ttg Lys Pro Cys Leu Leu Tyr Phe Ala Ser Phe Pro Lys Asp Thr Ser Leu 770 775 780			2352
30	aca atc tat gag ttg aat gtt tat ttc ggt gct gaa gga ttt gtg gga Thr Ile Tyr Glu Leu Asn Val Tyr Phe Gly Ala Glu Gly Phe Val Gly 785 790 795 800			2400
35	aag acg gag atg aac agt atg gaa gaa gtg gtg aag att tat atg gat Lys Thr Glu Met Asn Ser Met Glu Glu Val Val Lys Ile Tyr Met Asp 805 810 815			2448
40	gat tta att tac agt agc ttg gta att tgt ttc aat gag ata ggt tat Asp Leu Ile Tyr Ser Ser Leu Val Ile Cys Phe Asn Glu Ile Gly Tyr 820 825 830			2496
45	gca ctg aat ttc caa att cat gat ctt gtg cat gac ttt tgt ttg ata Ala Leu Asn Phe Gln Ile His Asp Leu Val His Asp Phe Cys Leu Ile 835 840 845			2544
50	aaa gca aga aag gaa aat ttg ttt gat cag ata aga tca agt gct cca Lys Ala Arg Lys Glu Asn Leu Phe Asp Gln Ile Arg Ser Ser Ala Pro 850 855 860			2592
55	tca gat ttg ttg cct cgt caa att acc att gat tgt gat gag gag gag Ser Asp Leu Leu Pro Arg Gln Ile Thr Ile Asp Cys Asp Glu Glu Glu 865 870 875 880			2640
60	cac ttt ggg ctt aat ttt gtc atg ttc gat tca aat aag aaa agg cat His Phe Gly Leu Asn Phe Val Met Phe Asp Ser Asn Lys Lys Arg His 885 890 895			2688
65	tct ggt aaa cac ctc tat tct ttg agg ata att gga gac cag ctg gat Ser Gly Lys His Leu Tyr Ser Leu Arg Ile Ile Gly Asp Gln Leu Asp 900 905 910			2736
70	gac agt gct tct gat gca ttt cac cta aga cac ttg agg ctt ctt aga Asp Ser Val Ser Asp Ala Phe His Leu Arg His Leu Arg Leu Leu Arg 915 920 925			2784
75	gtg ttg gac ctg cat acg tct ttt atc atg gtg aaa gat tct ttg ctg Val Leu Asp Leu His Thr Ser Phe Ile Met Val Lys Asp Ser Leu Leu 930 935 940			2832
80	aat gaa ata tgc atg ttg aat cat ttg agg tac tta tcc att gac aca Asn Glu Ile Cys Met Leu Asn His Leu Arg Tyr Leu Ser Ile Asp Thr 945 950 955 960			2880
85	caa gtt aaa tat ctg cct ttg tct ttc tca aac ctc tgg aat cta gaa Gln Val Lys Tyr Leu Pro Leu Ser Phe Ser Asn Leu Trp Asn Leu Glu 965 970 975			2928
90	agc ctg ttt gtg tct acc aac aga tca atc ttg gta cta tta ccg aga Ser Leu Phe Val Ser Thr Asn Arg Ser Ile Leu Val Leu Leu Pro Arg 980 985 990			2976

PF 54801

31

	att	ttg	gat	ctt	gta	aag	ttg	cga	gtg	ctg	tcc	gtg	gat	gct	tgt	tct	3024
	Ile	Leu	Asp	Leu	Val	Lys	Leu	Arg	Val	Leu	Ser	Val	Asp	Ala	Cys	Ser	
			995					1000					1005				
5	ttc	ttt	gat	atg	gat	gca	gat		gaa	tca	ata	ttg	ata	gca	gag	gac	3069
	Phe	Phe	Asp	Met	Asp	Ala	Asp		Glu	Ser	Ile	Leu	Ile	Ala	Glu	Asp	
		1010					1015						1020				
10	aca	aag	tta	gag	aac	ttg	aga		ata	tta	atg	gaa	ctg	ttg	att	tcc	3114
	Thr	Lys	Leu	Glu	Asn	Leu	Arg		Ile	Leu	Thr	Glu	Leu	Leu	Ile	Ser	
		1025					1030						1035				
15	tat	tcg	aaa	gat	aca	aag	aat		att	ttc	aaa	agg	ttt	ccc	aat	ctt	3159
	Tyr	Ser	Lys	Asp	Thr	Lys	Asn		Ile	Phe	Lys	Arg	Phe	Pro	Asn	Leu	
		1040					1045						1050				
20	cag	ttg	ctt	tca	ttt	gaa	ctc		aag	gag	tca	tgg	gat	tat	tca	aca	3204
	Gln	Leu	Leu	Ser	Phe	Glu	Leu		Lys	Glu	Ser	Trp	Asp	Tyr	Ser	Thr	
		1055					1060						1065				
25	gag	caa	cat	tgg	ttc	tcg	gaa		ttg	gat	ttc	cta	act	gaa	cta	gaa	3249
	Glu	Gln	His	Trp	Phe	Ser	Glu		Leu	Asp	Phe	Leu	Thr	Glu	Leu	Glu	
		1070					1075						1080				
30	aca	ctc	tct	gta	ggt	ttt	aaa		agt	tca	aac	aca	aac	gat	agt	ggg	3294
	Thr	Leu	Ser	Val	Gly	Phe	Lys		Ser	Ser	Asn	Thr	Asn	Asp	Ser	Gly	
		1085					1090						1095				
35	tcc	tct	gta	gcg	aca	aat	cgg		cgg	tgg	gat	ttt	caa	ttc	cct	tca	3339
	Ser	Ser	Val	Ala	Thr	Asn	Arg		Pro	Trp	Asp	Phe	His	Phe	Pro	Ser	
		1100					1105						1110				
40	aat	ttg	aaa	ata	ctg	tgg	ttg		cgt	gaa	ttt	ccg	ctg	aca	tcc	gat	3384
	Asn	Leu	Lys	Ile	Leu	Trp	Leu		Arg	Glu	Phe	Pro	Leu	Thr	Ser	Asp	
		1115					1120						1125				
45	tca	cta	tca	aca	ata	gcg	aga		ctg	ccc	aac	ctt	gaa	gag	ttg	tcc	3429
	Ser	Leu	Ser	Thr	Ile	Ala	Arg		Leu	Pro	Asn	Leu	Glu	Glu	Leu	Ser	
		1130					1135						1140				
50	ctt	tat	cat	aca	atc	atc	cat		gga	gaa	gaa	tgg	aac	atg	ggg	gag	3474
	Leu	Tyr	His	Thr	Ile	Ile	His		Gly	Glu	Glu	Trp	Asn	Met	Gly	Glu	
		1145					1150						1155				
55	gaa	gac	acc	ttt	gag	aat	ctc		aaa	ttt	ttg	aac	tto	aat	caa	gtt	3519
	Glu	Asp	Thr	Phe	Glu	Asn	Leu		Lys	Phe	Leu	Asn	Phe	Asn	Gln	Val	
		1160					1165						1170				
60	agt	att	tcc	aag	tgg	gag	gtt		gga	gag	gaa	tcc	ttc	ccc	aat	ctt	3564
	Ser	Ile</															

Agrico B.V.

20030596

PF 54801

32

<210> 8
 <211> 1255
 5 <212> PRT
 <213> Lycopersicon lycopersicum
 10
 <220>
 <221> misc_feature
 15 <222> (178)..(178)
 <223> The 'Xaa' at location 178 stands for Leu.
 20 <400> 8
 Met Glu Lys Arg Lys Asp Asn Glu Glu Ala Asn Asn Ser Leu Val Leu
 1 5 10 15
 25 Phe Ser Ala Leu Ser Lys Asp Ile Ala Asp Val Leu Val Phe Leu Glu
 20 25 30
 30 Asn Glu Glu Asn Gln Lys Ala Leu Asp Lys Asp Gln Val Glu Lys Ile
 35 40 45
 35 Lys Leu Lys Met Ala Phe Ile Cys Thr Tyr Val Gln Leu Ser Cys Ser
 50 55 60
 40 Asp Phe Glu Gln Phe Glu Asp Ile Met Thr Arg Lys Arg Gln Glu Val
 65 70 75 80
 Glu Asn Leu Leu Gln Pro Leu Leu Asp Asp Val Phe Thr Ser Leu
 85 90 95
 45 Thr Ser Asn Met Asp Asp Cys Ile Ser Leu Tyr His Arg Ser Tyr Lys
 100 105 110
 50 Ser Asp Ala Ile Met Met Asp Glu Gln Leu Asp Phe Leu Leu Leu Asn
 115 120 125
 55 Leu Tyr His Leu Ser Lys His His Ala Glu Lys Ile Phe Pro Gly Val
 130 135 140
 Thr Gln Tyr Glu Val Leu Gln Asn Ile Cys Gly Asn Ile Arg Asp Phe
 145 150 155 160
 60 His Gly Leu Ile Val Asn Gly Cys Ile Lys His Glu Met Val Glu Asn
 165 170 175
 65 Val Xaa Pro Leu Phe Gln Leu Met Ala Asp Arg Val Gly His Phe Leu
 180 185 190
 70 Trp Asp Asp Gln Thr Asp Glu Asp Ser Arg Leu Ser Glu Leu Asp Glu
 195 200 205

Agrico B.V.

20030596

PF 54801

33

	Asp	Glu	Gln	Asn	Asp	Arg	Asp	Ser	Arg	Leu	Phe	Lys	Leu	Ala	His	Leu	
	210						215					220					
5	Leu	Leu	Lys	Ile	Val	Pro	Val	Glu	Leu	Glu	Val	Ile	His	Ile	Cys	Tyr	
	225					230					235					240	
10	Thr	Asn	Leu	Lys	Ala	Ser	Thr	Ser	Ala	Glu	Val	Gly	Leu	Phe	Ile	Lys	
					245					250					255		
15	Gln	Leu	Leu	Glu	Thr	Ser	Pro	Asp	Ile	Leu	Arg	Glu	Tyr	Leu	Ile	Pro	
				260					265					270			
20	Leu	Gln	Glu	His	Met	Val	Thr	Val	Ile	Thr	Pro	Ser	Thr	Ser	Gly	Ala	
			275					280					285				
25	Arg	Asn	Ile	His	Val	Met	Met	Glu	Phe	Leu	Leu	Leu	Ile	Leu	Ser	Asp	
	290						295					300					
30	Met	Pro	Lys	Asp	Phe	Ile	His	His	Asp	Lys	Leu	Phe	Asp	Leu	Leu	Asp	
	305					310					315					320	
35	Arg	Val	Gly	Val	Leu	Thr	Arg	Glu	Val	Ser	Thr	Leu	Val	Arg	Asp	Leu	
					325					330					335		
40	Glu	Glu	Glu	Pro	Arg	Asn	Lys	Glu	Gly	Asn	Asn	Gln	Thr	Asn	Cys	Ala	
				340					345					350			
45	Thr	Leu	Asp	Leu	Leu	Glu	Asn	Ile	Glu	Leu	Leu	Lys	Lys	Asp	Leu	Lys	
			355					360					365				
50	His	Val	Tyr	Leu	Lys	Ala	Leu	Asp	Ser	Ser	Gln	Cys	Cys	Phe	Pro	Met	
		370					375					380					
55	Ser	Asp	Gly	Pro	Leu	Phe	Met	His	Leu	Leu	His	Ile	His	Leu	Asn	Asp	
	385					390					395					400	
60	Leu	Leu	Asp	Ser	Asn	Ala	Tyr	Ser	Ile	Ala	Leu	Ile	Lys	Glu	Glu	Ile	
					405					410					415		
65	Glu	Leu	Val	Lys	Gln	Asp	Leu	Lys	Phe	Ile	Arg	Ser	Phe	Phe	Val	Asp	
				420					425					430			
70	Ala	Glu	Gln	Gly	Leu	Tyr	Lys	Asp	Leu	Trp	Ala	Arg	Val	Leu	Asp	Val	
			435					440					445				
75	Ala	Tyr	Glu	Ala	Lys	Asp	Val	Ile	Asp	Ser	Ile	Ile	Val	Arg	Asp	Asn	
		450					455					460					
80	Gly	Leu	Leu	His	Leu	Ile	Phe	Ser	Leu	Pro	Ile	Thr	Ile	Lys	Lys	Ile	
	465					470					475					480	
85	Lys	Leu	Ile	Lys	Glu	Glu	Ile	Ser	Ala	Leu	Asp	Glu	Asn	Ile	Pro	Lys	
					485					490					495		
90	Asp	Arg	Gly	Leu	Ile	Val	Val	Asn	Ser	Pro	Lys	Lys	Pro	Val	Glu	Arg	

Agrico B.V.

20030596

PF 54801

34

500

505

510

5 Lys Ser Leu Thr Thr Asp Lys Ile Thr Val Gly Phe Glu Glu Glu Thr
 515 520 525

10 Asn Leu Ile Leu Arg Lys Leu Thr Ser Gly Ser Ala Asp Leu Asp Val
 530 535 540

15 Ile Ser Ile Thr Gly Met Pro Gly Ser Gly Lys Thr Thr Leu Ala Tyr
 545 550 555 560

20 Lys Val Tyr Asn Asp Lys Ser Val Ser Ser Arg Phe Asp Leu Arg Ala
 565 570 575

25 Trp Cys Thr Val Asp Gln Gly Cys Asp Glu Lys Lys Leu Leu Asn Thr
 580 585 590

30 Ile Phe Ser Gln Val Ser Asp Ser Asp Ser Lys Leu Ser Glu Asn Ile
 595 600 605

35 Asp Val Ala Asp Lys Leu Arg Lys Gln Leu Phe Gly Lys Arg Tyr Leu
 610 615 620

40 Ile Val Leu Asp Asp Val Trp Asp Thr Thr Thr Trp Asp Glu Leu Thr
 625 630 635 640

45 Arg Pro Phe Pro Glu Ser Lys Lys Gly Ser Arg Ile Ile Leu Thr Thr
 645 650 655

50 Arg Glu Lys Glu Val Ala Leu His Gly Lys Leu Asn Thr Asp Pro Leu
 660 665 670

55 Asp Leu Arg Leu Leu Arg Pro Asp Glu Ser Trp Glu Leu Leu Glu Lys
 675 680 685

60 Arg Ala Phe Gly Asn Glu Ser Cys Pro Asp Glu Leu Leu Asp Val Gly
 690 695 700

65 Lys Glu Ile Ala Glu Asn Cys Lys Gly Leu Pro Leu Val Ala Asp Leu
 705 710 715 720

70 Ile Ala Gly Val Ile Ala Gly Arg Glu Lys Lys Arg Ser Val Trp Leu
 725 730 735

75 Glu Val Gln Ser Ser Leu Ser Ser Phe Ile Leu Asn Ser Glu Val Glu
 740 745 750

80 Val Met Lys Val Ile Glu Leu Ser Tyr Asp His Leu Pro His His Leu
 755 760 765

85 Lys Pro Cys Leu Leu Tyr Phe Ala Ser Phe Pro Lys Asp Thr Ser Leu
 770 775 780

90 Thr Ile Tyr Glu Leu Asn Val Tyr Phe Gly Ala Glu Gly Phe Val Gly
 785 790 795 800

Agrico B.V.

20030596

PF 54801

35

5 Lys Thr Glu Met Asn Ser Met Glu Glu Val Val Lys Ile Tyr Met Asp
 805 810 815
 Asp Leu Ile Tyr Ser Ser Leu Val Ile Cys Phe Asn Glu Ile Gly Tyr
 820 825 830
 10 Ala Leu Asn Phe Gln Ile His Asp Leu Val His Asp Phe Cys Leu Ile
 835 840 845
 15 Lys Ala Arg Lys Glu Asn Leu Phe Asp Gln Ile Arg Ser Ser Ala Pro
 850 855 860
 20 Ser Asp Leu Leu Pro Arg Gln Ile Thr Ile Asp Cys Asp Glu Glu Glu
 865 870 875 880
 His Phe Gly Leu Asn Phe Val Met Phe Asp Ser Asn Lys Lys Arg His
 885 890 895
 25 Ser Gly Lys His Leu Tyr Ser Leu Arg Ile Ile Gly Asp Gln Leu Asp
 900 905 910
 30 Asp Ser Val Ser Asp Ala Phe His Leu Arg His Leu Arg Leu Leu Arg
 915 920 925
 35 Val Leu Asp Leu His Thr Ser Phe Ile Met Val Lys Asp Ser Leu Leu
 930 935 940
 40 Asn Glu Ile Cys Met Leu Asn His Leu Arg Tyr Leu Ser Ile Asp Thr
 945 950 955 960
 Gln Val Lys Tyr Leu Pro Leu Ser Phe Ser Asn Leu Trp Asn Leu Glu
 965 970 975
 45 Ser Leu Phe Val Ser Thr Asn Arg Ser Ile Leu Val Leu Leu Pro Arg
 980 985 990
 50 Ile Leu Asp Leu Val Lys Leu Arg Val Leu Ser Val Asp Ala Cys Ser
 995 1000 1005
 55 Phe Phe Asp Met Asp Ala Asp Glu Ser Ile Leu Ile Ala Glu Asp
 1010 1015 1020
 60 Thr Lys Leu Glu Asn Leu Arg Ile Leu Thr Glu Leu Leu Ile Ser
 1025 1030 1035
 Tyr Ser Lys Asp Thr Lys Asn Ile Phe Lys Arg Phe Pro Asn Leu
 1040 1045 1050
 65 Gln Leu Leu Ser Phe Glu Leu Lys Glu Ser Trp Asp Tyr Ser Thr
 1055 1060 1065
 70 Glu Gln His Trp Phe Ser Glu Leu Asp Phe Leu Thr Glu Leu Glu
 1070 1075 1080

Agrico B.V.

20030596

PF 54801

36

Thr Leu Ser Val Gly Phe Lys Ser Ser Asn Thr Asn Asp Ser Gly
 1085 1090 1095
 5 Ser Ser Val Ala Thr Asn Arg Pro Trp Asp Phe His Phe Pro Ser
 1100 1105 1110
 10 Asn Leu Lys Ile Leu Trp Leu Arg Glu Phe Pro Leu Thr Ser Asp
 1115 1120 1125
 15 Ser Leu Ser Thr Ile Ala Arg Leu Pro Asn Leu Glu Glu Leu Ser
 1130 1135 1140
 20 Leu Tyr His Thr Ile Ile His Gly Glu Glu Trp Asn Met Gly Glu
 1145 1150 1155
 Glu Asp Thr Phe Glu Asn Leu Lys Phe Leu Asn Phe Asn Gln Val
 1160 1165 1170
 25 Ser Ile Ser Lys Trp Glu Val Gly Glu Glu Ser Phe Pro Asn Leu
 1175 1180 1185
 30 Glu Lys Leu Lys Leu Arg Gly Cys His Lys Leu Glu Glu Ile Pro
 1190 1195 1200
 35 Pro Ser Phe Gly Asp Ile Tyr Ser Leu Lys Ser Ile Lys Ile Val
 1205 1210 1215
 40 Lys Ser Pro Gln Leu Glu Asp Ser Ala Leu Lys Ile Lys Glu Tyr
 1220 1225 1230
 Ala Glu Asp Met Arg Gly Gly Asp Glu Leu Gln Ile Leu Gly Gln
 1235 1240 1245
 45 Lys Asn Ile Pro Leu Phe Lys
 1250 1255
 50 <210> 9
 <211> 3774
 <212> DNA
 55 <213> Lycopersicon lycopersicum
 60 <220>
 <221> CDS
 <222> (1)..(3774)
 65 <223> Mil.2 from tomato
 70 <400> 9
 atg gaa aaa cga aaa gat att gaa gaa gca aac aac tca ttcg gtg tta
 Met Glu Lys Arg Lys Asp Ile Glu Glu Ala Asn Asn Ser Leu Val Leu
 1 5 10 15

Agrico B.V.

20030596

PF 54801

37

	ttt tct gct ctt agc aag gac att gcc aat gtt cta att ttc cta gag	96
	Phe Ser Ala Leu Ser Lys Asp Ile Ala Asn Val Leu Ile Phe Leu Glu	
	20 25 30	
5	aat gag gaa aat caa aaa gct ctt gac aaa gat caa gtt gaa aag cta	144
	Asn Glu Glu Asn Gln Lys Ala Leu Asp Lys Asp Gln Val Glu Lys Leu	
	35 40 45	
10	aaa ttg aaa atg gca ttt att tgt aca tat gtt cag ctt tct tat tcc	192
	Lys Leu Lys Met Ala Phe Ile Cys Thr Tyr Val Gln Leu Ser Tyr Ser	
	50 55 60	
15	gat ttt gag cag ttt gaa gat ata atg act aga aat aga caa gag gtt	240
	Asp Phe Glu Gln Phe Glu Asp Ile Met Thr Arg Asn Arg Gln Glu Val	
	65 70 75 80	
20	gag aat ctg ctt caa tca ctt ttg gat gat gat gtc ctt act agc ctc	288
	Glu Asn Leu Leu Gln Ser Leu Leu Asp Asp Asp Val Leu Thr Ser Leu	
	85 90 95	
	acc agt aat atg gat gac tgt atc age ttg tat cat cgt tct tat aaa	336
	Thr Ser Asn Met Asp Asp Cys Ile Ser Leu Tyr His Arg Ser Tyr Lys	
	100 105 110	
25	tca gat gcc atc atg atg gat gag caa ttg gac ttc ctc ctc ttg aat	384
	Ser Asp Ala Ile Met Met Asp Glu Gln Leu Asp Phe Leu Leu Leu Asn	
	115 120 125	
30	ctg tat cat cta tcc aag cat cac gct gaa aag ata ttt cct gga gtg	432
	Leu Tyr His Leu Ser Lys His His Ala Glu Lys Ile Phe Pro Gly Val	
	130 135 140	
35	act caa tat gaa gtt ctt cag aat gta tgt ggc aac ata aga gat ttc	480
	Thr Gln Tyr Glu Val Leu Gln Asn Val Cys Gly Asn Ile Arg Asp Phe	
	145 150 155 160	
40	cat ggg ttg ata ctg aat ggt tgc att aag cat gag atg gtt gag aat	528
	His Gly Leu Ile Leu Asn Gly Cys Ile Lys His Glu Met Val Glu Asn	
	165 170 175	
	gtc tta cct ctg ttt caa ctc atg gct gaa aga gta gga cac ttc ctt	576
	Val Leu Pro Leu Phe Gln Leu Met Ala Glu Arg Val Gly His Phe Leu	
	180 185 190	
45	tgg gag gat cag act gat gaa gac tct cgg ctc tcc gag cta gat gag	624
	Trp Glu Asp Gln Thr Asp Glu Asp Ser Arg Leu Ser Glu Leu Asp Glu	
	195 200 205	
50	gat gaa cac aat gat aga gac tct cga ctc ttc cag cta aca cat cta	672
	Asp Glu His Asn Asp Arg Asp Ser Arg Leu Phe Gln Leu Thr His Leu	
	210 215 220	
55	ctc ttg aag att gtt cca act gaa ctg gag gtt atg cac ata tgt tat	720
	Leu Leu Lys Ile Val Pro Thr Glu Leu Glu Val Met His Ile Cys Tyr	
	225 230 235 240	
60	aca aat ttg aaa gct tca act tca gca gaa gtt gga cgc ttc att aag	768
	Thr Asn Leu Lys Ala Ser Thr Ser Ala Glu Val Gly Arg Phe Ile Lys	
	245 250 255	
	aag ctc ctg gaa acc tca ccg gat att ctc aga gaa tat atc att caa	816
	Lys Leu Leu Glu Thr Ser Pro Asp Ile Leu Arg Glu Tyr Ile Ile Gln	
	260 265 270	
65	cta caa gag cat atg tta act gtt att ccc cct agc act tta ggg gct	864
	Leu Gln Glu His Met Leu Thr Val Ile Pro Pro Ser Thr Leu Gly Ala	
	275 280 285	
70	cga aac att cat gtc atg atg gaa ttc cta tta ctt att ctt tct gat	912
	Arg Asn Ile His Val Met Met Glu Phe Leu Leu Leu Ile Leu Ser Asp	
	290 295 300	

Agrico B.V.

20030596

PF 54801

38

	atg ccc aag gac ttt att cat cat gac aaa ctt ttt gat ctc ttg gct	960
	Met Pro Lys Asp Phe Ile His His Asp Lys Leu Phe Asp Leu Leu Ala	
	305 310 315 320	
5	cat gtc gga aca ctt acc agg gag gta tgg act ctt gta cgt gac ttg	1008
	His Val Gly Thr Leu Thr Arg Glu Val Ser Thr Leu Val Arg Asp Leu	
	325 330 335	
10	gaa gag aaa tta agg aat aaa gag ggt aat aac caa aca aat tgt gca	1056
	Glu Glu Lys Leu Arg Asn Lys Glu Gly Asn Asn Gln Thr Asn Cys Ala	
	340 345 350	
15	acc cta gac ttg ctg gaa aat att gaa ctc ctc aag aaa gat ctc aaa	1104
	Thr Leu Asp Leu Leu Glu Asn Ile Glu Leu Leu Lys Lys Asp Leu Lys	
	355 360 365	
20	cat gtt tat ctg aaa gcc cca aat tca tct caa tgt tgo tta ccc atg	1152
	His Val Tyr Leu Lys Ala Pro Asn Ser Ser Gln Cys Cys Phe Pro Met	
	370 375 380	
25	agt gat gga cca ctc ttc atg cat ctt cta cac atg cac tta aat gat	1200
	Ser Asp Gly Pro Leu Phe Met His Leu Leu His Met His Leu Asn Asp	
	385 390 395 400	
30	ttg cta gat tct aat gct tat tca att tct ttg ata aag gaa gaa atc	1248
	Leu Leu Asp Ser Asn Ala Tyr Ser Ile Ser Leu Ile Lys Glu Glu Ile	
	405 410 415	
35	gag ttg gtg agt caa gaa ctg gaa ttc ata aga tca ttc ttt ggg gat	1296
	Glu Leu Val Ser Gln Glu Leu Glu Phe Ile Arg Ser Phe Phe Gly Asp	
	420 425 430	
40	gct gct gag caa gga ttg tat aaa gat atc tgg gca cgt gtt cta gat	1344
	Ala Ala Glu Gln Gly Leu Tyr Lys Asp Ile Trp Ala Arg Val Leu Asp	
	435 440 445	
45	gtg gct tat gag gca aaa gat gtc ata gat tca att att gtt cga gat	1392
	Val Ala Tyr Glu Ala Lys Asp Val Ile Asp Ser Ile Ile Val Arg Asp	
	450 455 460	
50	aac ggt ctc tta cat ctt att ttc tca ctt ccc att acc ata aag aag	1440
	Asn Gly Leu Leu His Leu Ile Phe Ser Leu Pro Ile Thr Ile Lys Lys	
	465 470 475 480	
55	atc aaa ctt atc aaa gaa gag atc tct gct tta gat gag aac att ccc	1488
	Ile Lys Leu Ile Lys Glu Glu Ile Ser Ala Leu Asp Glu Asn Ile Pro	
	485 490 495	
60	aag gac aga ggt cta atc gtt gtg aac tct ccc aag aaa cca gtt gag	1536
	Lys Asp Arg Gly Leu Ile Val Val Asn Ser Pro Lys Lys Pro Val Glu	
	500 505 510	
65	aga aag tca ttg aca act gat aaa ata att gta ggt ttt gag gag gag	1584
	Arg Lys Ser Leu Thr Thr Asp Lys Ile Ile Val Gly Phe Glu Glu Glu	
	515 520 525	
70	aca aac ttg ata ctt aga aag ctc acc agt gga ccc gca gat tta gat	1632
	Thr Asn Leu Ile Leu Arg Lys Leu Thr Ser Gly Pro Ala Asp Leu Asp	
	530 535 540	
75	gtc att tgg atc acc ggt atg ccg ggt tca ggt aaa act act ttg gca	1680
	Val Ile Ser Ile Thr Gly Met Pro Gly Ser Gly Lys Thr Thr Leu Ala	
	545 550 555 560	
80	tac aaa gta tac aat gat aag tca gtt tot aga cat ttt gac ctt cgt	1728
	Tyr Lys Val Tyr Asn Asp Lys Ser Val Ser Arg His Phe Asp Leu Arg	
	565 570 575	
85	gca tgg tgc acg gtc gat caa gga tat gac gac aag aag ttg ttg gat	1776
	Ala Trp Cys Thr Val Asp Gln Gly Tyr Asp Asp Lys Lys Leu Leu Asp	
	580 585 590	
90	aca att ttc agt caa gtt agt ggc tca gat tca aat ttg agt gag aat	1824

Agrico B.V.

20030596

PF 54801

39

	Thr	Ile	Phe	Ser	Gln	Val	Ser	Gly	Ser	Asp	Ser	Asn	Leu	Ser	Glu	Asn	
			595					600					605				
5	att	gat	gtt	gct	gat	aaa	ttg	cgg	aaa	caa	ctg	ttt	gga	aag	agg	tat	1872
	Ile	Asp	Val	Ala	Asp	Lys	Leu	Arg	Lys	Gln	Leu	Phe	Gly	Lys	Arg	Tyr	
	610						615					620					
10	ctt	att	gtc	cta	gat	gat	gtg	tgg	gat	act	act	aca	ttg	gat	gag	ttg	1920
	Leu	Ile	Val	Leu	Asp	Asp	Val	Trp	Asp	Thr	Thr	Thr	Leu	Asp	Glu	Leu	
	625					630					635				640		
15	aca	aga	cct	ttt	cot	gaa	gct	aag	aaa	gga	agt	agg	att	att	ttg	aca	1968
	Thr	Arg	Pro	Phe		Glu	Ala	Lys	Lys	Gly	Ser	Arg	Ile	Ile	Leu	Thr	
					645					650					655		
20	act	cga	gaa	aag	gaa	gtg	gct	ttg	cat	gga	aag	ctg	aac	act	gat	cct	2016
	Thr	Arg	Glu	Lys	Glu	Val	Ala	Leu	His	Gly	Lys	Leu	Asn	Thr	Asp	Pro	
					660				665					670			
25	ctt	gac	ctt	cga	ttg	cta	aga	cca	gat	gaa	agt	tgg	gaa	ctt	tta	gat	2064
	Leu	Asp	Leu	Arg	Leu	Leu	Arg	Pro	Asp	Glu	Ser	Trp	Glu	Leu	Leu	Asp	
			675					680					685				
30	aaa	agg	aca	ttt	ggt	aat	gag	agt	tgc	cct	gat	gaa	cta	tta	gat	gtc	2112
	Lys	Arg	Thr	Phe	Gly	Asn	Glu	Ser	Cys	Pro	Asp	Glu	Leu	Leu	Asp	Val	
							695					700					
35	ggt	aaa	gaa	ata	gcc	gaa	aat	tgt	aaa	ggg	ctt	cct	ctg	gtg	got	gat	2160
	Gly	Lys	Glu	Ile	Ala	Glu	Asn	Cys	Lys	Gly	Leu	Pro	Leu	Val	Ala	Asp	
	705					710					715				720		
40	ctg	att	got	gga	gtc	att	gct	ggg	agg	gaa	aag	aaa	agg	agt	gtg	tgg	2208
	Leu	Ile	Ala	Gly	Val	Ile	Ala	Gly	Arg	Glu	Lys	Lys	Arg	Ser	Val	Trp	
					725					730					735		
45	ctt	gaa	gtt	caa	agt	agt	ttg	agt	tct	ctt	att	ttg	aac	agt	gaa	gtg	2256
	Leu	Glu	Val	Gln	Ser	Ser	Leu	Ser	Ser	Phe	Ile	Leu	Asn	Ser	Glu	Val	
					740				745					750			
50	gaa	gtg	atg	aaa	gtt	ata	gaa	tta	agt	tat	gac	cat	tta	cca	cat	cac	2304
	Glu	Val	Met	Lys	Val	Ile	Glu	Leu	Ser	Tyr	Asp	His	Leu	Pro	His	His	
			755				760						765				
55	ctc	aag	cca	tgc	ttg	ctt	cac	ttt	gca	agt	tgg	cgg	aag	gac	act	cct	2352
	Leu	Lys	Pro	Cys	Leu	Leu	His	Phe	Ala	Ser	Trp	Pro	Lys	Asp	Thr	Pro	
			770				775					780					
60	ttg	aca	atc	tat	ttg	ttt	act	gtt	tat	ttg	ggt	gct	gaa	gga	ttt	gtg	2400
	Leu	Thr	Ile	Tyr	Leu	Phe	Thr	Val	Tyr	Leu	Gly	Ala	Glu	Gly	Phe	Val	
	785					790					795				800		
65	gaa	aag	acg	gag	atg	aag	ggt	ata	gaa	gaa	gtg	gtg	aag	att	tat	atg	2448
	Glu	Lys	Thr	Glu	Met	Lys	Gly	Ile	Glu	Glu	Val	Val	Lys	Ile	Tyr	Met	
					805				810						815		
70	gat	gat	cta	act	tcc	agt	agc	ttg	gta	att	tgt	ttc	aat	gag	ata	ggt	2496
	Asp	Asp	Leu	Ile	Ser	Ser	Ser	Leu	Val	Ile	Cys	Phe	Asn	Glu	Ile	Gly	
				820					825					830			
75	gat	ata	ctg	aat	ttc	caa	att	cat	gat	ctt	gtg	cat	gac	ttt	tgt	ttg	2544
	Asp	Ile	Leu	Asn	Phe	Gln	Ile	His	Asp	Leu	Val	His	Asp	Phe	Cys	Leu	
			835					840					845				
80	ata	aaa	gca	aga	aag	gaa	aat	ttg	ttt	gat	cgg	ata	aga	tca	agt	gct	2592
	Ile	Lys	Ala	Arg	Lys	Glu	Asn	Leu	Phe	Asp	Arg	Ile	Arg	Ser	Ser	Ala	
			850				855					860					
85	cca	tca	gat	ttg	ttg	cct	cgt	caa	att	acc	att	gat	tat	gat	gag	gag	2640
	Pro	Ser	Asp	Leu	Leu	Pro	Arg	Gln	Ile	Thr	Ile	Asp	Tyr	Asp	Glu	Glu	
						870					875				880		
90	gag	gag	cac	ttt	ggg	ctt	aat	ttt	gtc	atg	ttc	gat	tca	aat	aag	aaa	2688
	Glu	Glu	His	Phe	Gly	Leu	Asn	Phe	Val	Met	Phe	Asp	Ser	Asn	Lys	Lys	

Agrico B.V.

20030596

PF 54801

40

	885	890	895	
5	agg cat tct ggt aaa cac ctc tat tct ttg agg ata aat gga gac cag Arg His Ser Gly Lys His Leu Tyr Ser Leu Arg Ile Asn Gly Asp Gln 900 905 910			2736
10	ctg gat gac agt gtt tot gat gca ttt cac cta aga cac ttg agg ctt Leu Asp Asp Ser Val Ser Asp Ala Phe His Leu Arg His Leu Arg Leu 915 920 925			2784
15	att aga gtg ttg gac ctg gaa ccc tct tta atc atg gtg aat gat tct Ile Arg Val Leu Asp Leu Glu Pro Ser Leu Ile Met Val Asn Asp Ser 930 935 940			2832
20	ttg ctg aat gaa ata cgc atg ttg aat cat ttg agg tac tta aga att Leu Leu Asn Glu Ile Cys Met Leu Asn His Leu Arg Tyr Leu Arg Ile 945 950 955 960			2880
25	cgg aca caa gtt aaa tat ctg cct ttc tct ttc tca aac ctc tgg aat Arg Thr Gln Val Lys Tyr Leu Pro Phe Ser Phe Ser Asn Leu Trp Asn 965 970 975			2928
30	cta gaa agt ctg ttt gtg tot aac aaa gga tca atc ttg gta cta tta Leu Glu Ser Leu Phe Val Ser Asn Lys Gly Ser Ile Leu Val Leu Leu 980 985 990			2976
35	ccg aga att ttg gat ctt gta aag ttg cga gtg ctg tcc gtg ggt gct Pro Arg Ile Leu Asp Leu Val Lys Leu Arg Val Leu Ser Val Gly Ala 995 1000 1005			3024
40	tgt tot ttc ttt gat atg gat gca gat gaa tca ata ttg ata gca Cys Ser Phe Phe Asp Met Asp Ala Asp Glu Ser Ile Leu Ile Ala 1010 1015 1020			3069
45	aag gac aca aag tta gag aac ttg aga ata tta ggg gaa ctg ttg Lys Asp Thr Lys Leu Glu Asn Leu Arg Ile Leu Gly Glu Leu Leu 1025 1030 1035			3114
50	att tcc tat tcy aaa gat aca atg aat att ttc aaa agg ttt ccc Ile Ser Tyr Ser Lys Asp Thr Met Asn Ile Phe Lys Arg Phe Pro 1040 1045 1050			3159
55	aat ctt cag gtg ctt cag ttt gaa ctc aag gag tca tgg gat tat Asn Leu Gln Val Leu Gln Phe Glu Leu Lys Glu Ser Trp Asp Tyr 1055 1060 1065			3204
60	tca aca gag caa cat tgg ttc ccg aaa ttg gat tgc cta act gaa Ser Thr Glu Gln His Trp Phe Pro Lys Leu Asp Cys Leu Thr Glu 1070 1075 1080			3249
65	cta gaa aca ctc tgt gta ggt ttt aaa agt tca aac aca aac cac Leu Glu Thr Leu Cys Val Gly Phe Lys Ser Ser Asn Thr Asn His 1085 1090 1095			3294
70	tgt ggg tcc tot gtt gtg aca aat cgg cgg tgg gat ttt cac ttc Cys Gly Ser Ser Val Val Thr Asn Arg Pro Trp Asp Phe His Phe 1100 1105 1110			3339
75	cct tca aat ttg aaa gaa ctg ttg ttg tat gac ttt cct ctg aca Pro Ser Asn Leu Lys Glu Leu Leu Leu Tyr Asp Phe Pro Leu Thr 1115 1120 1125			3384
80	tcc gat tca cta tca aca ata gcg aga ctg ccc aac ctt gaa aat Ser Asp Ser Leu Ser Thr Ile Ala Arg Leu Pro Asn Leu Glu Asn 1130 1135 1140			3429
85	ttg tcc ctt tat gat aca atc atc cag gga gaa gaa tgg aac atg Leu Ser Leu Tyr Asp Thr Ile Ile Gln Gly Glu Glu Trp Asn Met 1145 1150 1155			3474
90	ggg gag gaa gac act ttt gag aat ctc aaa ttt ttg aac ttg cgt Gly Glu Glu Asp Thr Phe Glu Asn Leu Lys Phe Leu Asn Leu Arg 1160 1165 1170			3519

Agrico B.V.

20030596

PF 54801

41

5 cta ctg act ctt tcc aag tgg gag gtt gga gag gaa tcc ttc ccc 3564
 Leu Leu Thr Leu Ser Lys Trp Glu Val Gly Glu Glu Ser Phe Pro
 1175 1180 1185

10 aat ctt gag aaa tta aaa ctg cag gaa tgt ggt aag ctt gag gag 3609
 Asn Leu Glu Lys Leu Lys Leu Gln Glu Cys Gly Lys Leu Glu Glu
 1190 1195 1200

15 att cca cct agt ttt gga gat att tat tca ttg aaa ttt atc aaa 3654
 Ile Pro Pro Ser Phe Gly Asp Ile Tyr Ser Leu Lys Phe Ile Lys
 1205 1210 1215

20 att gta aag agt cct caa ctt gaa gat tct gct ctc aag att aag 3699
 Ile Val Lys Ser Pro Gln Leu Glu Asp Ser Ala Leu Lys Ile Lys
 1220 1225 1230

25 aaa tac gct gaa gat atg aga gga ggg aac gat ctt cag atc ctt 3744
 Lys Tyr Ala Glu Asp Met Arg Gly Gly Asn Asp Leu Gln Ile Leu
 1235 1240 1245

30 ggc cag aag aat atc ccc tta ttt aag tag 3774
 Gly Gln Lys Asn Ile Pro Leu Phe Lys
 1250 1255

35 <210> 10

40 <211> 1257

45 <212> PRT

50 <213> Lycopersicon lycopersicum

55 <400> 10

60 Met Glu Lys Arg Lys Asp Ile Glu Glu Ala Asn Asn Ser Leu Val Leu
 1 5 10 15

65 Phe Ser Ala Leu Ser Lys Asp Ile Ala Asn Val Leu Ile Phe Leu Glu
 20 25 30

70 Asn Glu Glu Asn Gln Lys Ala Leu Asp Lys Asp Gln Val Glu Lys Leu
 35 40 45

75 Lys Leu Lys Met Ala Phe Ile Cys Thr Tyr Val Gln Leu Ser Tyr Ser
 50 55 60

80 Asp Phe Glu Gln Phe Glu Asp Ile Met Thr Arg Asn Arg Gln Glu Val
 65 70 75 80

85 Glu Asn Leu Leu Gln Ser Leu Leu Asp Asp Asp Val Leu Thr Ser Leu
 85 90 95

90 Thr Ser Asn Met Asp Asp Cys Ile Ser Leu Tyr His Arg Ser Tyr Lys
 100 105 110

95 Ser Asp Ala Ile Met Met Asp Glu Gln Leu Asp Phe Leu Leu Leu Asn
 115 120 125

100 Leu Tyr His Leu Ser Lys His His Ala Glu Lys Ile Phe Pro Gly Val
 130 135 140

Agrico B.V.

20030596

PF 54801

42

Thr Gln Tyr Glu Val Leu Gln Asn Val Cys Gly Asn Ile Arg Asp Phe
 145 150 155 160
 5 His Gly Leu Ile Leu Asn Gly Cys Ile Lys His Glu Met Val Glu Asn
 165 170 175
 10 Val Leu Pro Leu Phe Gln Leu Met Ala Glu Arg Val Gly His Phe Leu
 180 185 190
 15 Trp Glu Asp Gln Thr Asp Glu Asp Ser Arg Leu Ser Glu Leu Asp Glu
 195 200 205
 20 Asp Glu His Asn Asp Arg Asp Ser Arg Leu Phe Gln Leu Thr His Leu
 210 215 220
 Leu Leu Lys Ile Val Pro Thr Glu Leu Glu Val Met His Ile Cys Tyr
 225 230 235 240
 25 Thr Asn Leu Lys Ala Ser Thr Ser Ala Glu Val Gly Arg Phe Ile Lys
 245 250 255
 30 Lys Leu Leu Glu Thr Ser Pro Asp Ile Leu Arg Glu Tyr Ile Ile Gln
 260 265 270
 35 Leu Gln Glu His Met Leu Thr Val Ile Pro Pro Ser Thr Leu Gly Ala
 275 280 285
 40 Arg Asn Ile His Val Met Met Glu Phe Leu Leu Leu Ile Leu Ser Asp
 290 295 300
 Met Pro Lys Asp Phe Ile His His Asp Lys Leu Phe Asp Leu Leu Ala
 305 310 315 320
 45 His Val Gly Thr Leu Thr Arg Glu Val Ser Thr Leu Val Arg Asp Leu
 325 330 335
 50 Glu Glu Lys Leu Arg Asn Lys Glu Gly Asn Asn Gln Thr Asn Cys Ala
 340 345 350
 55 Thr Leu Asp Leu Leu Glu Asn Ile Glu Leu Leu Lys Lys Asp Leu Lys
 355 360 365
 60 His Val Tyr Leu Lys Ala Pro Asn Ser Ser Gln Cys Cys Phe Pro Met
 370 375 380
 Ser Asp Gly Pro Leu Phe Met His Leu Leu His Met His Leu Asn Asp
 385 390 395 400
 65 Leu Leu Asp Ser Asn Ala Tyr Ser Ile Ser Leu Ile Lys Glu Glu Ile
 405 410 415
 70 Glu Leu Val Ser Gln Glu Leu Glu Phe Ile Arg Ser Phe Phe Gly Asp
 420 425 430

Agrico B.V.

20030596

PF 54801

43

Ala Ala Glu Gln Gly Leu Tyr Lys Asp Ile Trp Ala Arg Val Leu Asp
435 440 445

5 Val Ala Tyr Glu Ala Lys Asp Val Ile Asp Ser Ile Ile Val Arg Asp
450 455 460

10 Asn Gly Leu Leu His Leu Ile Phe Ser Leu Pro Ile Thr Ile Lys Lys
465 470 475 480

Ile Lys Leu Ile Lys Glu Glu Ile Ser Ala Leu Asp Glu Asn Ile Pro
485 490 495

15 Lys Asp Arg Gly Leu Ile Val Val Asn Ser Pro Lys Lys Pro Val Glu
500 505 510

20 Arg Lys Ser Leu Thr Thr Asp Lys Ile Ile Val Gly Phe Glu Glu Glu
515 520 525

25 Thr Asn Leu Ile Leu Arg Lys Leu Thr Ser Gly Pro Ala Asp Leu Asp
530 535 540

30 Val Ile Ser Ile Thr Gly Met Pro Gly Ser Gly Lys Thr Thr Leu Ala
545 550 555 560

Tyr Lys Val Tyr Asn Asp Lys Ser Val Ser Arg His Phe Asp Leu Arg
565 570 575

35 Ala Trp Cys Thr Val Asp Gln Gly Tyr Asp Asp Lys Lys Leu Leu Asp
580 585 590

40 Thr Ile Phe Ser Gln Val Ser Gly Ser Asp Ser Asn Leu Ser Glu Asn
595 600 605

45 Ile Asp Val Ala Asp Lys Leu Arg Lys Gln Leu Phe Gly Lys Arg Tyr
610 615 620

50 Leu Ile Val Leu Asp Asp Val Trp Asp Thr Thr Thr Leu Asp Glu Leu
625 630 635 640

Thr Arg Pro Phe Pro Glu Ala Lys Lys Gly Ser Arg Ile Ile Leu Thr
645 650 655

55 Thr Arg Glu Lys Glu Val Ala Leu His Gly Lys Leu Asn Thr Asp Pro
660 665 670

60 Leu Asp Leu Arg Leu Leu Arg Pro Asp Glu Ser Trp Glu Leu Leu Asp
675 680 685

65 Lys Arg Thr Phe Gly Asn Glu Ser Cys Pro Asp Glu Leu Leu Asp Val
690 695 700

70 Gly Lys Glu Ile Ala Glu Asn Cys Lys Gly Leu Pro Leu Val Ala Asp
705 710 715 720

Leu Ile Ala Gly Val Ile Ala Gly Arg Glu Lys Lys Arg Ser Val Trp

Agrico B.V.

20030596

PF 54801

44

725

730

735

5 Leu Glu Val Gln Ser Ser Leu Ser Ser Phe Ile Leu Asn Ser Glu Val
 740 745 750
 Glu Val Met Lys Val Ile Glu Leu Ser Tyr Asp His Leu Pro His His
 755 760 765
 10 Leu Lys Pro Cys Leu Leu His Phe Ala Ser Trp Pro Lys Asp Thr Pro
 770 775 780
 15 Leu Thr Ile Tyr Leu Phe Thr Val Tyr Leu Gly Ala Glu Gly Phe Val
 785 790 795 800
 20 Glu Lys Thr Glu Met Lys Gly Ile Glu Glu Val Val Lys Ile Tyr Met
 805 810 815
 25 Asp Asp Leu Ile Ser Ser Ser Leu Val Ile Cys Phe Asn Glu Ile Gly
 820 825 830
 30 Asp Ile Leu Asn Phe Gln Ile His Asp Leu Val His Asp Phe Cys Leu
 835 840 845
 Ile Lys Ala Arg Lys Glu Asn Leu Phe Asp Arg Ile Arg Ser Ser Ala
 850 855 860
 35 Pro Ser Asp Leu Leu Pro Arg Gln Ile Thr Ile Asp Tyr Asp Glu Glu
 865 870 875 880
 40 Glu Glu His Phe Gly Leu Asn Phe Val Met Phe Asp Ser Asn Lys Lys
 885 890 895
 45 Arg His Ser Gly Lys His Leu Tyr Ser Leu Arg Ile Asn Gly Asp Gln
 900 905 910
 50 Leu Asp Asp Ser Val Ser Asp Ala Phe His Leu Arg His Leu Arg Leu
 915 920 925
 Ile Arg Val Leu Asp Leu Glu Pro Ser Leu Ile Met Val Asn Asp Ser
 930 935 940
 55 Leu Leu Asn Glu Ile Cys Met Leu Asn His Leu Arg Tyr Leu Arg Ile
 945 950 955 960
 60 Arg Thr Gln Val Lys Tyr Leu Pro Phe Ser Phe Ser Asn Leu Trp Asn
 965 970 975
 65 Leu Glu Ser Leu Phe Val Ser Asn Lys Gly Ser Ile Leu Val Leu Leu
 980 985 990
 70 Pro Arg Ile Leu Asp Leu Val Lys Leu Arg Val Leu Ser Val Gly Ala
 995 1000 1005
 Cys Ser Phe Phe Asp Met Asp Ala Asp Glu Ser Ile Leu Ile Ala
 1010 1015 1020

Agrico B.V.

20030596

PF 54801

45

5 Lys Asp Thr Lys Leu Glu Asn Leu Arg Ile Leu Gly Glu Leu Leu
 1025 1030 1035
 10 Ile Ser Tyr Ser Lys Asp Thr Met Asn Ile Phe Lys Arg Phe Pro
 1040 1045 1050
 15 Asn Leu Gln Val Leu Gln Phe Glu Leu Lys Glu Ser Trp Asp Tyr
 1055 1060 1065
 20 Ser Thr Glu Gln His Trp Phe Pro Lys Leu Asp Cys Leu Thr Glu
 1070 1075 1080
 25 Leu Glu Thr Leu Cys Val Gly Phe Lys Ser Ser Asn Thr Asn His
 1085 1090 1095
 30 Cys Gly Ser Ser Val Val Thr Asn Arg Pro Trp Asp Phe His Phe
 1100 1105 1110
 35 Pro Ser Asn Leu Lys Glu Leu Leu Leu Tyr Asp Phe Pro Leu Thr
 1115 1120 1125
 40 Ser Asp Ser Leu Ser Thr Ile Ala Arg Leu Pro Asn Leu Glu Asn
 1130 1135 1140
 45 Leu Ser Leu Tyr Asp Thr Ile Ile Gln Gly Glu Glu Trp Asn Met
 1145 1150 1155
 50 Gly Glu Glu Asp Thr Phe Glu Asn Leu Lys Phe Leu Asn Leu Arg
 1160 1165 1170
 55 Leu Leu Thr Leu Ser Lys Trp Glu Val Gly Glu Glu Ser Phe Pro
 1175 1180 1185
 60 Asn Leu Glu Lys Leu Lys Leu Gln Glu Cys Gly Lys Leu Glu Glu
 1190 1195 1200
 65 Ile Pro Pro Ser Phe Gly Asp Ile Tyr Ser Leu Lys Phe Ile Lys
 1205 1210 1215
 70 Ile Val Lys Ser Pro Gln Leu Glu Asp Ser Ala Leu Lys Ile Lys
 1220 1225 1230
 75 Lys Tyr Ala Glu Asp Met Arg Gly Gly Asn Asp Leu Gln Ile Leu
 1235 1240 1245
 80 Gly Gln Lys Asn Ile Pro Leu Phe Lys
 1250 1255
 85 <210> 11
 90 <211> 20
 95 <212> DNA
 100 <213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

46

5 <220>
<221> mapping_marker
<222> (1) .. (20)

10 <223>

15 <400> 11
ttgtggttat cgatgagaat 20

<210> 12
20 <211> 22
<212> DNA
25 <213> Solanum bulbocastanum

<220>
30 <221> Mapping_marker
<222> (1) .. (22)
<223>
35

<400> 12
40 gaaacaacag caggatagtg ag 22

<210> 13
45 <211> 20
<212> DNA
50 <213> Solanum bulbocastanum

<220>
55 <221> Mapping_marker
<222> (1) .. (20)
<223>
60

<400> 13
65 ttgtggttat cgatgagaat 20

<210> 14
70 <211> 22
<212> DNA
75 <213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

47

<220>

5 <221> Mapping_marker

<222> (1)..(22)

<223>

10

<400> 14

gaaacaacag caggatagtg ag

22

15

<210> 15

<211> 20

20

<212> DNA

<213> Solanum bulbocastanum

25

<220>

<221> Mapping_Marker

30

<222> (1)..(20)

<223>

35

<400> 15

gaattcagca caaataccaa

20

40

<210> 16

<211> 20

45

<212> DNA

<213> Solanum bulbocastanum

50

<220>

<221> Mapping_Marker

55

<222> (1)..(20)

<223>

60

<400> 16

ttaacgttta ctatcacgag

20

65

<210> 17

<211> 23

<212> DNA

70

<213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

48

<220>
5 <221> Mapping_Marker
<222> (1)..(23)
<223>
10
<400> 17
gtagaaacag cagcctcata agc 23
15
<210> 18
<211> 20
20 <212> DNA
<213> Solanum bulbocastanum
25
<220>
<221> Mapping_marker
30 <222> (1)..(20)
<223>
35
<400> 18
ttctgcctaa ttgccctgtg 20
40 <210> 19
<211> 22
<212> DNA
45 <213> Solanum bulbocastanum
50 <220>
<221> Mapping_marker
<222> (1)..(22)
55 <223>
60 <400> 19
ggggttggga agacaacgac ac 22
65 <210> 20
<211> 23
<212> DNA
70 <213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

49

<220>
<221> Mapping_marker
5 <222> (1)..(23)
<223>

10
<400> 20
aattgcaaga tacagtcaaa tac 23

15 <210> 21
<211> 21
<212> DNA
20 <213> Solanum bulbocastanum

25 <220>
<221> Mapping_marker
<222> (1)..(21)
30 <223>

35 <400> 21
aggcaggatt aacagtagaa g 21

40 <210> 22
<211> 21
<212> DNA
45 <213> Solanum bulbocastanum

50 <220>
<221> Mapping_marker
<222> (1)..(21)
55 <223>

60 <400> 22
catgctttta ggaagaagct c 21

65 <210> 23
<211> 20
<212> DNA
70 <213> Solanum bulbocastanum

<220>

Agrico B.V.

20030596

PF 54801

50

<221> Mapping_marker
5 <222> (1)..(20)
<223>

10 <400> 23
ttgagacaaa gcagctccac 20

15 <210> 24
<211> 20
<212> DNA

20 <213> Solanum bulbocastanum

25 <400> 24
acgtttctca cacctacagg 20

<210> 25
30 <211> 20
<212> DNA

35 <213> Solanum bulbocastanum

<220>

40 <221> Mapping_marker
<222> (1)..(20)
<223>

45

50 <400> 25
tgatggcacg ttgatcgtg 20

<210> 26
55 <211> 20
<212> DNA

60 <213> Solanum bulbocastanum

<400> 26
65 taagatccaa accagccacc 20

<210> 27
<211> 21

70 <212> DNA
<213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

51

<220>

5 <221> Mapping_marker

<222> (1)..(21)

10 <223>

<400> 27
cattatcaca catgtggcta c 21

15 <210> 28

20 <211> 22

<212> DNA

<213> Solanum bulbocastanum

25 <220>

30 <221> mapping_marker

<222> (1)..(22)

<223>

35 <400> 28
attgaaacgg aggaagtaca ac 22

40 <210> 29

<211> 21

45 <212> DNA

<213> Solanum bulbocastanum

50 <220>

<221> mapping_marker

55 <222> (1)..(21)

<223>

60 <400> 29
tttttcatat ggcagaccaa c 21

65 <210> 30

<211> 20

<212> DNA

70 <213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

52

<220>
5 <221> mapping_marker
<222> (1)..(20)
<223>
10
<400> 30
ctactctgct gacatgcagg 20
15 <210> 31
<211> 22
20 <212> DNA
<213> Solanum bulbocastanum
25 <220>
<221> mapping_marker
30 <222> (1)..(22)
<223>
35 <400> 31
gagattotca aaggtgtctt ca 22
40 <210> 32
<211> 20
<212> DNA
45 <213> Solanum bulbocastanum
50 <220>
<221> mapping_marker
55 <222> (1)..(20)
<223>
60 <400> 32
aacctgtgct ttccattcg 20
65 <210> 33
<211> 21
<212> DNA
70 <213> Solanum bulbocastanum

Agrico B.V.

20030596

PF 54801

53

<220>
<221> mapping_marker
5 <222> (1)..(21)
<223>

10 <400> 33
ctttcacaag cgtcactttg g 21

15 <210> 34
<211> 22
<212> DNA
20 <213> Solanum bulbocastanum

25 <220>
<221> mapping_marker
<222> (1)..(22)
30 <223>

35 <400> 34
taaaaagaat caacagggca ac 22

40 <210> 35
<211> 20
<212> DNA
45 <213> Solanum bulbocastanum

50 <220>
<221> mapping_marker
<222> (1)..(20)
55 <223>

60 <400> 35
acgactgctc aaagttggcc 20

65 <210> 36
<211> 20
<212> DNA
70 <213> Solanum bulbocastanum

<220>

Agrico B.V.

20030596

PF 54801

54

5 <221> Mapping_marker
<222> (1)..(20)
<223>

10 <400> 36
ccaagaagcc agttgagagc 20

15 <210> 37
<211> 22
<212> DNA

20 <213> Solanum bulbocastanum

25 <220>
<221> mapping_marker
<222> (1)..(22)

30 <223>

35 <400> 37
gtagattaca ctatggatat gg 22

40 <210> 38
<211> 20
<212> DNA

45 <213> Solanum bulbocastanum

50 <220>
<221> mapping_marker
<222> (1)..(20)

55 <223>

60 <400> 38
cagttagcag caatgtcagc 20

65 <210> 39
<211> 22
<212> DNA

70 <213> Solanum bulbocastanum
<220>

Agrico B.V.

20030596

PF 54801

55

<221> mapping_marker

<222> (1)..(22)

5 <223>

10 <400> 39
cattcaacta ggccaaaagt gg

22

<210> 40

15 <211> 20

<212> DNA

20 <213> Solanum bulbocastanum

<220>

25 <221> mapping_marker

<222> (1)..(20)

30 <223>

35 <400> 40
ccaggtaggt gttttcttcc

20

<210> 41

40 <211> 20

<212> DNA

45 <213> Solanum bulbocastanum

<220>

50 <221> mapping_marker

<222> (1)..(20)

<223>

55

60 <400> 41
gttctaagtc agatgccacc

20

<210> 42

<211> 19

65 <212> DNA

<213> Solanum bulbocastanum

70

<220>

<221> mapping_marker

Agrico B.V.

20030596

PF 54801

56

5 <222> (1)..(19)
<223>

10 <400> 42
aagtgcctcca acacgagcc 19

15 <210> 43
<211> 19
<212> DNA
<213> Solanum bulbocastanum

20 <220>

25 <221> mapping_marker
<222> (1)..(19)
<223>

30 <400> 43
tgagttctctc tacccttccg 19

35 <210> 44
<211> 20

40 <212> DNA
<213> Solanum bulbocastanum

45 <220>

50 <221> mapping_marker
<222> (1)..(20)
<223>

55 <400> 44
ggatatccag catcaatgcc 20

60 <210> 45
<211> 20

65 <212> DNA
<213> Solanum bulbocastanum

70 <220>

<221> mapping_marker

Agrico B.V.

20030596

57

<222> (1)..(20)
<223>

5
<400> 45 20
ggtgagcctc cttgcattcc

10
<210> 46
<211> 19

15
<212> DNA
<213> Solanum bulbocastanum

20
<220>
<221> mapping_marker

25
<222> (1)..(19)
<223>

30
<400> 46 19
cctgagggaa gatgtcacg

35
<210> 47
<211> 21
<212> DNA

40
<213> Solanum bulbocastanum

45
<220>
<221> mapping_marker

50
<222> (1)..(21)
<223>

55
<400> 47 21
cctagttag agtgagtaga c

60
<210> 48
<211> 22
<212> DNA

65
<213> Solanum bulbocastanum

70
<220>
<221> mapping_marker
<222> (1)..(22)

Agrico B.V.

20030596

PF 54801

58

<223>

5 <400> 48
gtgatatatt gctcaaggat cc 22

10 <210> 49
 <211> 20
 <212> DNA

15 <213> Solanum bulbocastanum

20 <220>
 <221> mapping_marker
 <222> (1)..(20)

25 <223>

30 <400> 49
gttgctggct gtcactgatc 20

35 <210> 50
 <211> 20
 <212> DNA

40 <213> Solanum bulbocastanum

45 <220>
 <221> mapping_marker
 <222> (1)..(20)

50 <223>

55 <400> 50
gtgatgtgca gggttcaagg 20

 <210> 51

60 <211> 22
 <212> DNA

65 <213> Solanum bulbocastanum

 <220>

70 <221> mapping_marker
 <222> (1)..(22)

Agrico B.V.

20030596

PF 54801

59

<223>

5 <400> 51
gattagtgtg gatcttagct tg

22

10 <210> 52

<211> 22

<212> DNA

15 <213> Solanum bulbocastanum

20 <220>

<221> mapping_marker

<222> (1) .. (22)

25 <223>

30 <400> 52
aaatctctct cacaattatc cc

22

<210> 53

35 <211> 21

<212> DNA

40 <213> Solanum bulbocastanum

<220>

45 <221> mapping_marker

<222> (1) .. (21)

<223>

50

55 <400> 53
ctattgactg aacctgctga g

21

<210> 54

60 <211> 21

<212> DNA

<213> Solanum bulbocastanum

65

<220>

70 <221> mapping_marker

<222> (1) .. (21)

<223>

Agrico B.V.

20030596

PF 54801

60

5	<400> 54 tgaagtcatt tagtcacag c	21
	<210> 55	
10	<211> 20	
	<212> DNA	
15	<213> Solanum bulbocastanum	
	<220>	
20	<221> mapping_marker	
	<222> (1) .. (20)	
25	<223>	
	<400> 55 agatcggagt gtgaacatgg	20
30	<210> 56	
	<211> 21	
35	<212> DNA	
	<213> Solanum bulbocastanum	
40	<220>	
	<221> mapping_marker	
45	<222> (1) .. (21)	
	<223>	
50	<400> 56 cttctacttc tagtcgactg c	21
55	<210> 57	
	<211> 20	
60	<212> DNA	
	<213> Solanum bulbocastanum	
65	<220>	
	<221> mapping_marker	
70	<222> (1) .. (20)	
	<223>	

Agrico B.V.

20030596

PF 54801

61

5 <400> 57
 cgtagtccat ctgaagctcc 20

10 <210> 58
 <211> 20
 <212> DNA
 <213> Solanum bulbocastanum

15 <220>
 <221> mapping_marker
20 <222> (1)..(20)
 <223>

25 <400> 58
 tcttctctcg ctagtctgctg 20

30 <210> 59
 <211> 23
35 <212> DNA
 <213> Solanum bulbocastanum

40 <220>
 <221> mapping_marker
45 <222> (1)..(23)
 <223>

50 <400> 59
 actattctca cgtaagggga cac 23

55 <210> 60
 <211> 23
 <212> DNA
60 <213> Solanum bulbocastanum

65 <220>
 <221> mapping_marker
70 <222> (1)..(23)
 <223>

Agrico B.V.

20030596

PF 54801

62

5 <400> 60
gtgtacatgt atgaaactct agc 23

5 <210> 61
 <211> 22

10 <212> DNA
 <213> Solanum bulbocastanum

15 <220>
 <221> mapping_marker

20 <222> (1)..(22)
 <223>

25 <400> 61
gttcctttca atcagaaagt ag 22

30 <210> 62
 <211> 21

35 <212> DNA
 <213> Solanum bulbocastanum

40 <220>
 <221> mapping_marker

45 <222> (1)..(21)
 <223>

50 <400> 62
ctttggatga gtcaaaaggc t 21

55 <210> 63
 <211> 20

60 <212> DNA
 <213> Solanum bulbocastanum

65 <220>
 <221> mapping_marker

70 <222> (1)..(20)
 <223>

Agrico B.V.

20030596

PF 54801

63

<400> 63
caagttacgg caaccaagag 20

5 <210> 64
<211> 22
<212> DNA
10 <213> Solanum bulbocastanum

15 <220>
<221> mapping_marker
<222> (1) .. (22)
20 <223>

25 <400> 64
ctttgacaca gtgtagaat gc 22
<210> 65
30 <211> 20
<212> DNA
35 <213> Solanum bulbocastanum

40 <220>
<221> mapping_marker
<222> (1) .. (20)
45 <223>

50 <400> 65
cgtgatctag gagttacgac 20
<210> 66
55 <211> 25
<212> DNA
<213> Solanum bulbocastanum
60

<220>
65 <221> mapping_marker
<222> (1) .. (25)
<223>
70

<400> 66

Agrico B.V.

20030596

64

cttatttttaa atacaagaca tctgg 25

5 <210> 67
<211> 20
<212> DNA

10 <213> Solanum bulbocastanum

15 <220>
<221> mapping_marker
<222> (1)..(20)

20 <223>

25 <400> 67
cagaggaaag tcaaccaacg 20

<210> 68

30 <211> 20
<212> DNA

<213> Solanum bulbocastanum

35 <220>
<221> mapping_marker
<222> (1)..(20)

40 <223>

45

<400> 68
cagaggaaag tcaaccaacg 20

50 <210> 69
<211> 24
<212> DNA

55 <213> Solanum bulbocastanum

60 <220>
<221> mapping_marker

65 <222> (1)..(24)
<223>

70 <400> 69
tcggctatga ctgggcacaa caga 24

Agrico B.V.

20030596

PF 54801

65

<210> 70
5 <211> 24
<212> DNA
10 <213> Solanum bulbocastanum

<220>
15 <221> mapping_marker
<222> (1)..(24)
20 <223>

<400> 70
25 aagaaggcga tagaaggcga tgcg

<210> 71
30 <211> 18
<212> DNA
<213> Solanum bulbocastanum
35

<220>
40 <221> mapping_marker
<222> (1)..(18)
<223>
45

<400> 71
50 tgtaaaacga cggccagt

<210> 72
<211> 19
55 <212> DNA
<213> Solanum bulbocastanum
60

<220>
65 <221> mapping_marker
<222> (1)..(19)
<223>
70

<400> 72
ggaaacagct atgaccatg

24

18

19

Agrico B.V.

20030596

66

5 <210> 73
 <211> 18
 <212> DNA
 <213> Solanum bulbocastanum

10 <220>
 <221> primer
15 <222> (1)..(18)
 <223>

20 <400> 73
 ttcagcacaa ataccaat 18

25 <210> 74
 <211> 18

30 <212> DNA
 <213> Solanum bulbocastanum

35 <220>
 <221> primer

40 <222> (1)..(18)
 <223>

45 <400> 74
 gatgttcccc ttctttta 18

50 <210> 75
 <211> 20
 <212> DNA

55 <213> Solanum bulbocastanum

60 <220>
 <221> primer
 <222> (1)..(20)

65 <223>

70 <400> 75
 ttgtgggttat cgatgagaat 20

Agrico B.V.

20030596

PF 54801

67

<210> 76
<211> 20
5 <212> DNA
<213> Solanum bulbocastanum
10 <220>
<221> primer
15 <222> (1) .. (20)
<223>
20 <400> 76
acctggcggt ccttattttt
25 <210> 77
<211> 15
<212> DNA
30 <213> Solanum bulbocastanum
35 <220>
<221> misc_feature
<222> (1) .. (1)
40 <223> N= A+T+G+C
45 <220>
<221> primer
<222> (1) .. (15)
50 <223>
55 <220>
<221> misc_feature
<222> (7) .. (7)
60 <223> W=A+T
65 <220>
<221> misc_feature
<222> (6) .. (6)
70 <223> S=G+C

20

Agrico B.V.

20030596

PF 54801

68

<220>
5 <221> misc_feature
<222> (10)..(10)
<223> N=A+T+G+C
10
<220>
15 <221> misc_feature
<222> (12)..(12)
<223> W=A+T
20
<400> 77
ngtcaswgan awgaa 15
25
<210> 78
<211> 21
30 <212> DNA
<213> Solanum bulbocastanum
35
<220>
<221> primer
40 <222> (1)..(21)
<223>
45
<400> 78
gatggagcgg aaaagccggt g 21
50
<210> 79
<211> 21
<212> DNA
55
<213> Solanum bulbocastanum
60
<220>
<221> primer
<222> (1)..(21)
65
<223>
70
<400> 79
ggtgttttgt agcatctcca g 21

69

<210> 80
<211> 20
5 <212> DNA
<213> Solanum bulbocastanum

10 <220>
<221> primer
15 <222> (1)..(20)
<223>

20 <400> 80
ccatgattac gccaaagctgg

25 <210> 81
<211> 20
<212> DNA
30 <213> Solanum bulbocastanum

35 <220>
<221> primer
<222> (1)..(20)
40 <223>

45 <400> 81
ggtttttccca gtcacgacgt

50 <210> 82
<211> 20
<212> DNA
55 <213> Solanum bulbocastanum

60 <220>
<221> primer
<222> (1)..(20)
65 <223>

70 <400> 82
agaaagctca ccagtggacc

<210> 83

20

20

20

70

5 <211> 20
<212> DNA
<213> Solanum bulbocastanum

10 <220>
<221> primer
<222> (1)..(20)
15 <223>

20 <400> 83
atttatggct gcagaggacc 20

25 <210> 84
<211> 21
<212> DNA
30 <213> Solanum bulbocastanum

35 <400> 84
aagtccaatt gctcatccat c 21

<210> 85
40 <211> 19
<212> DNA
<213> Solanum bulbocastanum
45

<220>
50 <221> primer
<222> (1)..(19)
<223>
55

<400> 85
60 tgcacccatgc acgaaggta 19

<210> 86
65 <211> 22
<212> DNA
<213> Solanum bulbocastanum
70
<220>

Agrico B.V.

71

<221> primer
<222> (1)..(22)
5 <223>

10 <400> 86
caatwtgtggt tcccgaatt gg 22

<210> 87

15 <211> 25
<212> DNA
<213> Solanum bulbocastanum

20

<220>

25 <221> primer
<222> (1)..(25)
<223>

30

<400> 87
atggaaaaac gaaaagataa tgaag 25

35

<210> 88

40 <211> 25
<212> DNA
<213> Solanum bulbocastanum

45

<220>

50 <221> primer
<222> (1)..(25)
<223>

55

<400> 88
ctaacttaaat aacgggatat ccttc 25

60

<210> 89
<211> 20

65 <212> DNA
<213> Solanum bulbocastanum

70

<220>
<221> primer

Agrico B.V.

72

<222> (1)..(20)
<223>
5

<400> 89
cccatgactc cttaggtttg 20
10

<210> 90

<211> 21
15
<212> DNA

<213> Solanum bulbocastanum
20

<220>

<221> primer
25
<222> (1)..(21)
<223>
30

<400> 90
ggtgggggttg ggaagacaac g 21
35

<210> 91

<211> 19
40
<212> DNA

<213> Solanum bulbocastanum

45

<220>

<221> primer
50
<222> (1)..(19)
<223>

55

<400> 91
gtagactgcg taccaatc 19

60

<210> 92

<211> 16

<212> DNA
65
<213> solanum bulbocastanum

70

<220>

<221> primer

73

<222> (1)..(16)
<223>

5
<400> 92
gatgagtcct gagtaa 16

10
<210> 93
<211> 2913

15
<212> DNA
<213> Solanum bulbocastanum

20
<220>
<221> CDS

25
<222> (1)..(2913)
<223> Rpi-blb or RB (Song, PNAS, 2003, 9128-9133)

30
<400> 93
atg gct gaa gct ttc att caa gtt ctg cta gac aat ctc act tct ttc 48
Met Ala Glu Ala Phe Ile Gln Val Leu Leu Asp Asn Leu Thr Ser Phe
1 5 10 15

35
ctc aaa ggg gaa ctt gta ttg ctt ttc ggt ttt caa gat gag ttc caa 96
Leu Lys Gly Glu Leu Val Leu Leu Phe Gly Phe Gln Asp Glu Phe Gln
20 25 30

40
agg ott tca agc atg ttt tct aca att caa gcc gtc ctt gaa gat gct 144
Arg Leu Ser Ser Met Phe Ser Thr Ile Gln Ala Val Leu Glu Asp Ala
35 40 45

45
cag gag aag caa ctc aac aac aag cct cta gaa aat tgg ttg caa aaa 192
Gln Glu Lys Gln Leu Asn Asn Lys Pro Leu Glu Asn Trp Leu Gln Lys
50 55 60

50
ctc aat gct gct aca tat gaa gtc gat gac atc ttg gat gaa tat aaa 240
Leu Asn Ala Ala Thr Tyr Glu Val Asp Asp Ile Leu Asp Glu Tyr Lys
65 70 75 80

55
acc aag gcc aca aga ttc tcc cag tct gaa tat gcc cgt tat cat cca 268
Thr Lys Ala Thr Arg Phe Ser Gln Ser Glu Tyr Gly Arg Tyr His Pro
85 90 95

60
aag gtt atc cct ttc cgt cac aag gtc ggg aaa agg atg gac caa gtg 336
Lys Val Ile Pro Phe Arg His Lys Val Gly Lys Arg Met Asp Gln Val
100 105 110

65
atg aaa aaa cta aag gca att gct gag gaa aga aag aat ttt cat ttg 384
Met Lys Lys Leu Lys Ala Ile Ala Glu Glu Arg Lys Asn Phe His Leu
115 120 125

70
cac gaa aaa att gta gag aga caa gct gtt aga cgg gaa aca ggt tct 432
His Glu Lys Ile Val Glu Arg Gln Ala Val Arg Arg Glu Thr Gly Ser
130 135 140

70
gta tta acc gaa ccg cag gtt tat gga aga gac aaa gag aaa gat gag 480
Val Leu Thr Glu Pro Gln Val Tyr Gly Arg Asp Lys Glu Lys Asp Glu
145 150 155 160

70
ata gtg aaa atc cta ata aac aat gtt agt gat gcc caa cac ctt tca 528
Ile Val Lys Ile Leu Ile Asn Asn Val Ser Asp Ala Gln His Leu Ser

Agrico B.V.

20030596

PF 54801

74

		165		170		175		
5	gtc ctc cca ata ctt ggt atg ggg gga tta gga aaa acg act ctt gcc						576	
	Val Leu Pro Ile Leu Gly Met Gly Gly Leu Gly Lys Thr Thr Leu Ala	180		185		190		
10	caa atg gtc ttc aat gac cag aga gtt act gag cat ttc cat tcc aaa						624	
	Gln Met Val Phe Asn Asp Gln Arg Val Thr Glu His Phe His Ser Lys	195		200		205		
15	ata tgg att tgt gtc tcg gaa gat ttt gat gag aag agg tta ata aag						672	
	Ile Trp Ile Cys Val Ser Glu Asp Phe Asp Glu Lys Arg Leu Ile Lys	210		215		220		
20	gca att gta gaa tct att gaa gga agg cca cta ctt ggt gag atg gac						720	
	Ala Ile Val Glu Ser Ile Glu Gly Arg Pro Leu Leu Gly Glu Met Asp	225		230		235		
25	ttg gct cca ctt caa aag aag ctt cag gag ttg ctg aat gga aaa aga						768	
	Leu Ala Pro Leu Gln Lys Lys Leu Gln Glu Leu Leu Asn Gly Lys Arg	245		250		255		
30	tac ttg ctt gtc tta gat gat gtt tgg aat gaa gat caa cag aag tgg						816	
	Tyr Leu Leu Val Leu Asp Asp Val Trp Asn Glu Asp Gln Gln Lys Trp	260		265		270		
35	gct aat tta aga gca gtc ctg aag gtt gga gca agt ggt gct tct gtt						864	
	Ala Asn Leu Arg Ala Val Leu Lys Val Gly Ala Ser Gly Ala Ser Val	275		280		285		
40	cta acc act act cgt ctt gaa aag gtt gga tca att atg gga aca ttg						912	
	Leu Thr Thr Thr Arg Leu Glu Lys Val Gly Ser Ile Met Gly Thr Leu	290		295		300		
45	caa cca tat gaa ctg tca aat ctg tot caa gaa gat tgt tgg ttg ttg						960	
	Gln Pro Tyr Glu Leu Ser Asn Leu Ser Gln Glu Asp Cys Trp Leu Leu	305		310		315		
50	ttc atg caa cgt gca ttt gga cac caa gaa gaa ata aat cca aac ctt						1008	
	Phe Met Gln Arg Ala Phe Gly His Gln Glu Ile Asn Pro Asn Leu	325		330		335		
55	gtg gca atc gga aag gag att gtg aaa aaa agt ggt ggt gtg cct cta						1056	
	Val Ala Ile Gly Lys Glu Ile Val Lys Lys Ser Gly Gly Val Pro Leu	340		345		350		
60	gca gcc aaa act ctt gga ggt att ttg tgc ttc aag aga gaa gaa aga						1104	
	Ala Ala Lys Thr Leu Gly Gly Ile Leu Cys Phe Lys Arg Glu Glu Arg	355		360		365		
65	gca tgg gaa cat gtg aga gac agt ccg att tgg aat ttg cct caa gat						1152	
	Ala Trp Glu His Val Arg Asp Ser Pro Ile Trp Asn Leu Pro Gln Asp	370		375		380		
70	gaa agt tct att ctg cct gcc ctg agg ctt agt tac cat caa ctt cca						1200	
	Glu Ser Ser Ile Leu Pro Ala Leu Arg Leu Ser Tyr His Gln Leu Pro	385		390		395		
75	ctt gat ttg aaa caa tgc ttt gcg tat tgt gcg gtg ttc cca aag gat						1248	
	Leu Asp Leu Lys Gln Cys Phe Ala Tyr Cys Ala Val Phe Pro Lys Asp	405		410		415		
80	gcc aaa atg gaa aaa gaa aag cta atc tot ctc tgg atg gcg cat ggt						1296	
	Ala Lys Met Glu Lys Glu Lys Leu Ile Ser Leu Trp Met Ala His Gly	420		425		430		
85	ttt ctt tta tca aaa gga aac atg gag cta gag gat gtg ggc gat gaa						1344	
	Phe Leu Leu Ser Lys Gly Asn Met Glu Leu Glu Asp Val Gly Asp Glu	435		440		445		
90	gta tgg aaa gaa tta tac ttg agg tct ttt ttc caa gag att gaa gtt						1392	
	Val Trp Lys Glu Leu Tyr Leu Arg Ser Phe Phe Gln Glu Ile Glu Val	450		455		460		

Agrico B.V.

75

	aaa gat ggt aaa act tat ttc aag atg cat gat ctc atc cat gat ttg	1440
	Lys Asp Gly Lys Thr Tyr Phe Lys Met His Asp Leu Ile His Asp Leu	
	465 470 475 480	
5	gca aca tct ctg ttt tca gca aac aca tca agc agc aat atc cgt gaa	1488
	Ala Thr Ser Leu Phe Ser Ala Asn Thr Ser Ser Asn Ile Arg Glu	
	485 490 495	
10	ata aat aaa cac agt tac aca cat atg atg tcc att ggt ttc gcc gaa	1536
	Ile Asn Lys His Ser Tyr Thr His Met Met Ser Ile Gly Phe Ala Glu	
	500 505 510	
15	gtg gtg ttt ttt tac act ctt ccc ccc ttg gaa aag ttt atc tct tta	1584
	Val Val Phe Phe Tyr Thr Leu Pro Pro Leu Glu Lys Phe Ile Ser Leu	
	515 520 525	
20	aga gtg ctt aat cta ggt gat tct aca ttt aat aag tta cca tct tcc	1632
	Arg Val Leu Asn Leu Gly Asp Ser Thr Phe Asn Lys Leu Pro Ser Ser	
	530 535 540	
25	att gga gat cta gta cat tta aga tac ttg aac ctg tat ggc agt ggc	1680
	Ile Gly Asp Leu Val His Leu Arg Tyr Leu Asn Leu Tyr Gly Ser Gly	
	545 550 555 560	
30	atg cgt agt ctt cca aag cag tta tgc aag ctt caa aat ctg caa act	1728
	Met Arg Ser Leu Pro Lys Gln Leu Cys Lys Leu Gln Asn Leu Gln Thr	
	565 570 575	
35	ctt gat cta caa tat tgc acc aag ctt tgt tgt ttg cca aaa gaa aca	1776
	Leu Asp Leu Gln Tyr Cys Thr Lys Leu Cys Cys Leu Pro Lys Glu Thr	
	580 585 590	
40	agt aaa ctt ggt agt ctc cga aat ctt tta ctt gat ggt agc cag tca	1824
	Ser Lys Leu Gly Ser Leu Arg Asn Leu Leu Leu Asp Gly Ser Gln Ser	
	595 600 605	
45	ttg act tgt atg cca cca agg ata gga tca ttg aca tgc ctt aag act	1872
	Leu Thr Cys Met Pro Pro Arg Ile Gly Ser Leu Thr Cys Leu Lys Thr	
	610 615 620	
50	cta ggt caa ttt gtt gtt gga agg aag aaa ggt tat caa ctt ggt gaa	1920
	Leu Gly Gln Phe Val Val Gly Arg Lys Lys Gly Tyr Gln Leu Gly Glu	
	625 630 635 640	
55	cta gga aac cta aat ctc tat ggc tca att aaa atc tct cat ctt gag	1968
	Leu Gly Asn Leu Asn Leu Tyr Gly Ser Ile Lys Ile Ser His Leu Glu	
	645 650 655	
60	aga gtg aag aat gat aag gac gca aaa gaa gcc aat tta tct gca aaa	2016
	Arg Val Lys Asn Asp Lys Asp Ala Lys Glu Ala Asn Leu Ser Ala Lys	
	660 665 670	
65	ggg aat ctg cat tct tta agc atg agt tgg aat aac ttt gga cca cat	2064
	Gly Asn Leu His Ser Leu Ser Met Ser Trp Asn Asn Phe Gly Pro His	
	675 680 685	
70	ata tat gaa tca gaa gaa gtt aaa gtg ctt gaa gcc ctc aaa cca cac	2112
	Ile Tyr Glu Ser Glu Glu Val Lys Val Leu Glu Ala Leu Lys Pro His	
	690 695 700	
75	tcc aat ctg act tct tta aaa atc tat ggc ttc aga gga atc cat ctc	2160
	Ser Asn Leu Thr Ser Leu Lys Ile Tyr Gly Phe Arg Gly Ile His Leu	
	705 710 715 720	
80	cca gag tgg atg aat cac tca gta ttg aaa aat att gtc tct att cta	2208
	Pro Glu Trp Met Asn His Ser Val Leu Lys Asn Ile Val Ser Ile Leu	
	725 730 735	
85	att agc aac ttc aga aac tgc tca tgc tta cca ccc ttt ggt gat ctg	2256
	Ile Ser Asn Phe Arg Asn Cys Ser Cys Leu Pro Pro Phe Gly Asp Leu	
	740 745 750	

Agrico B.V.

20030596

76

	cct tgt cta gaa agt cta gag tta cac tgg ggg tct gcg gat gtg gag	2304
	Pro Cys Leu Glu Ser Leu Glu Leu His Trp Gly Ser Ala Asp Val Glu	
	755 760 765	
5	tat gtt gaa gaa gtg gat att gat gtt cat tct gga ttc ccc aca aga	2352
	Tyr Val Glu Glu Val Asp Ile Asp Val His Ser Gly Phe Pro Thr Arg	
	770 775 780	
10	ata agg ttt cca tcc ttg agg aaa ctt gat ata tgg gac ttt ggt agt	2400
	Ile Arg Phe Pro Ser Leu Arg Lys Leu Asp Ile Trp Asp Phe Gly Ser	
	785 790 795 800	
15	ctg aaa gga ttg ctg aaa aag gaa gga gaa gag caa ttc cct gtg ctt	2448
	Leu Lys Gly Leu Leu Lys Lys Glu Gly Glu Glu Gln Phe Pro Val Leu	
	805 810 815	
20	gaa gag atg ata att cac gag tgc cct ttt ctg acc ctt tct tct aat	2496
	Glu Glu Met Ile Ile His Glu Cys Pro Phe Leu Thr Leu Ser Ser Asn	
	820 825 830	
25	ctt agg gct ctt act tcc ctg aga att tgc tat aat aaa gta gct act	2544
	Leu Arg Ala Leu Thr Ser Leu Arg Ile Cys Tyr Asn Lys Val Ala Thr	
	835 840 845	
30	tca ttc cca gaa gag atg ttc aaa aac ctt gca aat ctg aaa tac ttg	2592
	Ser Phe Pro Glu Glu Met Phe Lys Asn Leu Ala Asn Leu Lys Tyr Leu	
	850 855 860	
35	aca atc tct cgg tgc aat aat ctg aaa gag ctg cct acc agc ttg gct	2640
	Thr Ile Ser Arg Cys Asn Asn Leu Lys Glu Leu Pro Thr Ser Leu Ala	
	865 870 875 880	
40	agt ctg aat gct ttg aaa agt cta aaa att caa ttg tgt tgc gca cta	2688
	Ser Leu Asn Ala Leu Lys Ser Leu Lys Ile Gln Leu Cys Cys Ala Leu	
	885 890 895	
45	gag agt ctg cct gag gaa ggg ctg gaa ggt tta tct tca ctg aca gag	2736
	Glu Ser Leu Pro Glu Glu Gly Leu Glu Gly Leu Ser Ser Leu Thr Glu	
	900 905 910	
50	tta ttt gtt gaa cac tgt aac atg cta aaa tgt tta cca gag gga ttg	2784
	Leu Phe Val Glu His Cys Asn Met Leu Lys Cys Leu Pro Glu Gly Leu	
	915 920 925	
55	cag cac cta aca acc ctg aca agt tta aaa att cgg gga tgt cca caa	2832
	Gln His Leu Thr Thr Leu Thr Ser Leu Lys Ile Arg Gly Cys Pro Gln	
	930 935 940	
60	ctg atc aag cgg tgt gag aag gga ata gga gaa gac tgg cac aaa att	2880
	Leu Ile Lys Arg Cys Glu Lys Gly Ile Gly Glu Asp Trp His Lys Ile	
	945 950 955 960	
65	tct cac att cct aat gtg aat ata tat att taa	2913
	Ser His Ile Pro Asn Val Asn Ile Tyr Ile	
	965 970	
	<210> 94	
	<211> 970	
	<212> PRT	
	<213> Solanum bulbocastanum	
	<400> 94	
70	Met Ala Glu Ala Phe Ile Gln Val Leu Leu Asp Asn Leu Thr Ser Phe	
	1 5 10 15	

Agrico B.V.

20030596

PF 54801

77

Leu Lys Gly Glu Leu Val Leu Leu Phe Gly Phe Gln Asp Glu Phe Gln
 20 25 30

5 Arg Leu Ser Ser Met Phe Ser Thr Ile Gln Ala Val Leu Glu Asp Ala
 35 40 45

10 Gln Glu Lys Gln Leu Asn Asn Lys Pro Leu Glu Asn Trp Leu Gln Lys
 50 55 60

15 Leu Asn Ala Ala Thr Tyr Glu Val Asp Asp Ile Leu Asp Glu Tyr Lys
 65 70 75 80

Thr Lys Ala Thr Arg Phe Ser Gln Ser Glu Tyr Gly Arg Tyr His Pro
 85 90 95

20 Lys Val Ile Pro Phe Arg His Lys Val Gly Lys Arg Met Asp Gln Val
 100 105 110

25 Met Lys Lys Leu Lys Ala Ile Ala Glu Glu Arg Lys Asn Phe His Leu
 115 120 125

30 His Glu Lys Ile Val Glu Arg Gln Ala Val Arg Arg Glu Thr Gly Ser
 130 135 140

35 Val Leu Thr Glu Pro Gln Val Tyr Gly Arg Asp Lys Glu Lys Asp Glu
 145 150 155 160

Ile Val Lys Ile Leu Ile Asn Asn Val Ser Asp Ala Gln His Leu Ser
 165 170 175

40 Val Leu Pro Ile Leu Gly Met Gly Gly Leu Gly Lys Thr Thr Leu Ala
 180 185 190

45 Gln Met Val Phe Asn Asp Gln Arg Val Thr Glu His Phe His Ser Lys
 195 200 205

50 Ile Trp Ile Cys Val Ser Glu Asp Phe Asp Glu Lys Arg Leu Ile Lys
 210 215 220

55 Ala Ile Val Glu Ser Ile Glu Gly Arg Pro Leu Leu Gly Glu Met Asp
 225 230 235 240

Leu Ala Pro Leu Gln Lys Lys Leu Gln Glu Leu Leu Asn Gly Lys Arg
 245 250 255

60 Tyr Leu Leu Val Leu Asp Asp Val Trp Asn Glu Asp Gln Gln Lys Trp
 260 265 270

65 Ala Asn Leu Arg Ala Val Leu Lys Val Gly Ala Ser Gly Ala Ser Val
 275 280 285

70 Leu Thr Thr Thr Arg Leu Glu Lys Val Gly Ser Ile Met Gly Thr Leu
 290 295 300

Gln Pro Tyr Glu Leu Ser Asn Leu Ser Gln Glu Asp Cys Trp Leu Leu

Agrico B.V.

20030596

PF 54801

78

	305		310		315		320
5	Phe Met Gln Arg	Ala 325	Phe Gly His Gln	Glu 330	Glu Ile Asn Pro	Asn 335	Leu
10	Val Ala Ile Gly	Lys 340	Glu Ile Val Lys	Lys 345	Ser Gly Gly Val	Pro 350	Leu
15	Ala Ala Lys Thr	Leu 355	Gly Gly Ile Leu	Cys 360	Phe Lys Arg Glu	Glu 365	Arg
20	Ala Trp Glu His	Val 370	Arg Asp Ser Pro	Ile 375	Trp Asn Leu Pro	Gln 380	Asp
25	Glu Ser Ser Ile	Leu 385	Pro Ala Leu Arg	Leu 390	Ser Tyr His Gln	Leu 395	Pro
30	Leu Asp Leu Lys	Gln 405	Cys Phe Ala Tyr	Cys 410	Ala Val Phe Pro	Lys 415	Asp
35	Ala Lys Met Glu	Lys 420	Glu Lys Leu Ile	Ser 425	Leu Trp Met Ala	His 430	Gly
40	Phe Leu Leu Ser	Lys 435	Gly Asn Met Glu	Leu 440	Glu Asp Val Gly	Asp 445	Glu
45	Val Trp Lys Glu	Leu 450	Tyr Leu Arg Ser	Phe 455	Phe Gln Glu Ile	Glu 460	Val
50	Lys Asp Gly Lys	Thr 465	Tyr Phe Lys Met	His 470	Asp Leu Ile His	Asp 475	Leu
55	Ala Thr Ser Leu	Phe 485	Ser Ala Asn Thr	Ser 490	Ser Ser Asn Ile	Arg 495	Glu
60	Ile Asn Lys His	Ser 500	Tyr Thr His Met	Met 505	Ser Ile Gly Phe	Ala 510	Glu
65	Val Val Phe Phe	Tyr 515	Thr Leu Pro Pro	Leu 520	Glu Lys Phe Ile	Ser 525	Leu
70	Arg Val Leu Asn	Leu 530	Gly Asp Ser Thr	Phe 535	Asn Lys Leu Pro	Ser 540	Ser
	Ile Gly Asp Leu	Val 545	His Leu Arg Tyr	Leu 550	Asn Leu Tyr Gly	Ser 555	Gly
	Met Arg Ser Leu	Pro 565	Lys Gln Leu Cys	Lys 570	Leu Gln Asn Leu	Gln 575	Thr
	Leu Asp Leu Gln	Tyr 580	Cys Thr Lys Leu	Cys 585	Cys Leu Pro Lys	Glu 590	Thr
	Ser Lys Leu Gly	Ser 595	Leu Arg Asn Leu	Leu 600	Leu Asp Gly Ser	Gln 605	Ser

Agrico B.V.

20030596

PF 54801

79

5 Leu Thr Cys Met Pro Pro Arg Ile Gly Ser Leu Thr Cys Leu Lys Thr
 610 615 620
 10 Leu Gly Gln Phe Val Val Gly Arg Lys Lys Gly Tyr Gln Leu Gly Glu
 625 630 635 640
 15 Arg Val Lys Asn Asp Lys Asp Ala Lys Glu Ala Asn Leu Ser Ala Lys
 650 665 670
 20 Gly Asn Leu His Ser Leu Ser Met Ser Trp Asn Asn Phe Gly Pro His
 675 680 685
 25 Ile Tyr Glu Ser Glu Glu Val Lys Val Leu Glu Ala Leu Lys Pro His
 690 695 700
 30 Ser Asn Leu Thr Ser Leu Lys Ile Tyr Gly Phe Arg Gly Ile His Leu
 705 710 715 720
 35 Ile Ser Asn Phe Arg Asn Cys Ser Cys Leu Pro Pro Phe Gly Asp Leu
 740 745 750
 40 Pro Cys Leu Glu Ser Leu Glu Leu His Trp Gly Ser Ala Asp Val Glu
 755 760 765
 45 Tyr Val Glu Glu Val Asp Ile Asp Val His Ser Gly Phe Pro Thr Arg
 770 775 780
 50 Ile Arg Phe Pro Ser Leu Arg Lys Leu Asp Ile Trp Asp Phe Gly Ser
 785 790 795 800
 55 Leu Lys Gly Leu Leu Lys Lys Glu Gly Glu Glu Gln Phe Pro Val Leu
 805 810 815
 60 Glu Glu Met Ile Ile His Glu Cys Pro Phe Leu Thr Leu Ser Ser Asn
 820 825 830
 65 Leu Arg Ala Leu Thr Ser Leu Arg Ile Cys Tyr Asn Lys Val Ala Thr
 835 840 845
 70 Ser Phe Pro Glu Glu Met Phe Lys Asn Leu Ala Asn Leu Lys Tyr Leu
 850 855 860
 Thr Ile Ser Arg Cys Asn Asn Leu Lys Glu Leu Pro Thr Ser Leu Ala
 865 870 875 880
 Ser Leu Asn Ala Leu Lys Ser Leu Lys Ile Gln Leu Cys Cys Ala Leu
 885 890 895

BASF AG GUX C100

Agrico B.V.

20030596

+49 621 6021183
PF 54801

5.221/221

80

Glu Ser Leu Pro Glu Glu Gly Leu Glu Gly Leu Ser Ser Leu Thr Glu
900 905 910

5

Leu Phe Val Glu His Cys Asn Met Leu Lys Cys Leu Pro Glu Gly Leu
915 920 925

10

Gln His Leu Thr Thr Leu Thr Ser Leu Lys Ile Arg Gly Cys Pro Gln
930 935 940

15

Leu Ile Lys Arg Cys Glu Lys Gly Ile Gly Glu Asp Trp His Lys Ile
945 950 955 960

20

Ser His Ile Pro Asn Val Asn Ile Tyr Ile
965 970

**This Page is Inserted by IFW Indexing and Scanning
Operations and is not part of the Official Record**

BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of the original documents submitted by the applicant.

Defects in the images include but are not limited to the items checked:

- ☒ **BLACK BORDERS**
- ☐ **IMAGE CUT OFF AT TOP, BOTTOM OR SIDES**
- ☐ **FADED TEXT OR DRAWING**
- ☐ **BLURRED OR ILLEGIBLE TEXT OR DRAWING**
- ☐ **SKEWED/SLANTED IMAGES**
- ☐ **COLOR OR BLACK AND WHITE PHOTOGRAPHS**
- ☐ **GRAY SCALE DOCUMENTS**
- ☐ **LINES OR MARKS ON ORIGINAL DOCUMENT**
- ☐ **REFERENCE(S) OR EXHIBIT(S) SUBMITTED ARE POOR QUALITY**
- ☐ **OTHER:** _____

IMAGES ARE BEST AVAILABLE COPY.

As rescanning these documents will not correct the image problems checked, please do not report these problems to the IFW Image Problem Mailbox.